

ISSN: 1017-060X (Print)



ISSN: 1735-8515 (Online)

**Bulletin of the**  
**Iranian Mathematical Society**

Vol. 42 (2016), No. 5, pp. 1221–1235

**Title:**

**$\mathcal{X}$ -injective and  $\mathcal{X}$ -projective complexes**

**Author(s):**

**T. Özen and E. Yıldırım**

Published by Iranian Mathematical Society  
<http://bims.ims.ir>

## $\mathcal{X}$ -INJECTIVE AND $\mathcal{X}$ -PROJECTIVE COMPLEXES

T. ÖZEN\* AND E. YILDIRIM

(Communicated by Omid Ali S. Karamzadeh)

**ABSTRACT.** Let  $\mathcal{X}$  be a class of  $R$ -modules. In this paper, we investigate  $\mathcal{X}$ -injective (projective) and DG- $\mathcal{X}$ -injective (projective) complexes which are generalizations of injective (projective) and DG-injective (projective) complexes. We prove that some known results can be extended to the class of  $\mathcal{X}$ -injective (projective) and DG- $\mathcal{X}$ -injective (projective) complexes for this general settings.

**Keywords:** Injective (Projective) complex, precover, preenvelope.

**MSC(2010):** Primary: 18G35.

### 1. Introduction

Throughout this paper,  $R$  denotes an associative ring with identity and all modules are unitary. Let  $\mathcal{X}$  be a class of  $R$ -modules. An  $R$ -module  $E$  is called  $\mathcal{X}$ -injective (see [6]), if  $\text{Ext}^1(B/A, E) = 0$  for every module  $B/A \in \mathcal{X}$  or equivalently if  $E$  is injective with respect to every exact sequence  $0 \rightarrow A \rightarrow B \rightarrow B/A \rightarrow 0$  where  $B/A \in \mathcal{X}$ . Dually we can define an  $\mathcal{X}$ -projective module. In Section 2, we define and characterize  $\mathcal{X}$ -injective,  $\mathcal{X}$ -projective, DG- $\mathcal{X}$ -injective and DG- $\mathcal{X}$ -projective complexes which are generalizations of injective, projective, DG-injective and DG-projective complexes, respectively (see [1] and [2]). By [2] we know that  $(\varepsilon, \text{DG-injective})$  is a cotorsion pair. We denote the class of all  $\mathcal{X}$ -complexes, that is, exact complexes with kernel in  $\mathcal{X}$ , by  $\varepsilon_{\mathcal{X}}$ , (in [5] the same class is denoted by  $\tilde{\mathcal{X}}$ ). We prove that if  $\mathcal{X}$  is extension closed, then  $\varepsilon_{\mathcal{X}}^{\perp}({}^{\perp}\varepsilon_{\mathcal{X}}) = \text{DG-}\mathcal{X}\text{-injective (projective)}$  which is proved in [5] when  $(\mathcal{X}, \mathcal{X}^{\perp})$  is a cotorsion pair.

In the last section, we investigate when a complex has an exact  $C(\mathcal{X}\text{-projective (injective)})\text{-precover (preenvelope)}$ . We know that an injective (projective) complex is exact, thus we give some conditions that an  $\mathcal{X}$ -injective (projective) complex is exact and in particular in  $\varepsilon_{\mathcal{X}\text{-injective(projective)}}$ . We prove that if  $\mathcal{X}\text{-injective (projective)} \subseteq \mathcal{X}$  and  $(\mathcal{X}, \mathcal{X}\text{-injective}) ((\mathcal{X}\text{-projective}, \mathcal{X}))$  is a

---

Article electronically published on October 31, 2016.

Received: 12 November 2014, Accepted: 7 August 2015.

\*Corresponding author.

complete cotorsion pair, then every complex has a monic (epic)  $C(\mathcal{X}$ -injective (projective))-preenvelope (precover) in  $\varepsilon_{\mathcal{X}\text{-injective}}$  ( $\varepsilon_{\mathcal{X}\text{-projective}}$ ) and hence  $C(\mathcal{X}$ -injective (projective)) and  $\varepsilon_{\mathcal{X}\text{-injective(projective)}}$  complexes are identical.

Since every complex has an injective and projective resolution, we can compute the right derived functors  $Ext^i(X, Y)$  of  $Hom(-, -)$  where  $Hom(X, Y)$  is the set of all chain maps from  $X$  to  $Y$ .

Moreover  $\mathcal{H}om(X, Y)$  is the complex defined by  $\mathcal{H}om(X, Y)_n = \prod_{p+q=n} (X_{-p}, Y_q)$ .

(See for more details and the other definitions [1, 2, 3, 7]).

## 2. DG- $\mathcal{X}$ -injective and DG- $\mathcal{X}$ -projective complexes

We begin with the following generalized definitions.

**Definition 2.1.** Let  $\mathcal{X}$  be a class of R-modules. A complex  $\mathcal{C} : \dots \rightarrow C^{n-1} \rightarrow C^n \rightarrow C^{n+1} \rightarrow \dots$  is called an  $\mathcal{X}^*$ -(cochain) complex, if  $C^i \in \mathcal{X}$  for all  $i \in \mathbb{Z}$ . A complex  $\mathcal{C} : \dots \rightarrow C_{n+1} \rightarrow C_n \rightarrow C_{n-1} \rightarrow \dots$  is called an  $\mathcal{X}^*$ -(chain) complex, if  $C_i \in \mathcal{X}$  for all  $i \in \mathbb{Z}$ . The class of all  $\mathcal{X}^*$ -complexes is denoted by  $C(\mathcal{X}^*)$ .

**Definition 2.2.** A complex  $\mathcal{C}$  is called an  $\mathcal{X}$ -injective complex, if  $Ext^1(Y/X, C) = 0$  for every complex  $Y/X \in C(\mathcal{X}^*)$ . Equivalently, a complex  $\mathcal{C}$  is an  $\mathcal{X}$ -injective complex if for any exact sequence  $0 \rightarrow X \rightarrow Y \rightarrow Y/X \rightarrow 0$  with a complex  $Y/X \in C(\mathcal{X}^*)$ , the sequence  $Hom(Y, C) \rightarrow Hom(X, C) \rightarrow 0$  is exact.

Dually we can define an  $\mathcal{X}$ -projective complex. A complex  $\mathcal{C}$  is called an  $\mathcal{X}$ -projective complex, if  $Ext^1(C, X) = 0$  for every complex  $X \in C(\mathcal{X}^*)$ , or equivalently a complex  $\mathcal{C}$  is an  $\mathcal{X}$ -projective complex if for any exact sequence  $0 \rightarrow X \rightarrow A \rightarrow B \rightarrow 0$  with a complex  $X \in C(\mathcal{X}^*)$ , the sequence  $Hom(C, A) \rightarrow Hom(C, B) \rightarrow 0$  is exact. We denote the class of all  $\mathcal{X}$ -injective (projective) complexes by  $C(\mathcal{X}\text{-injective (projective)})$ .

**Definition 2.3.** Let  $\varepsilon$  be the class of exact complexes. Then we can define  $\varepsilon_{\mathcal{X}}$  as the class of exact complexes with kernels in  $\mathcal{X}$ .

**Example 2.4.** If  $P$  is an  $\mathcal{X}$ -projective ( $\mathcal{X}$ -injective) module, then  $\bar{P} : \dots \rightarrow 0 \rightarrow P \rightarrow P \rightarrow 0 \rightarrow 0 \rightarrow \dots$  is an  $\mathcal{X}$ -projective ( $\mathcal{X}$ -injective) complex. Moreover any direct sum (product) of  $\mathcal{X}$ -projective ( $\mathcal{X}$ -injective) complexes is again an  $\mathcal{X}$ -projective ( $\mathcal{X}$ -injective) complex. Since  $C(\mathcal{X}\text{-injective (projective)})$  is closed under extensions, every bounded exact complex  $Y : \dots \rightarrow Y^0 \rightarrow \dots \rightarrow Y^n \rightarrow 0 \dots$  with kernels an  $\mathcal{X}$ -injective (projective) module is in  $C(\mathcal{X}\text{-injective (projective)})$ .

Since every right (left) bounded exact complex with kernels  $\mathcal{X}$ -injective (projective) module is an inverse (direct) limit of bounded exact complexes with kernels  $\mathcal{X}$ -injective (projective) module, then every left (right) bounded exact complex with kernels  $\mathcal{X}$ -injective (projective) module is in  $C(\mathcal{X}\text{-injective (projective)})$ .

Moreover if  $\mathcal{X}$ -injective  $\subseteq \mathcal{X}$ , then every  $\varepsilon_{\mathcal{X}}$ -injective complex is a direct sum of  $\mathcal{X}$ -injective complexes, which is the same as injective complexes. Similarly, if  $\mathcal{X}$ -projective  $\subseteq \mathcal{X}$ , then every  $\varepsilon_{\mathcal{X}}$ -projective complexes is a direct sum of  $\mathcal{X}$ -projective complexes. Thus,  $\varepsilon_{\mathcal{X}}$ -injective(projective)  $\subseteq C(\mathcal{X}$ -injective (projective)).

Notice that if  $P$  is an  $\mathcal{X}$ -injective ( $\mathcal{X}$ -projective) module and  $P$  is not in the class  $\mathcal{X}$ , then  $\overline{P}$  is an  $\mathcal{X}$ -injective complex, but not an  $\mathcal{X}^*$ -complex. So an  $\mathcal{X}$ -injective (projective) complex may not be an  $\mathcal{X}^*$ -complex.

**Lemma 2.5.** *Let  $X$  be an  $\mathcal{X}$ -injective complex such that  $\frac{E(X)}{X} \in C(\mathcal{X}^*)$  (or  $\frac{Y}{X} \in C(\mathcal{X}^*)$ ) where  $E(X)$  is an injective envelope of  $X$ . Then  $X = E(X)$  and so it is an injective complex ( $X$  is a direct summand of  $Y$ ).*

*Proof.* We know that every complex has an injective envelope, so  $X$  has an injective envelope  $E(X)$ . Then  $E(X)$  is an injective complex, and so it is exact. We have the following commutative diagram:

$$\begin{array}{ccccc}
 0 & \longrightarrow & X & \xrightarrow{i} & E(X) \\
 & & \downarrow id_x & \searrow \phi & \\
 & & X & & 
 \end{array}$$

such that  $\phi i = id_x$ . Therefore  $X$  is a direct summand of  $E(X)$ . So  $X$  is an injective complex and hence it is exact. Similarly, if  $\frac{Y}{X} \in C(\mathcal{X}^*)$ , then we can prove that  $X$  is a direct summand of  $Y$ . □

**Definition 2.6.** A complex  $I$  is called DG- $\mathcal{X}$ -injective, if each  $I^n$  is  $\mathcal{X}$ -injective and  $\mathcal{H}om(E, I)$  is exact for all  $E \in \varepsilon_{\mathcal{X}}$ . A complex  $I$  is called a DG- $\mathcal{X}$ -projective, if each  $I^n$  is  $\mathcal{X}$ -projective and  $\mathcal{H}om(I, E)$  is exact for all  $E \in \varepsilon_{\mathcal{X}}$ .

**Lemma 2.7.** *Let  $A \xrightarrow{\beta} B \xrightarrow{\theta} C$  be an exact sequence of modules (complexes) where  $\text{Ker}\beta \in \mathcal{X}$  ( $C(\mathcal{X}^*)$ ). Then for all  $\mathcal{X}$ -projective modules (complexes)  $I$ ,  $\text{Hom}(I, A) \longrightarrow \text{Hom}(I, B) \longrightarrow \text{Hom}(I, C)$  is exact.*

*Proof.* By the exact sequence  $0 \longrightarrow \text{Ker}\theta \xrightarrow{i} B \xrightarrow{\theta} C$ ,  $0 \longrightarrow \text{Hom}(I, \text{Ker}\theta) \longrightarrow \text{Hom}(I, B) \longrightarrow \text{Hom}(I, C)$  is exact. We have the following commutative diagram:

$$\begin{array}{ccccc}
 A & \xrightarrow{\beta} & \text{Im}\beta & \longrightarrow & 0 \\
 & \searrow f & \uparrow g & & \\
 & & I & & 
 \end{array}$$

such that  $\beta f = g$ . Since  $I$  is an  $\mathcal{X}$ -projective module (complex) and  $\text{Ker}\beta \in \mathcal{X}$  ( $C(\mathcal{X}^*)$ ),  $\text{Hom}(I, A) \rightarrow \text{Hom}(I, B) \rightarrow \text{Hom}(I, C)$  is exact.  $\square$

Dually we can give the following lemma:

**Lemma 2.8.** *Let  $A \xrightarrow{\beta} B \xrightarrow{\theta} C$  be an exact sequence of modules (complexes) where  $\frac{C}{I_m\theta} \in \mathcal{X}(C(\mathcal{X}^*))$ . Then for all  $\mathcal{X}$ -injective modules (complexes)  $I$ ,  $\text{Hom}(C, I) \rightarrow \text{Hom}(B, I) \rightarrow \text{Hom}(A, I)$  is exact.*

**Example 2.9.** Let  $I = \dots \rightarrow 0 \rightarrow I^0 \rightarrow 0 \rightarrow 0 \rightarrow \dots$  where  $I^0$  is an  $\mathcal{X}$ -injective ( $\mathcal{X}$ -projective) module. Then  $I$  is a DG- $\mathcal{X}$ -injective (DG- $\mathcal{X}$ -projective) complex.

*Proof.* Let  $E : \dots \rightarrow E^{-1} \xrightarrow{d^{-1}} E^0 \xrightarrow{d^0} E^1 \xrightarrow{d^1} E^2 \xrightarrow{d^2} E^3 \rightarrow \dots$  be exact and  $\text{Ker}d^n \in \mathcal{X}$ , then  $\mathcal{H}om(E, I) \cong \dots \text{Hom}(E^2, I^0) \rightarrow \text{Hom}(E^1, I^0) \rightarrow \text{Hom}(E^0, I^0) \dots$ . By Lemma 2.8,  $\mathcal{H}om(E, I)$  is exact.  $\square$

**Lemma 2.10.** *If a complex  $X : \dots \rightarrow X_{n+1} \rightarrow X_n \rightarrow X_{n-1} \rightarrow \dots$  is an  $\mathcal{X}$ -injective ( $\mathcal{X}$ -projective) complex, then for all  $n \in \mathbb{Z}$   $X_n$  is an  $\mathcal{X}$ -injective ( $\mathcal{X}$ -projective) module.*

*Proof.* Let  $0 \rightarrow N \xrightarrow{i} M$  be exact such that  $\frac{M}{N} \in \mathcal{X}$  and  $\alpha : N \rightarrow X_n$  be linear. Form the pushout:

$$\begin{array}{ccc} N & \xrightarrow{i} & M \\ \alpha \downarrow & & \downarrow \gamma_n \\ X_n & \xrightarrow{\theta_n} & \frac{X_n \oplus M}{A} \end{array}$$

where  $A = \{(\alpha(n), -i(n)) : n \in N\}$ . By the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & X_{n+1} & \longrightarrow & X_{n+1} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & X_n & \longrightarrow & \frac{M \oplus X_n}{A} & \longrightarrow & \frac{M}{N} \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & X_{n-1} & \longrightarrow & X_{n-1} & \longrightarrow & 0 \end{array}$$


we have the exact sequence  $0 \rightarrow X \rightarrow T \rightarrow S \rightarrow 0$  where  $T : \dots \rightarrow X_{n+2} \rightarrow X_{n+1} \rightarrow \frac{M \oplus X_n}{A} \rightarrow X_{n-1} \dots$  and  $S : \dots \rightarrow 0 \rightarrow 0 \rightarrow \frac{M}{N} \rightarrow 0 \dots$ . Since  $X$  is an  $\mathcal{X}$ -injective complex,  $Ext^1(S, X) = 0$ , and so  $0 \rightarrow Hom(S, X) \rightarrow Hom(T, X) \rightarrow Hom(X, X) \rightarrow Ext^1(S, X) = 0$ . Therefore there exists  $\beta_n : T_n = \frac{M \oplus X_n}{A} \rightarrow X_n$  such that  $\beta_n \theta_n = 1$ . So

$$\begin{aligned} \beta^n \theta^n(\alpha(n)) &= \alpha(n) \\ \beta^n((\alpha(n), 0) + A) &= \alpha(n) \\ \beta^n((0, i) + A) &= \alpha(n) \\ \beta^n \gamma_n i(n) &= \alpha(n) \end{aligned}$$


and hence  $\beta^n \gamma_n i = \alpha$ . So  $X_n$  is an  $\mathcal{X}$ -injective module. □

The following example shows that if  $X : \dots \rightarrow X_{n+1} \rightarrow X_n \rightarrow X_{n-1} \rightarrow \dots$  is a complex such that  $X_n$  are  $\mathcal{X}$ -injective ( $\mathcal{X}$ -projective) modules for all  $n \in \mathbb{Z}$ , then  $X$  does not need to be an  $\mathcal{X}$ -injective ( $\mathcal{X}$ -projective) complex.

**Example 2.11.** Let  $R \in \mathcal{X}$  be an  $\mathcal{X}$ -injective module and  $f : R \rightarrow R \oplus R$  be a morphism such that  $f(a) = (0, a)$  and  $g : R \oplus R \rightarrow R$  be a morphism such that  $g(a, b) = a$ . Then  $gf = 0$  where  $g \neq 0$ . Consider the following diagrams:



Then we have the diagram:



such that  $g1 = 0$ . But this is impossible. So  $R$  cannot be an  $\mathcal{X}$ -injective complex. Dually, we can give an example for  $\mathcal{X}$ -projectivity.

**Remark 2.12.** There exists a module which is both in  $\mathcal{X}$  and an  $\mathcal{X}$ -injective module. Let  $\mathcal{X}$  be a class of injective modules and  $R$  be an injective module, then  $R$  is both in  $\mathcal{X}$  and an  $\mathcal{X}$ -injective module. Moreover let  $M$  be a flat cotorsion module (see Theorem 5.3.28 in [3] for the existence of such a module) and  $\mathcal{X}$  be a class of flat modules, then  $M$  is both in  $\mathcal{X}$  and an  $\mathcal{X}$ -injective module.

**Lemma 2.13.** *If  $I \in \varepsilon_{\mathcal{X}}^{\perp}$ , then each  $I^n$  is an  $\mathcal{X}$ -injective module for each  $n \in \mathbb{Z}$ .*

*Proof.* Let  $S \subseteq M$  be a submodule of a module  $M$  with  $\frac{M}{S} \in \mathcal{X}$  and  $\alpha : S \rightarrow I_n$  be linear. Form the pushout:

$$\begin{array}{ccc} S & \xrightarrow{i} & M \\ \alpha \downarrow & & \downarrow i_1 \\ I^n & \xrightarrow{i_2} & \frac{I^n \oplus M}{A} = I^n \oplus_S M \end{array}$$

where  $A = \{(\alpha(s), -s) : s \in S\}$ . Thus  $i_2$  is one-to-one the same as  $i$ . Then  $\bar{I} : \dots \rightarrow I^{n-1} \rightarrow I^n \oplus_S M \rightarrow I^{n+1} \rightarrow I^{n+2} \rightarrow \dots$  is a complex.

$$\begin{array}{ccccccc} 0 & \longrightarrow & I^{n-1} & \longrightarrow & I^{n-1} & \longrightarrow & 0 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & I^n & \longrightarrow & I^n \oplus_S M & \longrightarrow & \frac{M}{S} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & I^{n+1} & \longrightarrow & I^{n+1} & \longrightarrow & \frac{M}{S} \longrightarrow 0 \end{array}$$

Therefore, we have an exact sequence  $0 \rightarrow I \rightarrow \bar{I} \rightarrow E \rightarrow 0$  where  $E : \dots \rightarrow \frac{M}{S} \rightarrow \frac{M}{S} \rightarrow 0 \rightarrow 0 \rightarrow 0 \rightarrow \dots$  and so we have an exact sequence  $0 \rightarrow \text{Hom}(E, I) \rightarrow \text{Hom}(\bar{I}, I) \rightarrow \text{Hom}(I, I) \rightarrow \text{Ext}^1(E, I) = 0$  since  $I \in \varepsilon_{\mathcal{X}}^{\perp}$ . This implies that we can find  $\bar{f} : \bar{I} \rightarrow I$  with  $\bar{f}f = 1$ . Therefore, there exists a function  $\bar{f}^n : I^n \oplus_S M \rightarrow I^n$  with  $\bar{f}^n f^n = 1$ . So,

$$\begin{aligned} \bar{f}^n f^n(\alpha(s)) &= \alpha(s) \\ \bar{f}^n((\alpha(s), 0) + A) &= \alpha(s) \\ \bar{f}^n((0, s) + A) &= \alpha(s) \\ \bar{f}^n i_1 i(s) &= \alpha(s) \end{aligned}$$

and hence  $\bar{f}_n i_1 i = \alpha$  and thus each  $I^n \in \mathcal{X}$ -injective.  $\square$

**Lemma 2.14.** *Let  $f : X \rightarrow Y$  be a morphism of complexes. Then the exact sequence  $0 \rightarrow Y \rightarrow M(f) \rightarrow X[1] \rightarrow 0$  associated with the mapping cone  $M(f)$  splits if and only if  $f$  is homotopic to 0.*

*Proof.* The proof follows from [2].  $\square$

**Lemma 2.15.** *Let  $X$  and  $I$  be complexes. If  $Ext^1(X, I[n]) = 0$  for all  $n \in \mathbb{Z}$ , then  $\mathcal{H}om(X, I)$  is exact.*

*Proof.* Since  $Ext^1(X, I[n]) = 0$ , if  $f : X[-1] \rightarrow I[n]$  is a morphism, then  $0 \rightarrow I[n] \rightarrow M(f) \rightarrow X \rightarrow 0$  splits.

By Lemma 2.14,  $f : X[-1] \rightarrow I[n]$  is homotopic to zero for all  $n$ . So  $f^1 : X \rightarrow I[n+1]$  is homotopic to zero for all  $n \in \mathbb{Z}$ . Thus  $\mathcal{H}om(X, I)$  is exact.  $\square$

In [5] the following proposition is proved in the case when  $(\mathcal{X}, \mathcal{X}^\perp)$  is a cotorsion pair.

**Proposition 2.16.** *Let  $\mathcal{X}$  be extension closed. Then  $\varepsilon_{\mathcal{X}}^\perp({}^\perp\varepsilon_{\mathcal{X}}) = DG\text{-}\mathcal{X}$ -injective (projective).*

*Proof.* By Lemma 2.13 and Lemma 2.15 we have that  $\varepsilon_{\mathcal{X}}^\perp({}^\perp\varepsilon_{\mathcal{X}}) \subseteq DG\text{-}\mathcal{X}$ -injective (projective). Let  $I \in DG\text{-}\mathcal{X}$ -injective. Therefore  $\mathcal{H}om(X, I)$  is exact for all  $X \in \varepsilon_{\mathcal{X}}$  and so for all  $n$ ,  $f : X \rightarrow I[n]$  is homotopic to zero. By Lemma 2.14  $A : 0 \rightarrow I[n] \rightarrow M(f) \rightarrow X[1] \rightarrow 0$  is split exact. We know that any exact complex  $B : 0 \rightarrow I[n] \rightarrow Y \rightarrow X[1] \rightarrow 0$  splits at module level since the  $I[n]^m$  are  $\mathcal{X}$ -injective modules and  $X^m \in \mathcal{X}$ . Therefore the exact sequences  $A$  and  $B$  are isomorphic. It is known that  $Ext^1(C, A) = 0$  if and only if every short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  splits. This implies that  $Ext^1(X, I[n]) = 0$  and thus the converse inclusion is proved.  $\square$

If we use Proposition 2.16, then we can give the following example since  $\mathcal{X}$  and  $\varepsilon_{\mathcal{X}}^\perp({}^\perp\varepsilon_{\mathcal{X}})$  are extension closed and every right(left) bounded complex is a direct (inverse) transfinite limit of bounded complexes.

**Example 2.17.** Let  $\mathcal{X}$  be extension closed. Then every  $\mathcal{X}$ -projective (injective) complex is  $DG\text{-}\mathcal{X}$ -projective (injective). Every right (left) bounded complex  $I$  where  $I_i$  is an  $\mathcal{X}$ -projective (injective) module is a  $DG\text{-}\mathcal{X}$ -projective (injective) complex. Moreover  $\varepsilon_{\mathcal{X}\text{-injective(projective)}} \subseteq DG\text{-}\mathcal{X}$ -injective (projective) since the direct (inverse) limit of  $DG\text{-}\mathcal{X}$ -injective complexes is also an inverse (direct) transfinite limit of bounded  $\varepsilon_{\mathcal{X}\text{-injective(projective)}}$  complexes.

$\varepsilon_{\mathcal{X}}$  and  $DG\text{-}\mathcal{X}$ -injective cannot be a cotorsion pair if  $\mathcal{X}$  is extension closed. We have the following theorem:



**Theorem 2.18.** *Let  $\mathcal{X}$  be extension closed and we have enough  $\mathcal{X}$ -object. Then  $(DG\text{-}\mathcal{Y}\text{-projective}, \varepsilon_{\mathcal{Y}})$  is cotorsion pair where  $\mathcal{Y} = \mathcal{X}$ -injective.*

*Proof.* It follows from the proof of Proposition 3.6 in [5] and Proposition 2.16. □

### 3. $C(\mathcal{X}$ -projective)-precovers and $C(\mathcal{X}$ -injective)-preenvelopes

In this section we prove that if a complex has a  $C(\mathcal{X}$ -projective)-precover or  $C(\mathcal{X}$ -injective)-preenvelope in  $C(\mathcal{X}^*)$ , then such precovers or preenvelopes are homotopic. Moreover we investigate when a complex has an exact  $C(\mathcal{X}$ -projective (injective))-precover (preenvelope) and we give some conditions when an  $\mathcal{X}$ -projective (injective) complex is exact and in particular in  $\varepsilon_{\mathcal{X}\text{-projective(injective)}}$ .

**Lemma 3.1.** *i) Let  $f : X \rightarrow Y$  be a chain morphism, let  $X$  be an  $\mathcal{X}^*$  complex and let  $Y$  be an  $\mathcal{X}$ -injective complex. Then  $f$  is homotopic to zero. Moreover if a complex has a  $C(\mathcal{X}$ -injective)-preenvelope in  $C(\mathcal{X}^*)$ , then such preenvelopes are homotopic.*

*ii) Let  $f : X \rightarrow Y$  be a chain homomorphism such that  $Y$  is an  $\mathcal{X}^*$  complex and  $X$  is an  $\mathcal{X}$ -projective complex. Then  $f$  is homotopic to zero. Moreover if a complex has a  $C(\mathcal{X}$ -projective)-precover in  $C(\mathcal{X}^*)$ , then such precovers are homotopic.*

*Proof.* i) Let  $id : X \rightarrow X$ , then we have the following exact sequence:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & X & \xrightarrow{i} & M(id) & \longrightarrow & X[1] \longrightarrow 0 \\
 & & \downarrow f & \nearrow \dots g & & & \\
 & & Y & & & & 
 \end{array}$$

where  $gi = f$ . Let  $i_1^n : X[1]^n \rightarrow M(id)^n$  be a canonical injection and  $s^n : X[1]^{n-1} \rightarrow Y^{n-1}$  such that  $s^n = g^{n-1}i_1^{n-1}$  for all  $n \in \mathbb{Z}$ . Let  $u$  be the differential of the complex  $M(id)$ . Then we have the following diagram

$$\begin{array}{ccccc}
 X^{n-1} \oplus X^{n-2} & \xrightarrow{u^{n-2}} & X^n \oplus X^{n-1} & \xrightarrow{u^{n-1}} & X^{n+1} \oplus X^n \\
 \downarrow g^{n-2} & & \downarrow g^{n-1} & & \downarrow g^n \\
 Y^{n-2} & \xrightarrow{\gamma^{n-2}} & Y^{n-1} & \xrightarrow{\gamma^{n-1}} & Y^n
 \end{array}$$

$s^{n+1}\lambda^n + \gamma^{n-1}s^n = g^n i_1^n \lambda^n + \gamma^{n-1} g^{n-1} i_1^{n-1} = g^n i_1^n \lambda^n + g^n u^{n-1} i_1^{n-1} = g^n (i_1^n \lambda^n + u^{n-1} i_1^{n-1}) = g^n i^n = f^n$ .

ii) Consider  $id : Y \rightarrow Y$  and the exact sequence  $0 \rightarrow Y[-1] \rightarrow M(id)[-1] \rightarrow$

$Y \rightarrow 0$ . Since  $X$  is an  $\mathcal{X}$ -projective complex, we have the following commutative diagram:

$$\begin{array}{ccccc}
 M(id)[-1] & \xrightarrow{\pi} & Y & \longrightarrow & 0 \\
 \uparrow \text{\scriptsize } g & \nearrow \text{\scriptsize } f & & & \\
 \vdots & & & & \\
 \vdots & & & & \\
 X & & & & 
 \end{array}$$

where  $\pi g = f$ . Let  $\pi_1^n : M(id)[-1]^n \rightarrow Y[-1]^n$  be a projection for all  $n \in \mathbb{Z}$ . Then if we take as  $s^n = \pi_1^n g^n$ , then for all  $n \in \mathbb{Z}$ ,  $s^{n+1}\lambda^n + \gamma^{n-1}s^n = f^n$  where  $\lambda$  and  $\gamma$  are boundary maps of the complexes of  $X$  and  $Y$ , respectively. So  $f$  is homotopic to zero.  $\square$

**Proposition 3.2.** *Let  $({}^\perp\mathcal{X}, \mathcal{X})$  ( $(\mathcal{X}, \mathcal{X}^\perp)$ ) be a cotorsion pair. Then every  $\mathcal{X}$ -projective ( $\mathcal{X}$ -injective) complex is exact.*

*Proof.* By [2] we see that every  $\mathcal{X}$ -projective (injective) complex has an exact precover (preenvelope) with kernel (cokernel) in DG-injective (projective). The result follows.  $\square$

**Lemma 3.3.** *Let  $\mathcal{X}$  be extension closed (and  $({}^\perp\mathcal{X}, \mathcal{X})$  be a cotorsion pair). Let every  $R$ -module have an epic  $\mathcal{X}$ -projective-precover with kernel in  $\mathcal{X}$ . Then every bounded complex in  $C(\mathcal{X}^*)$  has an epic exact  $C(\mathcal{X}$ -projective)-precover (which is also in  $\varepsilon_{\mathcal{X}\text{-projective}}$ ) with kernel in  $C(\mathcal{X}^*)$  (which is also in  $DG\text{-}\mathcal{X}\text{-projective-injective} = (\varepsilon_{\mathcal{X}\text{-projective}})^\perp$ ). Thus every bounded  $\mathcal{X}$ -projective complex in  $C(\mathcal{X}^*)$  is exact (which is also in  $\varepsilon_{\mathcal{X}\text{-projective}}$  and every bounded complex in  $C(\mathcal{X}^*)$  has an  $\varepsilon_{\mathcal{X}\text{-projective}}$ -precover).*

*Proof.* Let  $Y(n) : \dots \rightarrow 0 \rightarrow Y^0 \rightarrow Y^1 \rightarrow \dots \rightarrow Y^n \rightarrow 0 \rightarrow \dots \in C(\mathcal{X}^*)$ . We use induction on  $n$ . Let  $n = 0$ , then we have the following commutative diagram:

$$\begin{array}{cccccccc}
 D(0) : \dots & \longrightarrow & 0 & \longrightarrow & P^0 & \xrightarrow{id} & P^0 & \longrightarrow & 0 & \longrightarrow & \dots \\
 & & \downarrow & & \downarrow \text{\scriptsize } f^0 & & \downarrow & & \downarrow & & \\
 Y(0) : \dots & \longrightarrow & 0 & \longrightarrow & Y^0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \dots
 \end{array}$$

where  $P^0 \rightarrow Y^0 \rightarrow 0$  is an  $\mathcal{X}$ -projective-precover in  $\mathcal{X}$  with kernel in  $\mathcal{X}$  since  $\mathcal{X}$  is extension closed,  $D(0)$  is exact and  $Ker(D(0) \rightarrow Y(0)) \in C(\mathcal{X}^*)$ . We

consider the following diagram which is commutative:

$$\begin{array}{ccccccc}
 D(n) : \dots 0 & \longrightarrow & P^0 & \xrightarrow{\lambda^0} & P^0 \oplus P^1 & \xrightarrow{\lambda^1} & \dots P^{n-1} \oplus P^n \xrightarrow{\lambda_1^n} P^n \dots \\
 & & \downarrow f^0 & & \downarrow (0, f^1) & & \downarrow (0, f^n) & \downarrow \\
 Y(n) : \dots 0 & \longrightarrow & Y^0 & \xrightarrow{a^0} & Y^1 & \xrightarrow{a^1} & \dots Y^n \longrightarrow & 0 \dots
 \end{array}$$

where  $\lambda_1^n$  is onto,  $D(n)$  is an exact  $C(\mathcal{X}\text{-projective})$ -precover of  $Y(n)$  such that  $\text{Ker}(D(n) \rightarrow Y(n)) \in \mathcal{C}(\mathcal{X}^*)$  and the  $P^i \rightarrow Y^i \rightarrow 0$  are  $\mathcal{X}$ -projective-precovers in  $\mathcal{X}$  with kernels in  $\mathcal{X}$  for  $1 \leq i \leq n$ . Since  $D(n) \rightarrow Y(n) \rightarrow 0$  and  $\overline{P^{n+1}} \rightarrow \underline{Y^{n+1}} \rightarrow 0$  are  $C(\mathcal{X}\text{-projective})$ -precovers, we have the following commutative diagram:

$$\begin{array}{ccc}
 D(n) & \xrightarrow{s} & \overline{P^{n+1}} \\
 \downarrow & & \downarrow \\
 Y(n) & \longrightarrow & \underline{Y^{n+1}}
 \end{array}$$

Thus we have the diagram:

$$\begin{array}{ccccccc}
 D(n) : \dots 0 & \longrightarrow & P^0 & \xrightarrow{\lambda^0} & P^0 \oplus P^1 \dots & \xrightarrow{\lambda^{n-1}} & P^{n-1} \oplus P^n \xrightarrow{\lambda_1^n} P^n \dots \\
 & & \downarrow & & \downarrow & & \downarrow s^1 & \downarrow s^2 \\
 \overline{P^{n+1}} : \dots 0 & \longrightarrow & 0 & \longrightarrow & 0 \dots & \longrightarrow & P^{n+1} \xrightarrow{1} & P^{n+1} \dots
 \end{array}$$

where  $s^2 \lambda_1^n = s^1$  and  $s^1 \lambda^{n-1} = 0$ . Moreover we see that  $f^{n+1} s^1 = a^n(0, f^n)$  and  $f^{n+1} s^2 = 0$  by the following diagrams:

$$\begin{array}{ccc}
 P^{n-1} \oplus P^n & \xrightarrow{s^1} & P^{n+1} \\
 \downarrow (0, f^n) & & \downarrow f^{n+1} \\
 Y^n & \xrightarrow{a^n} & Y^{n+1}
 \end{array}$$

$$\begin{array}{ccc}
 P^n & \xrightarrow{s^2} & P^{n+1} \\
 \downarrow & & \downarrow f^{n+1} \\
 0 & \longrightarrow & Y^{n+1}
 \end{array}$$

Let  $\lambda^n(x, y) = (\lambda_1^n(x, y), s^1(x, y))$ ,  $\lambda_1^{n+1}(x, y) = s^2(x) - y$ . Then we have the commutative diagram:

$$\begin{array}{ccccccc}
 D(n+1) : \dots & \longrightarrow & P^0 \dots & \longrightarrow & P^{n-1} \oplus P^n & \xrightarrow{\lambda^n} & P^n \oplus P^{n+1} & \xrightarrow{\lambda_1^{n+1}} & P^{n+1} \dots \\
 & & \downarrow f^0 & & \downarrow (0, f^n) & & \downarrow (0, f^{n+1}) & & \downarrow \\
 Y(n+1) : \dots & \longrightarrow & Y^0 \dots & \longrightarrow & Y^n & \xrightarrow{a^n} & Y^{n+1} & \longrightarrow & 0 \dots
 \end{array}$$

where  $\text{Ker}(D(n+1) \rightarrow Y(n+1)) \in C(\mathcal{X}^*)$  and since  $\lambda_1^{n+1}$  is onto,  $\text{Im}(\lambda^n) = \text{Ker}(\lambda_1^{n+1})$ ,  $\text{Im}(\lambda^{n-1}) = \text{Ker}(\lambda^n)$  and  $D(n)$  is exact,  $D(n+1)$  is exact. Therefore,  $Y(n)$  has a  $C(\mathcal{X}\text{-projective})$ -precover.  $\square$

The following corollary is a direct consequence of Lemma 3.3.

**Corollary 3.4.** *i) Let  $\mathcal{X}$  be extension closed. If  $\mathcal{X}\text{-projective} \subseteq \mathcal{X}$  and every  $R$ -module has an epic  $\mathcal{X}$ -projective-precover with kernel in  $\mathcal{X}$  (and  $({}^\perp \mathcal{X}, \mathcal{X})$  is a cotorsion pair), then every bounded complex has an an epic exact  $C(\mathcal{X}\text{-projective})$ -precover (which is also in  $\varepsilon_{\mathcal{X}\text{-projective}}$ ) with kernel in  $C(\mathcal{X}^*)$ . Thus if  $({}^\perp \mathcal{X}, \mathcal{X})$  is a complete cotorsion pair, then  $\varepsilon_{\mathcal{X}\text{-projective}}$  bounded complexes and  $C(\mathcal{X}\text{-projective})$  bounded complexes are identical.*

*ii) If  $({}^\perp \mathcal{X}, \mathcal{X})$  is a complete cotorsion pair, then every bounded complex in  $C(\mathcal{X}^*)$  has an  $\varepsilon_{\mathcal{X}\text{-projective}}$ -precover.*

**Lemma 3.5.** *If  $\mathcal{X}$  is extension closed and every  $R$ -module has a monic  $\mathcal{X}$ -injective-preenvelope with cokernel in  $\mathcal{X}$  (and  $(\mathcal{X}, \mathcal{X}^\perp)$  is a cotorsion pair), then every bounded complex in  $C(\mathcal{X}^*)$  has a monic exact  $C(\mathcal{X}\text{-injective})$ -preenvelope (which is also in  $\varepsilon_{\mathcal{X}\text{-injective}}$ ) with cokernel in  $C(\mathcal{X}^*)$  (which is also in  $DG\text{-}\mathcal{X}\text{-injective-projective} = {}^\perp(\varepsilon_{\mathcal{X}\text{-injective}})$ ).*

*Thus every bounded  $\mathcal{X}$ -injective complex in  $C(\mathcal{X}^*)$  is exact (which is also in  $\varepsilon_{\mathcal{X}\text{-injective}}$  and hence every bounded complex in  $C(\mathcal{X}^*)$  has an  $\varepsilon_{\mathcal{X}\text{-injective}}$ -preenvelope).*

*Proof.* Let  $Y(n) : \dots \rightarrow 0 \rightarrow Y_n \rightarrow Y_{n-1} \rightarrow \dots \rightarrow Y_0 \rightarrow 0 \rightarrow \dots$ . We use induction on  $n$ . Let  $n = 0$ , then we have the following commutative diagram:

$$\begin{array}{ccccccc}
 Y(0) : \dots 0 & \longrightarrow & 0 & \longrightarrow & Y_0 & \longrightarrow & 0 \dots \\
 & & \downarrow & & \downarrow & & \downarrow \\
 E(0) : \dots 0 & \longrightarrow & E_0 & \xrightarrow{id} & E_0 & \longrightarrow & 0 \dots
 \end{array}$$

where  $0 \rightarrow Y_0 \rightarrow E_0$  is a monic preenvelope in  $\mathcal{X}$  with cokernel in  $\mathcal{X}$  and thus  $E(0)$  is an exact preenvelope of  $Y(0)$  with cokernel in  $C(\mathcal{X}^*)$ . We consider the

following diagram which is commutative:

$$\begin{array}{ccccccccccc}
 Y(n) : \dots & \longrightarrow & 0 & \longrightarrow & Y_n & \xrightarrow{a_n} & \dots & \longrightarrow & Y_0 & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow (f_n, 0) & & & & \downarrow f_0 & & \\
 E(n) : \dots & \longrightarrow & E_n & \xrightarrow{\lambda_n^1} & E_n \oplus E_{n-1} & \xrightarrow{\lambda_{n-1}} & \dots & \longrightarrow & E_0 & \longrightarrow & 0
 \end{array}$$

where the  $0 \rightarrow Y_i \rightarrow E_i$  are  $\mathcal{X}$ -injective-preenvelopes in  $\mathcal{X}$  with cokernel in  $\mathcal{X}$  for  $1 \leq i \leq n$ ,  $E(n)$  is exact with cokernel  $(Y(n) \rightarrow E(n)) \in C(\mathcal{X}^*)$ . Since  $0 \rightarrow Y_n \rightarrow E_n$  and  $0 \rightarrow \overline{Y_{n+1}} \rightarrow \overline{E_{n+1}}$  are  $C(\mathcal{X}$ -injective)-preenvelopes, we have the following commutative diagram:

$$\begin{array}{ccc}
 \overline{Y_{n+1}} & \longrightarrow & Y(n) \\
 \downarrow & & \downarrow \\
 \overline{E_{n+1}} & \longrightarrow & E(n)
 \end{array}$$

Then we have the diagram:

$$\begin{array}{ccccccccccc}
 \overline{E_{n+1}} : \dots & \longrightarrow & E_{n+1} & \xrightarrow{1} & E_{n+1} & \longrightarrow & 0 & \longrightarrow & \dots & & \dots \\
 & & \downarrow s_{n+1} & & \downarrow s_n & & \downarrow & & & & \\
 E(n) : \dots & \longrightarrow & E_n & \xrightarrow{\lambda_n^1} & E_n \oplus E_{n-1} & \xrightarrow{\lambda_{n-1}} & \dots & \longrightarrow & \dots & & \dots
 \end{array}$$

where  $s_n = \lambda_n^1 s_{n+1}$  and  $\lambda_{n-1} s_n = 0$ . Moreover we see that  $(f_n, 0)a_{n+1} = s_n f_{n+1}$  and  $\lambda_n^1 s_{n+1} = s_n$  by the following diagrams:

$$\begin{array}{ccc}
 Y_{n+1} & \xrightarrow{a_{n+1}} & Y_n \\
 \downarrow f_{n+1} & & \downarrow (f_n, 0) \\
 E_{n+1} & \xrightarrow{s_n} & E_n \oplus E_{n-1}
 \end{array}$$

$$\begin{array}{ccc}
 E_{n+1} & \xrightarrow{s_{n+1}} & E_n \\
 \downarrow 1 & & \downarrow \lambda_n^1 \\
 E_{n+1} & \xrightarrow{s_n} & E_n \oplus E_{n-1}
 \end{array}$$

Let  $\lambda_{n+1}^1(x) = (x, -s_{n+1}(x))$ ,  $\lambda_n(x, y) = s_n(x) + \lambda_n^1(y)$ . Then we have the following commutative diagram:

$$\begin{array}{ccccccccc}
 Y(n+1) : \dots & 0 & \longrightarrow & Y_{n+1} & \xrightarrow{a_{n+1}} & Y_n \dots & \longrightarrow & Y_0 \dots & \\
 & \downarrow & & \downarrow (f_{n+1}, 0) & & \downarrow (f_n, 0) & & \downarrow f^0 & \\
 E(n+1) : \dots & E_{n+1} & \xrightarrow{\lambda_{n+1}^1} & E_{n+1} \oplus E_n & \xrightarrow{\lambda_n} & E_n \oplus E_{n-1} \dots & \longrightarrow & E_0 \dots & 
 \end{array}$$

where  $E(n+1)$  is exact with cokernel  $(Y(n+1) \rightarrow E(n+1))$  in  $C(\mathcal{X}^*)$ . Therefore,  $Y(n)$  has a  $C(\mathcal{X}$ -injective)-preenvelope.  $\square$

**Corollary 3.6.** *i) Let  $\mathcal{X}$  be extension closed. If  $\mathcal{X}$ -injective  $\subseteq \mathcal{X}$  and every  $R$ -module has a monic  $\mathcal{X}$ -injective-preenvelope with kernel in  $\mathcal{X}$  (and  $(\mathcal{X}, \mathcal{X}^\perp)$  is a cotorsion pair), then every bounded complex has an a monic exact  $C(\mathcal{X}$ -injective)-preenvelope (which is also in  $\varepsilon_{\mathcal{X}\text{-injective}} \subseteq C(\mathcal{X}\text{-injective})$ ) with kernel in  $C(\mathcal{X}^*)$ . Thus  $\varepsilon_{\mathcal{X}\text{-injective}}$  and  $C(\mathcal{X}\text{-injective})$  bounded complexes are identical if  $(\mathcal{X}, \mathcal{X}^\perp)$  is a cotorsion pair .*

*ii) If  $(\mathcal{X}, \mathcal{X}^\perp)$  is a complete cotorsion pair, then every bounded complex in  $C(\mathcal{X}^*)$  has an  $\varepsilon_{\mathcal{X}\text{-injective}}$ -preenvelope.*

We know that the direct (inverse) limit of exact complexes is also exact. Then we can give the following theorem.

**Theorem 3.7.** *Let  $\mathcal{X}$  be closed under extensions. The following are satisfied:*  
*i) If every  $R$ -module has a monic (epic)  $\mathcal{X}$ -injective (projective)-preenvelope (precover) with cokernel (kernel) in  $\mathcal{X}$ , then every left (right) bounded complex in  $C(\mathcal{X}^*)$  has a monic (epic) exact  $C(\mathcal{X}$ -injective (projective))-preenvelope (precover) (which is also in  $\varepsilon_{\mathcal{X}\text{-injective(projective)}}$  if  $(\mathcal{X}, \mathcal{X}^\perp)$  ( $(^\perp\mathcal{X}, \mathcal{X})$ ) is a cotorsion pair). Moreover if  $\mathcal{X}$ -injective (projective)  $\subseteq \mathcal{X}$ , then every left (right) bounded complex has a monic (epic) exact  $C(\mathcal{X}$ -injective (projective))-preenvelope (precover).*

*ii) If every  $R$ -module has a monic (epic)  $\mathcal{X}$ -injective (projective)-preenvelope (precover) with cokernel (kernel) in  $\mathcal{X}$ , then every right (left) bounded complex in  $C(\mathcal{X}^*)$  has a monic (epic) exact  $C(\mathcal{X}$ -injective(projective))-preenvelope (precover).*

*Therefore every right (left) bounded  $\mathcal{X}$ -injective (projective) complex in  $C(\mathcal{X}^*)$  is exact (which is also in  $\varepsilon_{\mathcal{X}\text{-injective(projective)}}$  and every right (left) bounded complexes in  $C(\mathcal{X}^*)$  has an  $\varepsilon_{\mathcal{X}\text{-injective(projective)}}$ -preenvelope (precover) if  $(\mathcal{X}, \mathcal{X}^\perp)$  ( $(^\perp\mathcal{X}, \mathcal{X})$ ) is a cotorsion pair). Moreover if  $\mathcal{X}$ -injective (projective)  $\subseteq \mathcal{X}$ , then every right (left) bounded complex has a monic (epic) exact  $C(\mathcal{X}$ -injective (projective))-preenvelope (precover).*

*Proof.* i) Let  $Y : \dots \rightarrow 0 \rightarrow Y^0 \rightarrow Y^1 \rightarrow \dots$  and  $E(n)$  be a  $C(\mathcal{X}$ -injective)-preenvelope of  $Y(n) : \dots \rightarrow 0 \rightarrow Y^0 \rightarrow \dots \rightarrow Y^n \rightarrow 0 \rightarrow \dots$ . Then  $\varprojlim Y(n) = Y$ . By Lemma 3.5,  $Y(n)$  has a  $C(\mathcal{X}$ -injective)-preenvelope  $E(n)$  such that  $0 \rightarrow$

$Y(n) \rightarrow E(n)$  is exact. Then by Theorem 1.5.13 in [3] and the proof of Lemma 3.5  $0 \rightarrow \varprojlim Y(n) \rightarrow \varprojlim E(n)$  is exact with cokernel  $\varprojlim \frac{E(n)}{Y(n)} \in C(\mathcal{X}^*)$  which is also a direct transfinite limit of  $DG(\mathcal{X})$ -injective-projective complexes. Since  $\text{Ext}^1(\frac{A}{B}, \varprojlim E(n)) = 0$  where  $\frac{A}{B} \in C(\mathcal{X}^*)$  by Lemma 2.3 in [9],  $\varprojlim E(n)$  is an exact  $C(\mathcal{X}$ -injective)-preenvelope of  $Y$ . The other part is also proved similarly using  $C(\mathcal{X}$ -projective) is closed under direct transfinite limits by Theorem 1.2 in [4].

ii) Let  $Y : \dots \rightarrow Y_2 \rightarrow Y_1 \rightarrow Y_0 \rightarrow 0 \rightarrow \dots$  and  $E(n)$  be a  $C(\mathcal{X}$ -injective)-preenvelope of  $Y(n) : \dots \rightarrow 0 \rightarrow Y_n \rightarrow \dots \rightarrow Y_1 \rightarrow Y_0 \rightarrow 0 \rightarrow \dots$ . Then  $\varinjlim Y(n) = Y$ . By Lemma 3.5,  $Y(n)$  has a  $C(\mathcal{X}$ -injective)-preenvelope  $E(n)$  such that  $0 \rightarrow Y(n) \rightarrow E(n)$  is exact. Then by Theorem 1.5.6 in [3]  $0 \rightarrow \varinjlim Y(n) \rightarrow \varinjlim E(n)$  is exact with cokernel  $\varinjlim \frac{E(n)}{Y(n)} \in C(\mathcal{X}^*)$  (which is also in  $DG(\mathcal{X})$ -injective-projective if  $(\mathcal{X}, \mathcal{X}^\perp)$  is a cotorsion pair). Since  $\varinjlim E(n)$  is also an inverse transfinite limit of some bounded  $\mathcal{X}$ -injective complexes,  $\text{Ext}^1(\frac{A}{B}, \varinjlim E(n)) = 0$  where  $\frac{A}{B} \in C(\mathcal{X}^*)$ . So  $\varinjlim E(n)$  is an exact  $C(\mathcal{X}$ -injective)-preenvelope of  $Y$ .  $\square$

**Corollary 3.8.** *i) Let  $\mathcal{X}$  be closed under extensions. If every  $R$ -module has a monic (epic)  $\mathcal{X}$ -injective (projective)-preenvelope (precover) with cokernel (kernel) in  $\mathcal{X}$ , then every complex in  $C(\mathcal{X}^*)$  has a monic (epic) exact  $C(\mathcal{X}$ -injective (projective))-preenvelope (precover) (which is also in  $\varepsilon_{\mathcal{X}$ -injective(projective)} if  $(\mathcal{X}, \mathcal{X}^\perp)$  ( $(^\perp \mathcal{X}, \mathcal{X})$ ) is a cotorsion pair). Moreover if  $\mathcal{X}$ -injective (projective)  $\subseteq \mathcal{X}$ , then every complex has a monic (epic) exact  $C(\mathcal{X}$ -injective (projective))-preenvelope (precover) (which is in  $\varepsilon_{\mathcal{X}$ -injective ( $\varepsilon_{\mathcal{X}$ -projective)} if  $(\mathcal{X}, \mathcal{X}^\perp)$  ( $(^\perp \mathcal{X}, \mathcal{X})$ ) is a cotorsion pair, thus  $\varepsilon_{\mathcal{X}$ -injective(projective)} and  $C(\mathcal{X}$ -injective (projective)) complexes are identical).*

ii) *If  $(\mathcal{X}, \mathcal{X}^\perp)$  is a complete cotorsion pair, then every complex in  $C(\mathcal{X}^*)$  has a monic  $\varepsilon_{\mathcal{X}$ -injective-preenvelope.*

**Example 3.9.** Let  $\mathcal{X}$  be a class of  $R$ -modules closed under quotients, extensions and direct sums (for the existence of such classes, if  $\mathcal{X}$  is a class of injective modules on a hereditary noetherian ring which is constructed in [8], then  $\mathcal{X}$  is closed under quotients, extensions and direct limits and moreover if  $\mathcal{X}$  is the class of min-injective modules and simple ideals of ring  $R$  are projective, then it is closed under quotients, extensions and direct sums). If  $A$  and  $B$  are in  $\mathcal{X}$  such that  $\phi : A \rightarrow B$  is a homomorphism, then by Theorem 2.10 in [6], we have monic  $\mathcal{X}$ -injective-preenvelopes such that  $f : A \rightarrow E_A$  and  $g : B \rightarrow E_B$  with cokernels in  $\mathcal{X}$ . Then there exists a homomorphism  $s : E_A \rightarrow E_B$  such that  $g\phi = sf$ . Using Lemma 3.5 we can determine an exact  $C(\mathcal{X}$ -injective)-preenvelope  $E(1)$  of complex  $Y(1)$  as follows:

$$\begin{array}{ccccccccc}
Y(1) : \dots 0 & \longrightarrow & 0 & \longrightarrow & A & \xrightarrow{\phi} & B & \longrightarrow & 0 \dots \\
\downarrow & & \downarrow & & \downarrow (f,0) & & \downarrow g & & \\
E(1) : \dots 0 & \longrightarrow & E_A & \xrightarrow{\alpha} & E_A \oplus E_B & \xrightarrow{\beta} & E_B & \longrightarrow & 0 \dots
\end{array}$$

where  $\alpha(x) = (x, -s(x))$  and  $\beta(x, y) = s(x) + y$ . Then every complex in  $C(\mathcal{X}^*)$  has a monic exact C(X-injective)-preenvelope by Corollary 3.8.

#### REFERENCES

- [1] L. L. Avramov and H. B. Foxby, Homological dimensions of unbounded complexes, *J. Pure Appl. Algebra* **71** (1991), no. 2-3, 129–155.
- [2] E. E. Enochs, O. M. G. Jenda and J. Xu, Orthogonality in the category of complexes, *Math. J. Okayama Univ.* **38** (1996) 25–46.
- [3] E. E. Enochs and O. M. G. Jenda, *Relative Homological Algebra*, de Gruyter Ex. Math., 30, Walter de Gruyter & Co., Berlin, 2000.
- [4] P. Eklof, Homological algebra and set theory, *Trans. Amer. Math. Soc.* **227** (1977) 207–225.
- [5] J. Gillespie, The flat model structure on  $\text{Ch}(\mathbb{R})$ , *Trans. Amer. Math. Soc.* **356** (2004), no. 8, 3369–3390.
- [6] L. X. Mao and N. Q. Ding, L-injective hulls of modules, *Bull. Aust. Math. Soc.* **74** (2006), no. 1, 37–44.
- [7] Joseph J. Rotman, *An Introduction to Homological Algebra*, Springer, New York, 2009.
- [8] J. T. Stafford and R. B. Warfield, Construction of Hereditary Noetherian Rings, *Proc. Lond. Math. Soc. (3)* **51** (1985), no. 1, 1–20.
- [9] J. Trlifaj, Ext and inverse limits, *Illinois J. Math.* **47** (2003), no. 1-2, 529–538.

(Tahire Özen) DEPARTMENT OF MATHEMATICS, ABANT İZZET BAYSAL UNIVERSITY, GÖLKÖY KAMPÜSÜ BOLU, TURKEY.

*E-mail address:* ozen.t@ibu.edu.tr

(Emine Yıldırım) DEPARTMENT OF MATHEMATICS, ABANT İZZET BAYSAL UNIVERSITY, GÖLKÖY KAMPÜSÜ BOLU, TURKEY.

*E-mail address:* emineyildirim@gmail.com