

THE STRUCTURE AND AMENABILITY OF ℓ^p -MUNN ALGEBRAS

S. NASERI* AND H. SAMEA

Communicated by Gholamhossein Esslamzadeh

ABSTRACT. We introduce the notion of $\mathcal{LM}_I^p(\mathcal{A})$, where \mathcal{A} is a Banach space, I is an index set and $1 \leq p < \infty$. We find necessary and sufficient conditions for which $\mathcal{LM}_I^p(\mathcal{A})$ is a Banach algebra and investigate amenability of this Banach algebra. Applications to $\ell^p(S)$ ($1 \leq p < \infty$), where S is a Brandt semigroup, are also given.

1. Introduction

Some properties of ℓ^1 -Munn algebras were investigated by Esslamzadeh [3], where the author introduced the notion and used them as a tool for studying certain semigroup algebras. For more information, see [2-4]. Our aim here is to introduce and investigate the properties of ℓ^p -Munn algebras. It enables us to study some properties of ℓ^p -spaces on Brandt semigroups. This paper is organized as follows. Our notations are introduced in the present section. In section 2, we introduce and investigate the structure of $\mathcal{LM}_I^p(\mathcal{A})$, for the Banach space \mathcal{A} , the index set I , and $1 \leq p < \infty$. The Banach space $\mathcal{LM}_I^p(\mathcal{A})$ is the vector space of all $I \times I$ -matrices A over \mathcal{A} such that $\|A\|_p = \left(\sum_{i,j \in I} \|A_{ij}\|^p \right)^{\frac{1}{p}} < \infty$. We find

MSC(2010): Primary: 43A07.

Keywords: Banach algebra, amenability, semigroup.

Received: 30 September 2008. Accepted: 30 July 2009.

*Corresponding author

© 2010 Iranian Mathematical Society.

necessary and sufficient conditions for which $\mathcal{LM}_I^p(\mathcal{A})$ is a Banach algebra. We prove that if \mathcal{A} is a unital Banach algebra, then $\mathcal{LM}_I^p(\mathcal{A})$ is a Banach algebra if and only if $1 \leq p \leq 2$. Moreover, it is proved that if G is a group and S is a Brandt semigroup over G with index set I , then the Banach space $\ell^p(S)/\mathbb{C}\delta_0$ is isometrically isomorphic with $\mathcal{LM}_I^p(\ell^p(G))$. Moreover, if G is a finite group, and I is finite, then $(\ell^p(S), *)/\mathbb{C}\delta_0$ is isometrically isomorphic with $\mathcal{LM}_I^p(\ell^p(G), *)$. Finally, in Section 3 we study the amenability of the Banach algebra $\mathcal{LM}_I^p(\mathcal{A})$ ($1 \leq p \leq 2$) over a Banach algebra \mathcal{A} with a unit. We prove that $\mathcal{LM}_I^p(\mathcal{A})$ ($1 \leq p \leq 2$) is amenable, if and only if \mathcal{A} is amenable, and I is finite.

The following are some of the notations which we use here.

Let \mathcal{A} be a Banach algebra. If \mathcal{A} admits a unit $e_{\mathcal{A}}$ ($ae_{\mathcal{A}} = e_{\mathcal{A}}a = a$, for all $a \in \mathcal{A}$) and $\|e_{\mathcal{A}}\| = 1$, we say that \mathcal{A} is a *unital normed algebra*. For a Banach algebra \mathcal{A} , an \mathcal{A} -bimodule will always refer to a *Banach \mathcal{A} -bimodule* X ; that is, a Banach space which is algebraically an \mathcal{A} -bimodule, and for which there is a constant $C_X \geq 0$ such that for $a \in \mathcal{A}$, $x \in X$, $\|a.x\| \leq C_X\|a\|\|x\|$, $\|x.a\| \leq C_X\|x\|\|a\|$. A *derivation* $D : \mathcal{A} \rightarrow X$ is a linear map, always taken to be continuous, satisfying $D(ab) = D(a).b + a.D(b)$, for $a, b \in \mathcal{A}$. For every $x \in X$, we define ad_x by $ad_x(a) = a.x - x.a$, for $a \in \mathcal{A}$. Note that ad_x is a derivation which is called an *inner derivation*. A Banach algebra \mathcal{A} is called *amenable* if and only if, for any \mathcal{A} -bimodule X , every derivation $D : \mathcal{A} \rightarrow X^*$ is inner.

2. The structure of the Banach space $\mathcal{LM}_I^p(\mathcal{A})$ ($1 \leq p < \infty$) over a Banach algebra \mathcal{A}

Definition 2.1. Let \mathcal{A} be a Banach space, $1 \leq p < \infty$, and I be an arbitrary index set, and let $\mathcal{LM}_I^p(\mathcal{A})$ be the vector space of all $I \times I$ -matrices A over \mathcal{A} such that

$$\|A\|_p = \left(\sum_{i,j \in I} \|A_{ij}\|^p \right)^{\frac{1}{p}} < \infty.$$

Then, it is easy to check that $\mathcal{LM}_I^p(\mathcal{A})$ with scalar multiplication, matrix addition, and the norm $\|\cdot\|_p$ is a Banach space. This Banach space is called *ℓ^p -Munn Banach space* over \mathcal{A} . If $\mathcal{LM}_I^p(\mathcal{A})$ is a Banach algebra, then $\mathcal{LM}_I^p(\mathcal{A})$ is called the *ℓ^p -Munn Banach algebra* over \mathcal{A} with index set I .

The space $\mathcal{LM}_I^1(\mathcal{A})$ over a unital Banach algebra \mathcal{A} is called the ℓ^1 -Munn Banach algebra over \mathcal{A} with index set I (see [3]). If I is finite, then $\|\cdot\|_{\mathcal{LM}_I^1(\mathbb{C})}$ is called *Frobenius norm*.

A Brandt semigroup S over a group G with index set I consists of all canonical $I \times I$ matrix units over $G \cup \{0\}$ and a zero matrix 0 . Note that an $I \times I$ matrix whose entries are zero except one, is called a canonical matrix unit.

Let G be a group and S be a Brandt semigroup over G . For $f \in \ell^p(S)$, and $i, j \in I$, define $f_{ij} : G \rightarrow \mathbb{C}$ by

$$f_{ij}(g) = f((g)_{ij}),$$

where $(g)_{ij}$ is the matrix with (k, l) -entry equal to g if $(k, l) = (i, j)$ and 0 if $(k, l) \neq (i, j)$. Since for every $i, j \in I$,

$$\sum_{g \in G} |f((g)_{ij})|^p \leq \sum_{s \in S} |f(s)|^p < \infty,$$

then we have $f_{ij} \in \ell^p(G)$. It is clear that if $A = [f_{ij}]$, then $A \in \mathcal{LM}_I^p(\ell^p(G))$. Now, as in Proposition 5.6 of [3], let

$$\Phi : \ell^p(S) \rightarrow \mathcal{LM}_I^p(\ell^p(G)) : f \mapsto [f_{ij}].$$

It is clear that Φ is a well-defined linear map with $\|\Phi\| \leq 1$. Suppose $A \in \mathcal{LM}_I^p(\ell^p(G))$ and $A = [f_{ij}]$. Define $f : S \rightarrow \mathbb{C}$ by $f(0) = 0$ and $f((g)_{ij}) = f_{ij}(g)$, for $g \in G$ and $i, j \in I$. Since

$$\sum_{s \in S} |f(s)|^p = \sum_{i, j \in I} \sum_{g \in G} |f((g)_{ij})|^p = \sum_{i, j \in I} \|f_{ij}\|_p^p < \infty,$$

Then $f \in \ell^p(S)$. Clearly $\Phi(f) = A$. Hence, Φ is onto. Therefore, there is an isometrical isomorphism from $\ell^p(S)/\mathbb{C}\delta_0$ onto $\mathcal{LM}_I^p(\ell^p(G))$. Thus, we have the following result.

Proposition 2.2. *Let G be a group and S be a Brandt semigroup over G with the index set I . Then, the Banach space $\ell^p(S)/\mathbb{C}\delta_0$ is isometrically isomorphic with $\mathcal{LM}_I^p(\ell^p(G))$.*

For the rest of the paper, we assume that \mathcal{A} is a Banach algebra.

Theorem 2.3. *Let $1 \leq p \leq 2$. The Banach space $\mathcal{LM}_I^p(\mathcal{A})$ with matrix multiplication and norm $\|\cdot\|_p$ is a Banach algebra.*

Proof. Let $A, B \in \mathcal{LM}_I^p(\mathcal{A})$, and $i, j \in I$. Since $1 \leq p \leq 2$, then for q with $\frac{1}{p} + \frac{1}{q} = 1$, we have $q \geq 2 \geq p$. Hence, $\ell^p(I) \subseteq \ell^q(I)$ and

$\|f\|_q^p \leq \|f\|_p^p$ ($f \in \ell^p(I)$). We denote the function $f : I \rightarrow \mathbb{C}$, by $(f(i))_i$. Now, we have

$$\begin{aligned} \left(\sum_{k \in I} \|A_{ik}\| \|B_{kj}\| \right)^p &= \|(\|A_{ik}\|)_k (\|B_{kj}\|)_k\|_1^p \\ &\leq \|(\|A_{ik}\|)_k\|_p^p \|(\|B_{kj}\|)_k\|_q^p \\ &\leq \|(\|A_{ik}\|)_k\|_p^p \|(\|B_{kj}\|)_k\|_p^p \\ &= \left(\sum_{k \in I} \|A_{ik}\|^p \right) \left(\sum_{l \in I} \|B_{lj}\|^p \right). \end{aligned}$$

Therefore,

$$\begin{aligned} \|AB\|_p^p &= \sum_{i,j \in I} \left\| \sum_{k \in I} A_{ik} B_{kj} \right\|^p \\ &\leq \sum_{i,j \in I} \left(\sum_{k \in I} \|A_{ik}\| \|B_{kj}\| \right)^p \\ &\leq \sum_{i,j \in I} \left(\sum_{k \in I} \|A_{ik}\|^p \right) \left(\sum_{l \in I} \|B_{lj}\|^p \right) \\ &= \left(\sum_{i,k \in I} \|A_{ik}\|^p \right) \left(\sum_{j,l \in I} \|B_{lj}\|^p \right) \\ &= \|A\|_p^p \|B\|_p^p. \end{aligned}$$

Hence, $\|AB\|_p \leq \|A\|_p \|B\|_p$. This shows that $\|\cdot\|_p$ is an algebra norm. Hence, $\mathcal{LM}_I^p(\mathcal{A})$ is a Banach algebra. \square

Example 2.4. Let \mathcal{A} be a non-zero Banach space. Define

$$a.b = 0 \quad (a, b \in \mathcal{A}).$$

With this multiplication \mathcal{A} is a Banach algebra. Now, let I be an arbitrary set and $1 \leq p < \infty$. Then for each $A, B \in \mathcal{LM}_I^p(\mathcal{A})$, $AB = 0$. Hence, $\mathcal{LM}_I^p(\mathcal{A})$ is a Banach algebra.

Proposition 2.5. *Let I be an infinite set and \mathcal{A} be a Banach algebra such that $\mathcal{A}^2 \neq 0$. Then, for each $2 < p < \infty$, $\mathcal{LM}_I^p(\mathcal{A})$ is not an algebra.*

Proof. Since $\mathcal{A}^2 \neq 0$, then there exist $a, b \in \mathcal{A}$ such that $ab \neq 0$. Let $\{i_n\}_{n \in \mathbb{N}}$ be an infinite subset of distinct elements of I . Define the $I \times I$ -matrix A over \mathcal{A} by $A_{i_1 i_n} = \frac{1}{\sqrt{n}}a$ ($n \in \mathbb{N}$) and $A_{ij} = 0$, for other $i, j \in I$. Also, define the $I \times I$ -matrix B over \mathcal{A} by $B_{i_n i_1} = \frac{1}{\sqrt{n}}b$ ($n \in \mathbb{N}$) and $B_{ij} = 0$, for other $i, j \in I$. It is easy to see that $A, B \in \mathcal{LM}_I^p(\mathcal{A})$. But AB is not even well defined, since

$$(AB)_{i_1 i_1} = \sum_{n \in \mathbb{N}} A_{i_1 i_n} B_{i_n i_1} = \left(\sum_{n \in \mathbb{N}} \frac{1}{n} \right) ab.$$

□

Proposition 2.6. *Let I be a set with at least two elements, and \mathcal{A} be a unital Banach algebra. Then, $\mathcal{LM}_I^p(\mathcal{A})$ is a Banach algebra if and only if $1 \leq p \leq 2$.*

Proof. By Theorem 2.3, if $1 \leq p \leq 2$, then $\mathcal{LM}_I^p(\mathcal{A})$ is a Banach algebra. By Proposition 2.5, if I is infinite, and $2 < p < \infty$, then $\mathcal{LM}_I^p(\mathcal{A})$ is not a Banach algebra. Now, suppose I is finite. Let $i_1, i_2 \in I$ and $i_1 \neq i_2$. Define the $I \times I$ -matrix A over \mathcal{A} by $A_{i_1 i_1} = A_{i_1 i_2} = e_{\mathcal{A}}$ and $A_{ij} = 0$, for other $i, j \in I$. Also, define the $I \times I$ -matrix B over \mathcal{A} by $B_{i_1 i_1} = B_{i_2 i_1} = e_{\mathcal{A}}$ and $B_{ij} = 0$, for other $i, j \in I$. Then,

$$\|AB\|_p = 2 > 2^{\frac{2}{p}} = 2^{\frac{1}{p}} 2^{\frac{1}{p}} = \|A\|_p \|B\|_p,$$

and so $\|\cdot\|_p$ is not an algebra norm. Hence, $\mathcal{LM}_I^p(\mathcal{A})$ is not a Banach algebra. □

Remark 2.7. (a) Let I be finite and \mathcal{A} be a Banach algebra with the unit $e_{\mathcal{A}}$. Suppose $\text{Card}(I) = m$. If $(A_1, \dots, A_m) \in \mathcal{A}^m$, $1 \leq p < \infty$ and $\frac{1}{p} + \frac{1}{q} = 1$, then

$$\|(A_i)_i\|_1 = \|(A_i e_{\mathcal{A}})_i\|_1 \leq \|(A_i)_{i \in I}\|_p \|(e_{\mathcal{A}})_{i \in I}\|_q = m^{\frac{1}{q}} \|e_{\mathcal{A}}\| \|(A_i)_{i \in I}\|_p.$$

Thus, for arbitrary $I \times I$ matrices A, B on \mathcal{A} ,

$$\begin{aligned} \|AB\|_p^p &= \sum_{i,j \in I} \left\| \sum_{k \in I} A_{ik} B_{kj} \right\|^p \leq \sum_{i,j \in I} \left(\sum_{k \in I} \|A_{ik}\| \|B_{kj}\| \right)^p \\ &= \sum_{i,j \in I} \left\| (\|A_{ik}\|)_k (\|B_{kj}\|)_k \right\|_1^p \leq \sum_{i,j \in I} \left\| (\|A_{ik}\|)_k \right\|_1^p \left\| (\|B_{lj}\|)_l \right\|_1^p \\ &\leq m^{\frac{2p}{q}} \|e_{\mathcal{A}}\|^{2p} \sum_{i,j \in I} \left\| (\|A_{ik}\|)_k \right\|_p^p \left\| (\|B_{lj}\|)_l \right\|_p^p = m^{\frac{2p}{q}} \|e_{\mathcal{A}}\|^{2p} \|A\|_p^p \\ &\quad \|B\|_p^p. \end{aligned}$$

Hence, $\|AB\|_p \leq m^{\frac{2}{q}} \|e_{\mathcal{A}}\|^2 \|A\|_p \|B\|_p$.

(b) Let I be finite and \mathcal{A} be a Banach algebra with the unit $e_{\mathcal{A}}$. Suppose $\text{Card}(I) = m$. By (a), it is easy to see that $(\mathcal{L}M_I^p(\mathcal{A}), m^{\frac{2}{q}} \|e_{\mathcal{A}}\|^2 \|\cdot\|_p)$ is a Banach algebra.

(c) Let I be finite and \mathcal{A} be a Banach algebra with the unit $e_{\mathcal{A}}$. Suppose $\text{Card}(I) = m$. Define the norm $\|\cdot\|$ on \mathcal{A} by $\|a\| = C\|a\|$ ($a \in \mathcal{A}$), where $C \geq m^{\frac{2}{q}} \|e_{\mathcal{A}}\|^2$. Let $\tilde{\mathcal{A}}$ denote the algebra \mathcal{A} with the norm $\|\cdot\|$ and A be an $I \times I$ -matrix over $\tilde{\mathcal{A}}$. Then,

$$\|A\|_{\mathcal{L}M_I^p(\tilde{\mathcal{A}})} = C\|A\|_{\mathcal{L}M_I^p(\mathcal{A})}.$$

From this equality and (a), for each $A, B \in \mathcal{L}M_I^p(\tilde{\mathcal{A}})$ we obtain,

$$\begin{aligned} \|AB\|_{\mathcal{L}M_I^p(\tilde{\mathcal{A}})} &= C\|AB\|_{\mathcal{L}M_I^p(\mathcal{A})} \leq C m^{\frac{2}{q}} \|e_{\mathcal{A}}\|^2 \|A\|_{\mathcal{L}M_I^p(\mathcal{A})} \|B\|_{\mathcal{L}M_I^p(\mathcal{A})} \\ &= \frac{m^{\frac{2}{q}} \|e_{\mathcal{A}}\|^2}{C} \|A\|_{\mathcal{L}M_I^p(\tilde{\mathcal{A}})} \|B\|_{\mathcal{L}M_I^p(\tilde{\mathcal{A}})} \leq \|A\|_{\mathcal{L}M_I^p(\tilde{\mathcal{A}})} \\ &\quad \|B\|_{\mathcal{L}M_I^p(\tilde{\mathcal{A}})}. \end{aligned}$$

Therefore, $\mathcal{L}M_I^p(\tilde{\mathcal{A}})$ is a Banach algebra.

Example 2.8. The algebra $\mathcal{A} = \mathbb{C}$ with the norm $\|A\| = 3|A|$ ($A \in \mathcal{A}$) is a Banach algebra with a the unit that is not unital (since $\|1\| = 3 \neq 1$). Then, by notations of Remark 2.7, $\mathcal{A} = \mathbb{C}$ with $C = 3$. Let $I = \{1, 2\}$. Since $C \geq 2^{2\frac{2}{3}}|1|$, then by remark 2.7, $\mathcal{L}M_I^3(\mathcal{A})$ is a Banach algebra. This example shows that we can not replace the condition “ \mathcal{A} is unital” by “ \mathcal{A} has a unit” in the Proposition 2.6.

Proposition 2.9. *Let G be a finite group with $\text{Card}(G) = m$, $1 < p < \infty$, and S be a Brandt semigroup over G with the index set I . Then, $\ell^p(S)$ is closed under convolution if and only if I is finite. Moreover, if I is finite, then there exists a constant C such that $\ell^p(S)$ with the product*

$$\delta_s * \delta_t = \delta_{st} \quad (s, t \in S),$$

and the norm $C\|\cdot\|_p$ defines a Banach algebra. Also, $\ell^p(G)$ with the norm $C\|\cdot\|_p$ is a Banach algebra under convolution, and $\ell^p(S)/\mathbb{C}\delta_0$ is an isometric Banach algebra-isomorphic with $\mathcal{LM}_I^p(\ell^p(G))$.

Proof. Suppose I is infinite. Let $\{i_n\}_{n \in \mathbb{N}}$ be an infinite subset of distinct elements of I . Let $f = \sum_{n=1}^{\infty} \frac{1}{n} \delta_{(e)_{i_1 i_n}}$, and $g = \sum_{n=1}^{\infty} \frac{1}{n} \delta_{(e)_{i_n i_1}}$. Clearly, $f, g \in \ell^p(S)$. But

$$f * g(0) = \sum_{m, n \in \mathbb{N}, m \neq n} \frac{1}{mn} = \infty.$$

Hence, $\ell^p(S)$ is not closed under convolution.

Suppose I is finite with $\text{Card}(I) = l$. It is easy to see that the Banach space $\ell^p(G)$ with the norm $\|\cdot\|_{\ell^p(G)} = m^{1-\frac{1}{p}} \|\cdot\|_p$ and the product

$$\delta_x * \delta_y = \delta_{xy} \quad (x, y \in G),$$

defines a convolution Banach algebra. Note that δ_e (e is the unit of G) is the unit of $\ell^p(G)$ with $\|\delta_e\|_{\ell^p(G)} = m^{1-\frac{1}{p}}$.

By Theorem 2.3, for $p \leq 2$, $\mathcal{LM}_I^p(\ell^p(G))$ defines a Banach algebra. In this case, let $C = m^{1-\frac{1}{p}}$. For $p > 2$, by Remark 2.7(c),

$$\|\cdot\| = l^{\frac{2}{q}} \|\delta_e\|_{\ell^p(G)}^2 \|\cdot\|_{\ell^p(G)} = l^{\frac{2}{q}} \|\delta_e\|_{\ell^p(G)}^2 m^{1-\frac{1}{p}} \|\cdot\|_p = l^{\frac{2}{q}} m^{3(1-\frac{1}{p})} \|\cdot\|_p.$$

Thus, $\mathcal{LM}_I^p(\ell^p(G))$ defines a Banach algebra. In this case, let $C = l^{\frac{2}{q}} m^{3(1-\frac{1}{p})}$. It is easy to see that for the mapping

$$\Phi : (\ell^p(S), C\|\cdot\|_p) \longrightarrow \mathcal{LM}_I^p((\ell^p(G), C\|\cdot\|_p)) : f \mapsto [f_{ij}],$$

$\Phi(\delta_s * \delta_t) = \Phi(\delta_s)\Phi(\delta_t)$. Hence, Φ is an algebra homomorphism. Therefore, by Proposition 2.2, the Banach algebra $(\ell^p(S), C\|\cdot\|_p)/\mathbb{C}\delta_0$ is isometrically algebra isomorphic with $\mathcal{LM}_I^p((\ell^p(G), C\|\cdot\|_p))$. \square

Remark 2.10. By Proposition 5.6 of [3], for a Brandt semigroup S over a group G with an index set I , $\ell^1(S)/\mathbb{C}\delta_0$ is isometrically algebra isomorphic with $\mathcal{LM}_I(\ell^1(G))$.

3. Amenability of the Banach algebra $\mathcal{LM}_I^p(\mathcal{A})$ ($1 \leq p \leq 2$) over a Banach algebra \mathcal{A} with unit

Throughout this section, we suppose \mathcal{A} has a unit which we denote by $e_{\mathcal{A}}$.

Lemma 3.1. *Let \mathcal{A} be a Banach algebra with unit $e_{\mathcal{A}}$, and $1 \leq p \leq 2$. The following conditions are equivalent:*

- (1) $\mathcal{LM}_I^p(\mathcal{A})$ has a bounded approximate identity.
- (2) I is finite.

Proof. (1) \Rightarrow (2) Suppose on the contrary that I is infinite and $(E_{\alpha})_{\alpha}$ is an approximate identity for $\mathcal{LM}_I^p(\mathcal{A})$. For every finite subset F of I , define E_F by $(E_F)_{ii} = e_{\mathcal{A}}$ if $i \in F$, $(E_F)_{ii} = 0$ if $i \in I - F$ and $(E_F)_{ij} = 0$ if $i \neq j$. Then,

$$\begin{aligned} (\text{Card}F)^{\frac{1}{p}} &= \left(\sum_{i \in F} \|e_{\mathcal{A}}\|^p \right)^{\frac{1}{p}} = \|E_F\|_p \\ &= \lim_{\alpha} \|E_F E_{\alpha}\|_p = \lim_{\alpha} \left(\sum_{i \in F, j \in I} \|(E_{\alpha})_{ij}\|^p \right)^{\frac{1}{p}} \\ &\leq \liminf \|E_{\alpha}\|_p. \end{aligned}$$

Therefore, $\lim_{\alpha} \|E_{\alpha}\|_p = \infty$. Thus, $\mathcal{LM}_I^p(\mathcal{A})$ does not have a bounded approximate identity.

(2) \Rightarrow (1) Suppose I is finite. Then, it is clear that E_I is a unit for $\mathcal{LM}_I^p(\mathcal{A})$. \square

Theorem 3.2. *Let \mathcal{A} be a Banach algebra with a unit and $1 \leq p \leq 2$. The following conditions are equivalent:*

- (i) $\mathcal{LM}_I^p(\mathcal{A})$ is amenable.
- (ii) \mathcal{A} is amenable and I is finite.

Proof. (i) \Rightarrow (ii): Since $\mathcal{LM}_I^p(\mathcal{A})$ is amenable, then by Proposition(2.2.1) of [5], $\mathcal{LM}_I^p(\mathcal{A})$ has a bounded approximate identity and by Lemma 3.1, I is a finite set. By Corollary 4 of Section 4 of [1], there exists an equivalent norm $\|\cdot\|$ on \mathcal{A} such that $\tilde{\mathcal{A}} = (\mathcal{A}, \|\cdot\|)$ is unital. Since I is finite, then the identity map $A \mapsto A$; $\mathcal{LM}_I^p(\mathcal{A}) \longrightarrow \mathcal{LM}_I(\tilde{\mathcal{A}})$ is continuous. Indeed, if $c\|a\| \leq \|a\| \leq C\|a\|$ ($a \in \mathcal{A}$), then by Remark

2.7(a), $\|A\|_{\mathcal{LM}_I(\tilde{\mathcal{A}})} \leq \frac{C}{c^2} (\text{Card}(I))^{\frac{2}{q}} \|e_{\mathcal{A}}\|^2 \|A\|_{\mathcal{LM}_I^p(\mathcal{A})}$. So, $\mathcal{LM}_I^p(\mathcal{A})$ is Banach algebra isomorphic with $\mathcal{LM}_I(\tilde{\mathcal{A}})$. Hence, $\mathcal{LM}_I(\tilde{\mathcal{A}})$ is amenable, and so by Theorem 4.1 of [3], $\tilde{\mathcal{A}}$ is amenable. Therefore, \mathcal{A} is amenable. (ii) \Rightarrow (i): We apply the notations of the above paragraph. By Theorem 4.1 of [3], $\mathcal{LM}_I(\tilde{\mathcal{A}})$ is amenable. Hence, $\mathcal{LM}_I^p(\mathcal{A})$ is amenable. \square

Remark 3.3. The above theorem remains valid, if we replace the condition “ $1 \leq p \leq 2$ ” by “ $\mathcal{LM}_I^p(\mathcal{A})$ is a Banach algebra”.

Example 3.4. Let S be a Brandt semigroup over a finite group G with a finite index I . Then, by Proposition 2.9, Theorem 3.2, and Remark 3.3, the convolution Banach algebra $\ell^p(S)$ is amenable.

Acknowledgments

The first author is grateful to the Office of Graduate Studies of the University of Isfahan for its support and the second author also thanks the University of Bu-Ali Sina (Hamedan) for its support. Also, the authors thank the referee for his invaluable comments.

REFERENCES

- [1] F. Bonsall, J. Duncan, Complete Normed Algebras, Springer-Verlag, New York, 1973.
- [2] H. G. Dales, A. T. M. Lau D. Strauss, *Banach algebras on semigroups and their compactifications*, To appear in the Memoirs of AMS.
- [3] G. H. Esslamzadeh, *Banach algebra structure and amenability of a class of matrix algebras with applications*, J. Funct. Anal. **161** (1999) 364-383.
- [4] G. H. Esslamzadeh, *Ideals and representations of certain semigroup algebras*, Semigroup Forum **69** (2004) 51-56.
- [5] V. Runde, Lectures on Amenability, Lecture Notes in Mathematics, Vol. **1774**, Springer, Berlin, 2002.

S. Naseri

Department of Mathematics, University of Isfahan, Isfahan, Iran.

Email: naserisaber@yahoo.com

H. Samea

Department of Mathematics, University of Bu-Ali Sina, Hamedan, Iran.

Email: h-samea@basu.ac.ir