Bulletin of the Iranian Mathematical Society Vol. 34 No. 2 (2008), pp 39-47.

# ON FINITE GROUPS WITH TWO IRREDUCIBLE CHARACTER DEGREES

A. HEYDARI AND B. TAERI\*

Communicated by Jamshid Moori

ABSTRACT. We characterize non-abelian finite groups with only two irreducible character degree and prime number of non-linear irreducible characters.

## 1. Introduction

Let  $\operatorname{Irr}(G)$  be the set of irreducible complex characters of a finite group G and  $\operatorname{cd}(G) = \{\chi(1) \mid \chi \in \operatorname{Irr}(G)\}$  be the set of irreducible character degrees of G. The question:

Given the set cd(G), what can be said about the structure of G? have been studied by several people. In case  $cd(G) = \{1, m\}$ , for some integer m, the basic tools for studying the question can be found in Chapter 12 of the well known book of Isaacs [8]. Isaacs proved in Theorem 12.5 and Corollary 12.6 the following results.

**Theorem A.** Let G be a finite group and  $cd(G) = \{1, m\}$ . Then, at least one of the followings occurs: (a) G has an abelian normal subgroup of index m.

MSC(2000): Primary: 20C15, 20E99

Keywords: Groups, irreducible characters, degrees, non-linear characters  $% \left( {{{\left( {{{\left( {{{\left( {{{c}}} \right)}} \right.} \right.}} \right)}_{\rm characters}}} \right)$ 

Received: 26 September 2007, Accepted: 9 February 2008

 $* Corresponding \ author$ 

© 2008 Iranian Mathematical Society.

#### 39

(b)  $m = p^e$  for some prime p and G is a direct product of a p-group and an abelian group.

**Theorem B.** Let G be a finite group and  $cd(G) = \{1, m\}$ . Then, G' is an abelian group, where G' is the derived subgroup of G.

If G is non-nilpotent and  $cd(G) = \{1, m\}$ , then clearly (a) of Theorem A holds. Recently, Bianchi *et.al.* [4], characterized non-nilpotent groups G with  $cd(G) = \{1, m\}$ , and proved the following result.

**Theorem C.** Let G be a finite non-nilpotent group. Then,  $cd(G) = \{1, m\}$  if and only if G' is abelian and one of the following holds: (a) m is a prime, and F(G), the fitting subgroup of G, is abelian of index m in G. (b)  $G' \cap Z(G) = 1$  and G/Z(G) is a Frobenius group with kernel  $\frac{G' \times Z(G)}{Z(G)}$ 

(b)  $G' \cap Z(G) = 1$  and G/Z(G) is a Frobenius group with kernel  $\frac{Z(G)}{Z(G)}$ and a cyclic complement of order  $|G: G' \times Z(G)| = m$ , where Z(G)denotes the center of G.

Our notations and terminologies are standard and mainly taken from [8]. We also use the following notations:

$$\begin{split} & \operatorname{Lin}(\mathrm{G}) = \text{the set of linear characters of } G. \\ & \operatorname{Irr}_1(G) = \operatorname{Irr}(G) \setminus \operatorname{Lin}(G). \\ & ES(c,2) = \text{the extra-special group of the order } 2^{2c+1}. \\ & (H,N) = \text{the Frobenius group with kernel } N \text{ and complement } H. \\ & C_n = \text{the cyclic group of order } n. \\ & E(q^n) = \text{the elementary abelian } q\text{-group of rank } n. \\ & Z_i = Z_i(G) = \text{the i-th center of } G. \end{split}$$

Here, we characterize groups G with two irreducible character degrees, that is  $cd(G) = \{1, m\}$ , such that the number of irreducible non-linear characters is a prime p. In [1], Berkovich characterized solvable groups for which the degrees of the characters are distinct, except for one pair (having the same degrees). Thus, by [1] we may assume that  $p \geq 3$ . Firstly, we consider nilpotent groups and prove the following result.

**Theorem 1.** Let G be a finite nilpotent group,  $p \ge 3$ , a prime and m, a positive integer. Then  $cd(G) = \{1, m\}$  and G has p non-linear

*irreducible characters of degree m if and only if one of the followings holds:* 

(a)  $G = C_p \times ES(c, 2)$ ; that is, G is the direct product of a cyclic group of order p and an extra-special group of order  $2^{2c+1}$ .

(b) G is an special 2-group of order  $m^2(p+1)$ , and for each subgroup M of index 2 of G',  $G/M \simeq ES(c, 2)$ , where  $2^c = m$ .

(c) G is dihedral, semidihedral or a generalized quaternion 2-group and |G| = 4(p+1).

If  $cd(G) = \{1, m\}$ , then by Theorem B, G is metabelian. Theorem C characterizes non-nilpotent groups G with  $cd(G) = \{1, m\}$ . We complete the characterization of groups G with  $cd(G) = \{1, m\}$  having p non-linear characters of degree m, by considering the non-nilpotent case in the following theorem.

**Theorem 2.** Let G be a finite non-nilpotent group,  $p \ge 3$ , a prime and m, a positive integer. Then  $cd(G) = \{1, m\}$  and G has p non-linear characters of the degree m if and only if G' is abelian and one of the followings holds:

(a)  $G = (C_m, G')$ , where G is a Frobenius group with kernel of order mp + 1 and a cyclic complement of order m.

(b)  $G/Z(G) = (C_m, (G/Z(G))'), Z(G) = C_p \text{ and } G' = E(q^n), \text{ where } q^n - 1 = m.$  Therefore,  $G = C_p \times (C_{q^n-1}, E(q^n))$  or  $G = C_{p(q^n-1)} \ltimes E(q^n)$ ; that is, G is a semidirect product with kernel  $E(q^n)$ .

(c) *m* is a prime,  $G' \cap Z_{t-1} = m^{t-2} = p+1$  and  $G/Z_{t-1} \simeq (C_m, (G/Z_{t-1})')$ , a Frobenius group with kernel of order *b*, such that  $|G| = m^t b$  and gcd(m, b) = 1.

## 2. Proofs

To prove Theorem 1, firstly we prove the following Lemma for p-groups.

**Lemma 1.** Let G be a finite p-group, where p is a prime,  $cd(G) = \{1, m\}$ and  $|Irr_1(G)|$  is odd. Then, at least one of the followings holds: (a) G is an special 2-group, the index of G' is  $m^2$ , and for each subgroup M of index 2 of G',  $G/M \simeq ES(c, 2)$  such that  $2^c = m$ .

(b) G is dihedral, semidihedral or a generalized quaternion 2-group and |G| = 4(p+1).

**Proof.** Let  $p \neq 2$ . Then, by [8, Exercise 3.16], the number of non-linear characters of G is even, which is a contradiction. Thus,  $|G| = 2^n$  and  $m = 2^c$ . Let  $|\operatorname{Irr}_1(G)| = x$ . From

$$|G| = \sum_{\chi \in \operatorname{Irr}(G)} \chi(1)^2$$

we have

$$2^n = x2^{2c} + |G:G'| = x2^{2c} + 2^k$$

for some positive integer k. Since x is odd, then 2c = k and  $|G| = (x+1)2^{2c}$ . Hence,  $x+1 = 2^r = |G'|$  and  $|G:G'| = 2^{2c}$ . Now, we distinguish two cases below.

**Case 1:**  $G' \leq Z(G)$ . Since G is nilpotent then  $2^{2c} = m^2 = \chi(1)^2$ divides |G : Z(G)|, for all  $\chi \in \operatorname{Irr}_1(G)$  (see [6, Theorem 8.2]). Since  $|G : G'| = 2^{2c}$ , it follows that Z(G) = G'. Now, let M be an arbitrary normal subgroup of index 2 in G'. Note that M is normal in G, since G' is central. If  $|\operatorname{Irr}_1(G/M)| = y$ , then

$$|G/M| = 2^{2c+1} = y2^{2c} + 2^{2c}.$$

So y = 1 and by [2, Main Theorem],  $G/M \simeq ES(c, 2)$ . Thus,  $\Phi(G/M) = (G/M)'$ , where  $\Phi(G)$  is the Frattini subgroup of G, and  $\frac{G/M}{\Phi(G/M)} = \frac{G/M}{G'/M} \simeq G/G'$  is elementary. So,  $\Phi(G)G'/G' = \Phi(G/G') = 1$ , and hence  $\Phi(G) \leq G'$ . Therefore,  $\Phi(G) = G'$ . Now, by [5, Lemma III, 3.14],  $G^2 = G' \leq Z(G)$ . So for each  $g_1, g_2 \in G$ ,  $[g_1, g_2]^2 = [g_1^2, g_2] = 1$ , and hence G' is an elementary abelian group. Thus, G is an special 2-group. Therefore, (a) holds.

**Case 2:**  $G' \notin Z(G)$ . Let  $c \neq 1$ . By [8, Exercise 5.14], each subgroup of index  $2^c$  in G contains G'. Then, each  $\chi \in \operatorname{Irr}_1(G)$  can be induced from a linear character of some normal subgroup of G. Thus, G is an nM-group and by [7, Theorem 3], G' is an elementary abelian subgroup.

(Recall that a group is said nM-group if all its irreducible characters are induced from a linear character of some normal subgroup of G.)

If N is a normal subgroup of G such that G/N is a non-abelian group, since  $\operatorname{cd}(G/N) = \{1, 2^c\}$ , by [6, Theorem 8.2],  $2^{2c}$  divides |G/N|. So, from  $|G/N| = \sum_{\chi \in \operatorname{Irr}(G/N)} \chi(1)^2$ , we have that  $2^{2c}$  divides t = |G/N|: (G/N)'|. But  $t = |G : G'N| \leq |G : G'| = 2^{2c}$ , so that  $t = 2^{2c}$  and  $N \leq G'$ . Therefore,  $Z(G) \leq G'$  and for  $\chi \in \operatorname{Irr}_1(G)$  if  $K = \operatorname{ker}(\chi)$ , then  $K \leq G'$ .

Now, suppose that  $\bar{G} = \frac{G}{K}$ . Since  $cd(\bar{G}) = \{1, 2^c\}$ , then from  $|\bar{G}| = \sum_{\bar{\chi} \in Irr(\bar{G})} \bar{\chi}(1)^2$  we have that  $|Irr_1(G)|$  is odd. By a similar argument for the group  $\bar{G}$  instead of G, we have  $Z(\bar{G}) \leq \bar{G}' = G'/K$ . Since  $\bar{G}$  is an nM-group for each  $\bar{\chi} \in Irr_1(\bar{G})$ , then there exists a normal subgroup  $\bar{H}$  of  $\bar{G}$  of index  $2^c$  and  $\bar{\theta} \in \text{Lin}(\bar{H})$  such that  $\bar{\chi} = \bar{\theta}^{\bar{G}}$ . Let  $\ker(\bar{\theta}) \neq 1$ . From  $Z(\bar{G}) \leq \bar{G}' \leq \bar{H}$ , we have  $\ker(\bar{\theta}) \cap Z(\bar{G}) \neq 1$ , and so  $\bigcap_{x \in \bar{G}} (\ker(\bar{\theta}))^x \neq 1$ . Thus,  $\ker(\bar{\chi}) \neq 1$  (see [8, Lemma 5.11]). But, for each  $xK \in \bar{G}$  we have  $\bar{\chi}(xK) = \chi(x)$  and thus  $\ker(\bar{\chi}) = 1$ , which is a contradiction. Therefore,  $\ker(\bar{\theta}) = 1$  and  $\bar{H}' = \bigcap\{\ker(\lambda) \mid \lambda \in \text{Lin}(\bar{H})\} = 1$ . So,  $\bar{H}$  is abelian and by [8, Theorem 2.32],  $\bar{H}$  is cyclic. Since  $\bar{G}' \leq \bar{H}'$ , then  $\bar{G}'$  is cyclic. Thus,  $\bar{G}'$  is a cyclic elementary abelian group and hence  $|\bar{G}'| = 2$ . Therefore,  $Z(\bar{G}) = \bar{G}'$  and  $G' = Z(\chi)$ , where  $Z(\chi) = \{g \in G \mid |\chi(g)| = \chi(1)\}$ . So,

$$G' \subseteq \bigcap_{\chi \in Irr_1(G)} Z(\chi) = Z(G),$$

which is a contradiction. Therefore, c = 1, |G/G'| = 4, and by [5, Chapter III, Theorem 11.9(a)], G is dihedral, semidihedral or generalized quaternion 2-group. So, (b) holds.

Now, we are ready to prove Theorem 1.

**Proof of Theorem 1:** Since G is a non-abelian nilpotent group, then  $G = Q \times A$ , for some non-abelian Sylow q-subgroup Q and a subgroup A of G. Let A be non-abelian. Let  $\chi_1 \in \operatorname{Irr}_1(Q)$ ,  $\chi_2 \in \operatorname{Irr}_1(A)$  and  $\lambda \in \operatorname{Lin}(A)$ . If  $\chi_1(1) = x_1$  and  $\chi_2(1) = x_2$ , then  $\chi_1\lambda(1) = x_1$  and  $\chi_1\chi_2(1) = x_1x_2$ . So,  $x_1, x_1x_2 \in \operatorname{cd}(G)$ , which is a contradiction. Thus A is abelian.

Now, every irreducible character of G is of the form  $\chi\lambda$ , where  $\chi \in Irr(Q), \lambda \in Irr(A)$ . Since the number of non-linear characters is equal to p, then either Q has only one non-linear character and A has p linear characters or else A = 1.

If Q has only one non-linear character then by [2, Main theorem],  $Q \simeq ES(c, 2)$ , for some positive integer c. Thus,  $G \simeq ES(c, 2) \times C_p$  and (a) holds.

If A = 1 then G is a q-group and by by Lemma 1, G satisfies (b) or (c).

Conversely, suppose  $G = C_p \times ES(c, 2)$ . Since ES(c, 2) has only one non-linear character, then G has p non-linear characters of the same degree. If G satisfies (c), then G has a normal abelian subgroup of index 2, and thus  $cd(G) = \{1, 2\}$  and  $|Irr_1(G)| = p$ .

If G is an special 2-group such that  $|G:G'| = m^2 = 2^{2c}$ , then G' has  $2^r - 1$  distinct subgroups of index 2, where  $|G'| = 2^r$  for some positive integer r. If M is one of such subgroups, then  $G/M \simeq ES(c, 2)$  has only one non-linear character. Then, for  $\chi \in \operatorname{Irr}_1(G)$  there exists only one maximal subgroup  $M_{\chi}$  of G' such that  $M_{\chi} \subseteq \ker \chi$ . Thus, G has at least  $2^r - 1$  non-linear characters of degree  $2^c$  and  $|G:G'| = 2^{2c}$ . Then, from  $\sum_{\chi \in \operatorname{Irr}(G)} \chi(1)^2 = |G|$ , we have  $\operatorname{cd}(G) = \{1, 2^c\}$  and  $|\operatorname{Irr}_1(G)| = p$ .  $\Box$ 

**Proof of Theorem 2:** Since G has only p non-linear characters of degree m, then

$$|G| = \sum_{\chi \in \operatorname{Irr}(G)} \chi(1)^2 = pm^2 + |G:G'|.$$

Since G' is an abelian normal subgroup G, then by Ito's Theorem, see [8, Theorem 6.15],  $\chi(1)$  divides |G:G'|. If  $|G:G'| = mp^2$ , then |G'| = 2 and G is nilpotent, which is a contradiction. So, |G:G'| is one of the followings:

(i) |G:G'| = my, such that  $y \neq p$  and y divides m. If m is not a prime, then by [3, Lemma 2] we have  $G = (C_m, G')$ . If m is not a prime, then by Theorem C,  $G' \cap Z(G) = 1$  and  $G/Z(G) = (C_m, \frac{G' \times Z(G)}{Z(G)})$ . So, On finite groups with two irreducible character degrees

|Z(G)| = y. If  $|\operatorname{Irr}_1(G/Z(G))| = x$ , then

$$|G/Z(G)| = \frac{pm^2 + my}{y} = xm^2 + \left|\frac{G}{Z(G)} : \frac{G'}{Z(G)}\right| = xm^2 + m.$$

So y = 1 and  $G = (C_m, G')$  and (a) holds.

(ii) |G:G'| = pmy, such that y divides m. We have  $G = pm^2 + pmy$ . If  $y \neq 1$  then  $|G'| \leq m$ , and by [8, Exercise 5.14], G is nilpotent, which is a contradiction. So, |G| = pm(m+1). Suppose m is a prime. By Theorem A, G has a normal abelian subgroup M of index m. By [8, Theorem 12.12],  $|M \cap Z(G)| = p$ . If  $Z(G) \nleq M$ , then G = Z(G)M. So, G is abelian, which is a contradiction. So,  $Z(G) \leq M$  and  $Z(G) = C_p$ . Now, suppose m is not a prime. Then, by Theorem C,  $G' \cap Z(G) = 1$  and  $G/Z(G) = \binom{G' \times Z(G)}{Z(G)}$ . Since |G| = pm(m+1), then |Z(G)| = p and  $Z(G) = C_p$ . If  $Z(G) \leqslant G'$ , then  $|\frac{G}{Z(G)} : (\frac{G}{Z(G)})'| = pm$ . Hence, by the above argument,  $m(m+1) = xm^2 + pm$  and m+1 = xm+p, which is a contradiction. So,  $Z(G) \nleq G'$  and  $|\frac{G}{Z(G)} : (\frac{G}{Z(G)})'| = m$ . Let x be the number of non-linear characters of G/Z(G). Then,

$$|G/Z(G)| = \sum_{\chi \in \operatorname{Irr}(G/Z(G))} \chi(1)^2 = xm^2 + m,$$

and hence x = 1. Therefore, by [2, Main Theorem],  $G/Z(G) = (C_{q^n-1}, E(q^n))$  is a Frobenius group with elementary abelian kernel  $E(q^n) = (G/Z(G))' \simeq G'$  and cyclic complement  $C_{q^n-1}$ , for some prime q and integer n such that  $m = q^n - 1$ .

Now G' is a minimal normal subgroup of G. Since H is a proper subgroup of G' which is normal in G, then  $|\frac{G}{H}: (\frac{G}{H})'| = |G:G'| = mp$ . Hence, if G/H, has r non-linear characters of degree m, then  $|G/H| = rm^2 + pm$ . So,  $(rm^2 + pm) | mp(m+1) = |G|$  and (rm+p) | m(p-r) so (rm+p) | (m+1), which is a contradiction. Since G is a non-nilpotent group, then there exists one maximal normal subgroup M of G. Now, G'is a minimal normal abelian subgroup of G, and thus  $G' \cap M = 1$ , and  $G = M \ltimes G'$ . Now,  $G/Z(G) = (C_{q^n-1}, E(q^n))$  and  $G' \leq (C_{q^n-1}, E(q^n))$ . Hence,  $G = Z(G) \times (C_{q^n-1}, E(q^n))$  or  $G = C_{p(q^n-1)} \ltimes E(q^n)$ , a semidirect product with kernel  $G' = E(q^n)$ . So, (b) holds. (iii)  $|G:G'| = m^2$ . We have  $|G| = m^2(p+1)$ . Let m be a prime. Now G has a normal abelian group A of index m and by [8, Lemma 12.2],  $|Z(G) \cap A| = m$ . If  $Z(G) \notin A$ , then G is abelian, and so  $Z(G) \leq A$  and |Z(G)| = m. Suppose that  $|G| = m^t b$ , such that gcd(m, b) = 1. From  $\sum_{\chi \in Irr(G/Z(G))} \chi(1)^2 = |G/Z(G)|$ , we have  $mb = xm^2 + |\frac{G}{Z(G)} : (\frac{G}{Z(G)})'|$ , where  $|Irr_1(G/Z(G))| = x$ . Thus,  $m^2$  divide  $|\frac{G}{Z(G)} : (\frac{G}{Z(G)})'| = |\frac{G}{G'Z(G)}|$ . Now,  $|G:G'| = m^2$ , and so  $Z(G) \leq G'$ . For  $1 \leq i < t$ ,  $G/Z_i$  is a non-nilpotent group and  $cd(G/Z_i) = \{1, m\}$ . Thus, by the above argument,  $|Z(G/Z_i)| = m$ . Hence,  $Z_{t-1}(G)$  is a subgroup of order  $m^{t-1}$  of G. If  $G = G'Z_{t-1}$ , then  $G/Z_{t-1}$  is abelian, which is a contradiction. Therefore  $|G' \cap Z_{t-1}| = m^{t-2}$ .

Now, from  $\sum_{\chi \in \operatorname{Irr}(G/Z_{t-1})} \chi(1)^2 = |G/Z_{t-1}|$ , we have  $mb = xm^2 + |\frac{G}{Z_{t-1}} : (\frac{G}{Z_{t-1}})'|$ , where  $|\operatorname{Irr}_1(G/Z_{t-1})| = x$ . Hence,  $|\frac{G}{Z_{t-1}} : (\frac{G}{Z_{t-1}})'| = m$ . Thus, by [3, Lemma 2],  $G/Z_{t-1} = (C_m, (G/Z_{t-1})')$ .

Now, let *m* be not a prime. Then, by Theorem C,  $G/Z(G) = (C_m, \frac{G' \times Z(G)}{Z(G)})$ , and  $|\frac{G' \times Z(G)}{Z(G)}| = p + 1$ . Then,  $m \mid p$  and m = p is prime, which is a contradiction.

Conversely, if G satisfies (a) or (b), then  $Z(G) \cap G' = 1$  and  $G/Z(G) = (C_m, \frac{G' \times Z(G)}{Z(G)})$ . So, by Theorem C,  $\operatorname{cd}(G) = \{1, m\}$  and  $|\operatorname{Irr}_1(G)| = p$ . If G satisfies (c), then  $|G| = m^t b$  and G' is a normal abelian subgroup of index  $m^2$  of G. So, G' contains N, a normal abelian subgroup of order b of G, and thus  $NZ_{t-1}$  is a normal abelian subgroup of index m of G. So, by [8, Theorem 6.15],  $\operatorname{cd}(G) = \{1, m\}$ . Now, since  $|G : G'| = m^2$ , then  $|\operatorname{Irr}_1(G)| = p$ . This completes the proof.  $\Box$ 

#### Acknowledgment

This work was partially supported by Center of Excellence of Mathematics of Isfahan University of Technology (CEAMA).

### References

- Y. Berkovich, Finte solvable groups in which only two non-linear irreducible characters have equal degree, J. Algebra 184 (1996) 584-603.
- [2] Y. Berkovich, D. Chillag and M. Herzog, Finite groups in which the degrees of the non-linear irreducible characters are distinct, *Proc. Amer. Math. Soc.* 115 (1992) 955-959.

- [3] Y. Berkovich and A. Mann, On sums of degrees of irreducible characters, J. Algebra 199 (1998) 646-665.
- [4] M. Bianchi, A. G. B. Mauri, M. Herzog, G. Qian and W. Shi, Characterization of non-nilpotent groups with two irreducible character degrees, J. Algebra 284 (2005) 326-332.
- [5] B. Huppert, Endliche Gruppen I, Speringer-Verlag, 1967.
- [6] B. Huppert, Character Theory of Finite Groups, de Gruyter, Berlin, 1998.
- [7] A. Mann, Normally monomial p-groups, J. Algebra 300 (2006) 2-9.
- [8] I. M. Isaacs, Character Theory of Finite Groups, Academic Prees, New York, 1976.

### Abbas Heydari

Department of Mathematical Sciences, Isfahan University of Technology, Isfahan 84156-83111, Iran.

Email: a-heydari@math.iut.ac.ir

#### Bijan Taeri

Department of Mathematical Sciences, Isfahan University of Technology, Isfahan 84156-83111, Iran.

Email: b.taeri@cc.iut.ac.ir