LINEAR PRESERVING GD-MAJORIZATION FUNCTIONS FROM $M_{n,m}$ TO $M_{n,k}$

A. ARMANDNEJAD* AND H. HEYDARI

Communicated by Heydar Radjavi

ABSTRACT. Let $\mathbf{M}_{n,m}$ be the vector space of all $n \times m$ real matrices. For $A, B \in \mathbf{M}_{n,m}$, it is said that B is gd-majorized by A (written $A \succ_{gd} B$) if for every $x \in \mathbb{R}^n$ there exists a g-doubly stochastic matrix D_x such that $Bx = D_x(Ax)$. Here, we show that if $A \succ_{gd} B$, then there exists a g-doubly stochastic matrix D (independent of x) such that B = DA. Also, the possible structures of linear preserving gd-majorization functions from $\mathbf{M}_{n,m}$ to $\mathbf{M}_{n,k}$ are found. Finally, all linear strongly preserving gd-majorization functions from $\mathbf{M}_{n,m}$ to $\mathbf{M}_{n,k}$ are characterized.

1. Introduction

Let $T: \mathbf{M}_{n,m} \to \mathbf{M}_{n,k}$ be a linear function and let \sim be a relation on both $\mathbf{M}_{n,m}$ and $\mathbf{M}_{n,k}$. We say that T preserves \sim when $X \sim Y$ implies $TX \sim TY$; if in addition, $TX \sim TY$ implies $X \sim Y$, we say that T strongly preserves \sim . For $x, y \in \mathbb{R}^n$, it is said that x is vector majorized by y (written $y \succ x$) if there exists a doubly stochastic matrix D such that x = Dy. For given $X, Y \in \mathbf{M}_{n,m}$, it is said that X is directionally majorized by Y (written $Y \succ_d X$) if $Yv \succ Xv$, for all $v \in \mathbb{R}^m$. The linear preservers of \succ_d on $\mathbf{M}_{n,m}$ have been characterized in [7]. Some

MSC(2010): Primary: 15A04; Secondary: 15A21, 15A51.

Keywords: Doubly stochastic matrix, g-doubly stochastic matrix, girectional majorization, gd-majorization, linear preserver.

Received: 27 April 2009, Accepted: 28 December 2009.

^{*}Corresponding author

^{© 2011} Iranian Mathematical Society.

types of majorization and their linear preservers are presented in [1], [5] and [6]. Throughout the paper, the notation \mathbf{M}_n is fixed for the algebra of all $n \times n$ real matrices. The space $\mathbf{M}_{n,1}$ of all $n \times 1$ real vectors is denoted by the usual notation \mathbb{R}^n . The collection of all $n \times n$ permutation matrices is denoted by \mathcal{P}_n . The notation $X = [x_1|\cdots|x_m]$ is used for an $n \times m$ matrix with $x_j \in \mathbb{R}^n$ as the jth column of X $(1 \leq j \leq m)$. The letters \mathbf{J} and e stand for the square matrix and the vector, which respectively all of their entries are 1, and the dimensions of the matrix \mathbf{J} and the vector e are understood from the context. The standard basis of \mathbb{R}^n is denoted by $\{e_1, \dots, e_n\}$. The notation A^t stands for the transpose of a given matrix A. For a given vector $x \in \mathbb{R}^n$, tr(x) is the sum of all components of x. Now, we state an extension of a familiar result [7, Theorem 2] about linear functions preserving directional majorization from $\mathbf{M}_{n,m}$ to $\mathbf{M}_{n,k}$.

Proposition 1.1. [2, Theorem 1.3] A linear function $T: \mathbf{M}_{n,m} \to \mathbf{M}_{n,k}$ preserves directional majorization if and only if one of the following holds.

(i) There exist
$$A_1, \ldots, A_m \in \mathbf{M}_{n,k}$$
 such that $T(X) = \sum_{j=1}^m (trx_j)A_j$,

where, $X = [x_1| \cdots |x_m]$.

(ii) There exist $R, S \in M_{m,k}$ and $P \in \mathcal{P}_n$ such that T(X) = PXR + JXS.

A (not necessarily nonnegative) matrix $D \in \mathbf{M}_n$ with the properties De=e and $D^te=e$ is said to be a g-doubly stochastic matrix. This generalization of stochastic matrices was introduced in [4]. We denote the set of all $n \times n$ g-doubly stochastic matrices by \mathbf{GD}_n . For matrices $A, B \in \mathbf{M}_{n,m}$, it is said that B is gs-majorized by A (written $A \succ_{gs} B$) if there exists an $n \times n$ g-doubly stochastic matrix D such that B=DA. In [3], the authors found the possible structures of all linear operators preserving \succ_{gs} on $\mathbf{M}_{n,m}$ as follows.

Proposition 1.2. [3, Theorem 3.3] Let $T : \mathbf{M}_{n,m} \to \mathbf{M}_{n,m}$ be a linear operator that preserves \succ_{gs} . Then, one of the following holds.

- (i) There exist $A_1, \dots, A_m \in \mathbf{M}_{n,m}$ such that $T(X) = \sum_{j=1}^m tr(x_j) A_j$, where, $X = [x_1 | \dots | x_m]$.
- (ii) There exist $S \in \mathbf{M}_m$, $a_1, \ldots, a_m \in \mathbb{R}^m$ and invertible matrices $D_1, \ldots, D_m \in \mathbf{GD}_n$ such that $T(X) = [D_1 X a_1 | \cdots | D_m X a_m] + JXS$.

For $A, B \in \mathbf{M}_{n,m}$, it is said that B is gd-majorized by A (written $A \succ_{gd} B$) if $Ax \succ_{gs} Bx$, for all $x \in \mathbb{R}^m$. In fact, $A \succ_{gd} B$ if and only

if, for every $x \in \mathbb{R}^m$, there exists a g-doubly stochastic matrix D_x such that $Bx = D_x(Ax)$. Here we prove the following theorem which gives the possible structures of all linear functions preserving \succ_{gd} from $\mathbf{M}_{n,m}$ to $\mathbf{M}_{n,k}$.

Theorem 1.3. Let $T : \mathbf{M}_{n,m} \to \mathbf{M}_{n,k}$ be a linear function that preserves \succ_{qd} . Then, one of the following holds.

- (i) There exist $A_1, \ldots, A_m \in \mathbf{M}_{n,k}$ such that $T(X) = \sum_{j=1}^m tr(x_j)A_j$, where, $X = [x_1 | \ldots | x_m]$.
- (ii) There exist $R, S \in \mathbf{M}_{m,k}$ and an invertible matrix $D \in \mathbf{GD}_n$ such that $T(X) = DXR + \mathbf{J}XS$.
- (iii) There exist $S \in \mathbf{M}_{m,k}$, $a \in \mathbb{R}^m$, $r_1, \ldots, r_k \in \mathbb{R}$ and invertible matrices $D_1, \ldots, D_k \in \mathbf{GD}_n$ such that $T(X) = [r_1D_1Xa| \ldots |r_kD_kXa] + \mathbf{J}XS$.

2. Gd-Majorization

In this section, we present some properties of \succ_{gd} and then show that the relation implies \succ_{gs} on $\mathbf{M}_{n,m}$.

Lemma 2.1. Let x and y be two distinct vectors in \mathbb{R}^n . Then, $x \succ_{gs} y$ if and only if $x \notin span\{e\}$ and tr(x) = tr(y).

Proposition 2.2. Let $A = [a_1| \cdots |a_m], B = [b_1| \cdots |b_m] \in \mathbf{M}_{n,m}$. Then, B is gd-majorized by A if and only if the following conditions hold.

- (a) For every i $(1 \le i \le m)$, $tr(a_i) = tr(b_i)$; in other words, $A^t e = B^t e$.
 - (b) For every $x \in \mathbb{R}^m$ such that $Ax \in span\{e\}$, Ax = Bx.

Proof. It is clear that $A \succ_{gd} B$ implies the conditions (a) and (b). Conversely, assume (a) and (b) hold. For $x \in \mathbb{R}^m$, if $Ax \in span\{e\}$, then Bx = Ax, and hence $Ax \succ_{gs} Bx$. If $Ax \notin span\{e\}$, since $tr(b_i) = tr(a_i)$ for every i $(1 \le i \le n)$, then tr(Bx) = tr(Ax). So, $Ax \succ_{gs} Bx$, by Lemma 2.1, and therefore $A \succ_{gd} B$.

Remark 2.3. Let $X, Y \in \mathbf{M}_{n,m}$, $A, B \in \mathbf{GD}_n$, $C \in \mathbf{M}_m$ and $\alpha, \beta \in \mathbb{R}$ such that A, B and C are invertible and $\alpha \neq 0$. Then, the following conditions are equivalent:

- (1) $X \succ_{gd} Y$.
- (2) $AX \succ_{gd} BY$.
- (3) $\alpha X + \beta J_{nm} \succ_{gd} \alpha Y + \beta J_{nm}$, where $J_{nm} \in \mathbf{M}_{n,m}$ is the matrix with all entries equal to one.

(4) $XC \succ_{gd} YC$.

Now, we show that \succ_{gs} coincides with \succ_{gd} on $\mathbf{M}_{n,m}$.

Lemma 2.4. Let $A, B \in \mathbf{M}_n$. If A is invertible and $A \succ_{gd} B$, then $A \succ_{gs} B$.

Proof. Put $D = BA^{-1}$. Since DA = B, it is enough to show that D is a g-doubly stochastic matrix. By invertibility of A, there exists a unique $x_0 \in \mathbb{R}^n$ such that $Ax_0 = e$, and hence $Bx_0 = e$, by Proposition 2.2. So, $De = (BA^{-1})e = B(A^{-1}e) = Bx_0 = e$. On the other hand, $A^te = B^te$, by Proposition 2.2, and hence $D^te = (BA^{-1})^t(e) = (A^{-1})^t(B^te) = (A^{-1})^tA^te = e$.

Lemma 2.5. Let A = [C|D] and $B = [E|F] \in \mathbf{M}_{n,m}$, where $C, E \in \mathbf{M}_{n,k}$ and $D, F \in \mathbf{M}_{n,(m-k)}$. Suppose that the columns of D are generated by the columns of C. If $A \succ_{qd} B$ and $C \succ_{gs} E$, then $A \succ_{gs} B$.

Proof. Since $C \succ_{gs} E$, then there exits a g-doubly stochastic matrix $R \in \mathbf{GD}_n$ such that RC = E, and hence $Rc_i = e_i$ $(1 \le i \le k)$, where c_i and e_i are the ith columns of C and E, respectively. We claim that RA = B. Suppose that d is the first column of D. Then, there exist scalars $\alpha_1, \ldots, \alpha_k \in \mathbb{R}$ such that $d = \sum_{i=1}^k \alpha_i c_i$. Put $x_0 = (\alpha_1, \ldots, \alpha_k, -1, 0, \ldots, 0)^t \in \mathbb{R}^m$. Then, $Ax_0 = 0$, and hence $Bx_0 = 0$ by, Proposition 2.2. So, $f = \sum_{i=1}^k \alpha_i e_i$, where f is the first column of F. Thus, $Rd = \sum_{i=1}^k \alpha_i Rc_i = \sum_{i=1}^k \alpha_i e_i = f$. This argument is valid for other columns of D and F, and hence RA = B.

Theorem 2.6. The concepts of gs and gd-majorization on $\mathbf{M}_{n,m}$ are the same.

Proof. It is clear that \succ_{gs} implies \succ_{gd} , and so we prove only the converse. Let $A, B \in \mathbf{M}_{n,m}$ and $A \succ_{gd} B$. By Remark 2.3, $A \succ_{gd} B$ if and only if $AP \succ_{gd} BP$, for every permutation matrix $P \in \mathbf{M}_n$. Then, without loss of generality, we can assume that A = [C|D], where, $C = [c_1|\cdots|c_k] \in \mathbf{M}_{n,k}$ is a full rank matrix, $D = [d_1|\cdots|d_{m-k}] \in \mathbf{M}_{n,(m-k)}$ and $d_1, \cdots, d_{m-k} \in span\{c_1, \cdots, c_k\}$ (C or D can be vacuous). It is clear that $k \leq n$. Choose some vectors $c_{k+1}, \cdots, c_n \in \mathbb{R}^n$ such that $C' = [c_1|\cdots|c_n] \in \mathbf{M}_n$ is an invertible matrix. Consider the matrices $E \in \mathbf{M}_{n,k}$ and $F \in \mathbf{M}_{n,(m-k)}$ such that B = [E|F]. Since C' is invertible, then there exists a unique vector $x = (x_1, \cdots, x_n)^t \in \mathbb{R}^n$ such that C'x = e. Put $y = (x_1, \cdots, x_k)^t \in \mathbb{R}^k$. Since $A \succ_{gd} B$, then

 $C \succ_{gd} E$, by Proposition 2.2, and hence $Cy \succ_{gs} Ey$. So, there exists a g-doubly stochastic matrix $R \in \mathbf{M}_n$ such that RCy = Ey. Put $E' = [E|Rc_{k+1}|\cdots|Rc_n] \in \mathbf{M}_n$. Then, $E'x = x_1e_1 + \cdots + x_ke_k + x_{k+1}Rc_{k+1} + \cdots + x_nRc_n$, where e_i is the ith column of E. Since RCy = Ey then $x_1Rc_1 + \cdots + x_kRc_k = x_1e_1 + \cdots + x_ke_k$, then $E'x = x_1Rc_1 + \cdots + x_nRc_n = R(C'x) = Re = e$. On the other hand, $tr(e_i) = tr(c_i)$, for every i $(1 \le i \le k)$, and $tr(c_i) = tr(Rc_i)$, for every i $(k+1 \le i \le n)$. Then, $C' \succ_{gd} E'$, by Proposition 2.2. Therefore, $C' \succ_{gs} E'$, by Lemma 2.4, and hence $C \succ_{gs} E$. Since $d_1, \cdots, d_{m-k} \in span\{c_1, \cdots, c_k\}$ and $A \succ_{gd} B$, we get $A \succ_{gs} B$, by Lemma 2.5.

3. Linear Preservers

In this section, we prove the following statements which shed light on the structure of linear functions preserving \succ_{gd} from $\mathbf{M}_{n,m}$ to $\mathbf{M}_{n,k}$.

Theorem 3.1. Let $T: \mathbf{M}_{n,m} \to \mathbb{R}^n$ be a linear function. Then, T preserves \succ_{qs} if and only if one of the following holds.

- (a) There exist $a_1, \ldots, a_m \in \mathbb{R}^n$ such that $T(X) = \sum_{j=1}^m tr(x_j)a_j$, where, $X = [x_1 | \ldots | x_m]$.
- (b) There exist $a, b \in \mathbb{R}^m$ and an invertible matrix $A \in \mathbf{GD}_n$ such that $T(X) = AXa + \mathbf{J}Xb$.

Proof. The fact that each of the conditions (a) or (b) is sufficient for T to be a preserver of \succ_{gs} is easy to prove. So, we prove the necessity of the conditions. Define $T': \mathbf{M}_{n,m} \to \mathbf{M}_{n,m}$ by T'(X) = [T(X)|0], where 0 denotes an $n \times (m-1)$ zero block. Clearly, T' is a linear function which preserves \succ_{gs} . Then, by Proposition 1.2, T' has one of the following forms.

(i)
$$T'(X) = \sum_{j=1}^{m} tr(x_j)B_j$$
, for some $B_1, \ldots, B_m \in \mathbf{M}_{n,m}$. So, $T(X) =$

 $\sum_{j=1}^{m} tr(x_j)a_j$, where a_j is the first column of B_j , for every j $(1 \le j \le m)$, and hence (a) holds.

(ii) $T'(X) = [D_1Xa_1| \dots | D_mXa_m] + \mathbf{J}XS$, for some $S \in \mathbf{M}_m$, $a_1, \dots, a_m \in \mathbb{R}^n$ and invertible matrices $D_1, \dots, D_m \in \mathbf{GD}_n$. So, $TX = D_1Xa_1 + \mathbf{J}Xb$, where b is the first column of S, and hence (b) holds.

Lemma 3.2. [3, Lemma 3.1] Let $A \in GD_n$ be invertible. Then, the following conditions are equivalent.

- (a) $A = \alpha I + \beta J$, for some $\alpha, \beta \in \mathbb{R}$.
- (b) $(x+Ay) \succ_{qs} (Dx+ADy)$, for all $D \in \mathbf{GD}_n$ and for all $x, y \in \mathbb{R}^n$.

Remark 3.3. Assume that T_1 and T_2 are of the form (a) and (b) in Theorem 3.1, respectively. Then, $T_1 = T_2$ if and only if a = 0 and $a_j = \lambda_j e$, for every j $(1 \le j \le m)$, where a, a_j $(1 \le j \le m)$ and $b = (\lambda_1, \ldots, \lambda_m)^t$ are as in Theorem 3.1.

Lemma 3.4. Let $T_1, T_2 : \mathbf{M}_{n,m} \to \mathbb{R}^n$ be two linear preservers of \succ_{gs} such that $T_1 + T_2$ preserves \succ_{gs} . If $T_1(X) = DXa + JXb$, for some $a, b \in \mathbb{R}^m$, $a \neq 0$ and an invertible matrix $D \in \mathbf{GD}_n$, then $T_2(X) = D'Xc + JXd$, for some $c, d \in \mathbb{R}^m$ and an invertible matrix $D' \in \mathbf{GD}_n$.

Proof. Since $T_1 + T_2$ preserves \succ_{gs} , then $T_1 + T_2$ is of the form (a) or (b) in Theorem 3.1. Now, consider two cases.

Case 1: Suppose that $T_1 + T_2$ is of the form (a). Since T_2 preserves \succ_{gs} , it is of the form (a) or (b) in Theorem 3.1. Assume, if possible, T_2 is of the form (a). Then, $T_1 = (T_1 + T_2) - T_2$ is of the form (a), as well. So, by Remark 3.3, we obtain a = 0, which is a contradiction. Therefore, T_2 is of the form (b).

Case 2: Suppose that $T_1 + T_2$ is of the form (b). So, $(T_1 + T_2)(X) = BXa' + \mathbf{J}Xb'$, for some $a', b' \in \mathbb{R}^m$ and invertible matrix $B \in \mathbf{GD}_n$. Assume, if possible, T_2 is of the form (a) and is not of the form (b). Then, by Theorem 3.1 and Remark 3.3, there exist (not all in $span\{e\}$) $a_1, \ldots, a_m \in \mathbb{R}^n$ such that $T_2(X) = \sum_{j=1}^m tr(x_j)a_j$. Without loss of generality, suppose that $a_1 \notin span\{e\}$. Put $X := [e|0| \ldots |0] \in \mathbf{M}_{n,m}$. So,

(3.1)
$$na_1 = \sum_{j=1}^m tr(x_j)a_j = T_2(X)$$
$$= (T_1 + T_2 - T_1)(X)$$
$$= [a'_1 + nb'_1 - a_1 - nb_1]e,$$

where, a'_1 , b'_1 , a_1 and b_1 are the first entry of a', b', a and b, respectively, which is a contradiction. Therefore, T_2 is of the form (b), and hence there exist $c, d \in \mathbb{R}^m$ and an invertible matrix $D' \in \mathbf{GD}_n$ such that $T_2(X) = D'Xc + \mathbf{J}Xd$.

Now, we can prove Theorem 1.3.

Proof of Theorem 1.3. Suppose that T preserves \succ_{gd} . Then, for every i $(1 \le i \le k)$, $T_i = E_i \circ T : \mathbf{M}_{n,m} \to \mathbb{R}^n$ preserves \succ_{gd} , where,

 $E_i: \mathbf{M}_{n,k} \to \mathbb{R}^n$ is defined by $E_i(A) = A\epsilon_i$. Thus, T_i is of the form (a) or (b) in Theorem 3.1. Now, consider two cases.

Case 1: Assume T_i is of the form (a), for every i $(1 \le i \le k)$. Then, $T_i(X) = \sum_{j=1}^m tr(x_j)a_j^i$, for some $a_j^i \in \mathbb{R}^n$. Put $A_j := [a_j^1|\dots|a_j^k]$, for every

j $(1 \le j \le m)$. So, $T(X) = \sum_{j=1}^{m} tr(x_j)A_j$, and hence the condition (i) holds.

Case 2: Assume there exists p $(1 \le p \le k)$ such that $T_p(X) = D_pXa_p + \mathbf{J}Xb_p$, for some $a_p, b_p \in \mathbb{R}^k$, $a_p \ne 0$ and an invertible matrix $D_p \in \mathbf{GD}_n$. Since T preserves \succ_{gd} , so $T_p + T_j$ preserves \succ_{gd} , for every j $(1 \le j \le k)$. Then, by Lemma 3.4, $T_j(X) = D_jXa_j + \mathbf{J}Xb_j$, for some $a_j, b_j \in \mathbb{R}^m$, and an invertible matrix $D_j \in \mathbf{GD}_n$. So,

$$T(X) = [T_1(X)|\cdots|T_k(X)]$$

$$= [D_1Xa_1 + \mathbf{J}Xb_1|\cdots|D_mXa_m + \mathbf{J}Xb_k]$$

$$= [D_1Xa_1|\cdots|D_kXa_k] + \mathbf{J}X[b_1|\cdots|b_k].$$

If $rank[a_1|\cdots|a_k] \geq 2$, then, without loss of generality, we may assume that $rank[a_1|a_2] = 2$. Since for every $X \in \mathbf{M}_{n,m}$ and every $D \in \mathbf{GD}_n$, $X \succ_{gs} DX$, then $(T_1 + T_2)X \succ_{gs} (T_1 + T_2)(DX)$, and hence $D_1Xa_1 + D_2Xa_2 \succ_{gs} D_1DXa_1 + D_2DXa_2$. So, for every $D \in \mathbf{GD}_n$,

$$Xa_1 + (D_1^{-1}D_2)Xa_2 \succ_{gs} DXa_1 + (D_1^{-1}D_2)DXa_2, \forall X \in \mathbf{M}_{n,m}.$$

Since a_1 and a_2 are linearly independent, we may put some suitable X in the above relation and obtain the following:

$$x + (D_1^{-1}D_2)y \succ_{qs} Dx + (D_1^{-1}D_2)Dy , \forall x, y \in \mathbb{R}^m , \forall D \in \mathbf{GD}_n .$$

Then, by Lemma 3.2, $D_1^{-1}D_2 = \lambda_1 I + \mu_1 \mathbf{J}$, and hence $D_2 = \lambda_1 D_1 + \mu_1 \mathbf{J}$, for some $\lambda_1, \mu_1 \in \mathbb{R}$. For every i $(2 \leq i \leq k)$, with $a_i \neq 0$, it is clear that $\{a_1, a_i\}$ or $\{a_2, a_i\}$ is linearly independent, and so, by a similar argument as above, $D_i = \lambda_i D_1 + \mu_i \mathbf{J}$, for some $\lambda_i, \mu_i \in \mathbb{R}$. Set $D := D_1$. Then, for every i $(1 \leq i \leq k)$, $D_i = \lambda_i D + \mu_i \mathbf{J}$, for some $\lambda_i, \mu_i \in \mathbb{R}$, and hence $T(X) = DXR + \mathbf{J}XS$, where, $R = [\lambda_1 a_1 | \cdots | \lambda_k a_k]$ and $S = [\mu_1 a_1 + b_1 | \cdots | \mu_k a_k + b_k]$. Therefore, the condition (ii) holds. If $rank[a_1| \dots |a_k] \leq 1$, then there exist $a \in \mathbb{R}^m$ and $r_1, \dots, r_k \in \mathbb{R}$ such that for every i $(1 \leq i \leq k)$, $a_i = r_i a$. Therefore, $T(X) = [r_1 D_1 X a| \dots | r_k D_k X a] + \mathbf{J}XS$, where, $S = [b_1| \dots | b_k]$, and hence the condition (iii) holds.

It is easy to show that if T is of the form (i) or (ii) in Theorem 1.3, then T preserves \succ_{ad} . The following example shows that there is a linear function of the form (iii) not preserving \succ_{ad} .

Example 3.5. Suppose that $T: \mathbf{M}_{3,2} \to \mathbf{M}_{3,2}$ is defined by T(X) =Example 3.3. Suppose that 1 : A.3., 2 : A.3.

Theorem 1.3. Put
$$A := \begin{pmatrix} 1 & 1 \\ -1 & 2 \\ 1 & 0 \end{pmatrix}$$
 and $B := \begin{pmatrix} -1 & 1 \\ 1 & 0 \\ 1 & 2 \end{pmatrix}$. It is easy

to show that $B \succ_{ad} A$ and $TB \not\succ_{ad} TA$. Then, T does not preserve \succ_{ad} .

It is clear that the form (ii) is a special case of the form (iii) in Theorem 1.3 (put $D_1 = \cdots = D_k := D$ and $R := [r_1 a | \cdots | r_k a]$). The following example shows that there is a linear function preserving \succ_{qd} , which is of the form (iii) but is not of the form (ii).

Example 3.6. [3, Example 3.5] Let $T: \mathbf{M}_{3,2} \to \mathbf{M}_{3,2}$ be defined by $T(X) = [X\epsilon_1|PX\epsilon_1], \text{ where, } P = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$ Then, T preserves \succ_{ad} and T is not of the form (ii) in Theorem

Now, we state the following lemma which characterizes all strong linear preservers of \succ_{qd} from $\mathbf{M}_{n,m}$ to $\mathbf{M}_{n,k}$.

Lemma 3.7. [2, Lemma 2.4] Let $T: \mathbf{M}_{n,m} \to \mathbf{M}_{n,k}$ be a linear function of the form T(X) = XR + JXS, for some $R, S \in \mathbf{M}_{m,k}$. Then, T is injective if and only if R and R + nS are full-rank matrices.

Proof. It is easy to see that the matrix representation of T with respect to the standard bases of $M_{n,m}$ and $M_{n,k}$ is similar to the following block matrix:

$$\begin{pmatrix} R+nS & & * \\ & R & \\ & & \ddots & \\ 0 & & & R \end{pmatrix} \in M_{nk,nm} .$$

Therefore, T is injective if and only if R and R + nS are full-rank matrices.

If T is a strong linear preserver of \succ_{qd} and T(A) = 0, then $T(0) \succ_{qd}$ T(A). So, $0 \succ_{qd} A$, and hence A = 0.

Remark 3.8. Every strong linear preserver of \succ_{gd} from $M_{n,m}$ to $M_{n,k}$ is injective.

If m=1, then the following theorem is obtained from Theorem 3.1. So, in the proof we may assume $m\geq 2$.

Theorem 3.9. Let $T: \mathbf{M}_{n,m} \to \mathbf{M}_{n,k}$ be a linear function. Then, T strongly preserves \succ_{gd} if and only if there exist an invertible matrix $D \in \mathbf{GD}_n$ and matrices $R, S \in \mathbf{M}_{m,k}$ such that R and R + nS are full-rank matrices and $TX = DXR + \mathbf{J}XS$.

Proof. If T is of the form $TX = DXR + \mathbf{J}XS$, for some invertible matrix $D \in \mathbf{GD}_n$ and full-rank matrices $R, R + nS \in \mathbf{M}_{m,k}$, then it is easy to show that T is a strong linear preserver of \succ_{gd} . Conversely, assume T is a strong linear preserver of \succ_{gd} . So, T is of the form (i), (ii) or (iii) in Theorem 1.3. If T is of the form (i), then T is not injective, which is a contradiction. If T is of the form (iii), then we can choose $0 \neq b \in (span\{a\})^{\perp}$, by the assumption $m \geq 2$. Put $X_0 := [b \mid -b \mid 0 \mid \cdots \mid 0]^t \in \mathbf{M}_{n,m}$. So, $X_0 \neq 0$ and $T(X_0) = 0$, which is a contradiction. Therefore, T is of the form (ii), and by Lemma 3.7, R and R + nS are full-rank matrices.

Acknowledgments

The authors are very grateful to the anonymous referee for his/her helpful comments and for detecting an error in the first version of the article which helped us to state Theorem 2.6.

References

- [1] A. Armandnejad, Right GW-majorization on $\mathbf{M}_{n,m}$, Bull. Iranian Math. Soc. **35** (2009) 69-76.
- [2] A. Armandnejad and H. R. Afshin, Linear functions preserving multivariate and directional majorization, *Iran. J. Math. Sci. Inform.* **5** (2010) 1-5.
- [3] A. Armandnejad and A. Salemi, The structure of linear preservers of gs-majorization, *Bull. Iranian Math. Soc.* **32** (2006) 31-42.
- [4] H. Chiang and C.-K. Li, Generalized doubly stochastic matrices and linear preservers, *Linear Multilinear Algebra* 53 (2005) 1-11.
- [5] A. M. Hasani and M. Radjabalipuor, The structure of linear operators strongly preserving majorizations of matrices, *Electron. J. Linear Algebra* 15 (2006) 260-268.

- [6] F. Khalooei and A. Salemi, The structure of linear preservers of left matrix majorization on \mathbb{R}^p , Electron. J. Linear Algebra 18 (2009) 88-97.
- [7] C.-K. Li and E. Poon, Linear operators preserving directional majorization, *Linear Algebra Appl.* **325** (2001) 141-146.

Ali Armandnejad and Hossein Heydari

Department of Mathematics, Vali-e-Asr University of Rafsanjan, P.O.Box 7713936417, Rafsanjan, Iran

Email: armandnejad@mail.vru.ac.ir and

Email: hosein8212@yahoo.com