RINGS OF SINGULARITIES

H. LENZING

Communicated by Henning Krause

ABSTRACT. This paper is a slightly revised version of an introduction into singularity theory corresponding to a series of lectures given at the "Advanced School and Conference on homological and geometrical methods in representation theory" at the International Centre for Theoretical Physics (ICTP), Miramare - Trieste, Italy, 11-29 January 2010. We show how to associate to a triple of positive integers (p_1, p_2, p_3) a two-dimensional isolated graded singularity by an elementary procedure that works over any field k (with no assumptions on characteristic, algebraic closedness or cardinality). This assignment starts from the triangle singularity $x_1^{p_1} + x_2^{p_2} + x_3^{p_3}$ and when applied to the Platonic (or Dynkin) triples, it produces the famous list of A-D-E-singularities. As another particular case, the procedure yields Arnold's famous strange duality list consisting of the 14 exceptional unimodular singularities (and an infinite sequence of further singularities not treated here in detail). As we are going to show, weighted projective lines and various triangulated categories attached to them play a key role in the study of the triangle and associated singularities.

MSC(2010): Primary: 14J17; Secondary: 16G20, 16G50.

Keywords: Weighted projective line, (extended) canonical algebra, simple singularity,

Arnold's strange duality, stable category of vector bundles.

Received: 5 January 2011, Accepted: 5 May 2011.

^{© 2011} Iranian Mathematical Society.

1. Introduction

Weighted projective lines provide the mathematical environment to study triangle singularities $x_1^{p_1} + x_2^{p_2} + x_3^{p_3}$ (and suitable finite systems of such equations). To analyze the arising singularities, we attach to them an abelian hereditary k-linear category $\mathcal H$ with Serre duality having a tilting object T, whose endomorphism ring is a canonical algebra (with three arms). This category $\mathcal H$ has an interpretation as the category of coherent sheaves coh- $\mathbb X$ on a weighted projective line $\mathbb X$ whose weight type is just the triple of integers we started with.

Weighted projective lines and their defining equations of shape $f=x_1^{p_1}+x_2^{p_2}+x_3^{p_3}$ have a long history going back to Klein [13] and Poincaré [25]. Accordingly, their study has a high contact surface with many mathematical subjects, classical and modern. Among the many related subjects, we mention representation theory of algebras and groups, invariant theory, function theory, orbifolds, 3-manifolds, singularities and the study of nilpotent operators. The formal definition of the category of coherent sheaves is due to Geigle and the author in 1987; see [7]. In particular, through tilting theory a close link to the representation theory of finite dimensional algebras has been established. This concerns in particular, Ringel's study of canonical algebras and their representations, see [26]. The quoted paper further contains a link to commutative algebra, through the relationship between vector bundles on a weighted projective line and the (graded) Cohen-Macaulay modules over the algebra $S = k[x_1, x_2, x_3]/(f)$ and related algebras. This link to singularity theory is exploited in the present paper. For an alternative categorybased access to weighted projective lines, we refer to the papers [3, 20].

In the focus of this paper is the construction and analysis of three types of (usually not equivalent) triangulated categories which are naturally attached to coh-X. These categories all have a tilting object and thus each one yields an explicit link to the representation theory of finite dimensional algebras. One of the three categories is the bounded derived category of coh-X, and the other two are obtained from two (usually different) Frobenius structures on the category vect-X of vector bundles on X. Due to a general result of Happel [8], the associated stable categories are triangulated. Following the work of Buchweitz (1987), they are equivalent to the stable categories of the (suitably graded) Cohen-Macaulay modules; see [2].

The topics discussed in the final part of the paper are related to recent independent work (2006) of Kajiura-Saito-Takahashi-Ueda [11, 30] and Lenzing-de la Peña [21] on Fuchsian singularities. A key role in these developments is played by a theorem of Orlov (arXiv 2005, published in 2009) dealing with the analysis of singularities by means of the triangulated category of (graded) singularities (the stable derived category in Buchweitz's sense). An important feature of our treatment is its independence of the characteristic of the base field.

Historically, an important aspect of singularity theory is incorporated in the following table of simple singularities:

Dynkin's diagram Δ	simple singularity f_{Δ}
$A_n: \circ - \circ - \circ \cdots \circ - \circ$	$zy + x^{n+1}$
\mathbb{D}_n : \circ \circ \circ \circ \circ \circ	$z^2 + y^2 x + x^{n-1}$
\mathbb{E}_6 : \circ $ \circ$ $ \circ$ $ \circ$ $ \circ$	$z^2 + y^4 + x^3$
$\mathbb{E}_7: \circ - \circ - \overset{\circ}{\circ} - \circ - \circ - \circ$	$z^2 + y^3x + x^3$
$\mathbb{E}_8: \circ - \circ - \circ - \circ - \circ - \circ $ $The A-D-E-singularities$	

For the moment, the above singularities should be considered to be defined over the field \mathbb{C} of complex numbers, giving rise to the simple isolated singularities $R_{\Delta} = \mathbb{C}[x,y,z]/(f_{\Delta})$. Just as the Dynkin diagrams, these singularities appear in many mathematical contexts where here we only mention a few. They appear for instance in

(1) the classification of critical points of differential maps.

(2) rings of invariants under the natural action of finite subgroups of $\mathbf{SL}(2,\mathbb{Z})$ acting on $\mathbb{C}[[X,Y]]$. (A graded version with the action on $\mathbb{C}[X,Y]$ is also available.) This links the topic with the ancient classification of regular or Platonic solids and its modern treatment as the McKay correspondence; see [5].

(3) finite dimensional representation theory as suitable orbit algebras of the Auslander-Reiten translation for tame hereditary algebras.

For further information on the omnipresence of Dynkin diagrams and singularities we refer to [4, 10, 29].

A look at the table does not reveal any building law. And, in the setting discussed, the equations f_{Δ} are far from being unique, since the primary object of interest — in this context — is the ring R_{Δ} , which is not changed if we change the variables x, y and z by a linear base change with coefficients in \mathbb{C} . Our first aim of here is therefore to work in a graded setting in order


- (1) to present an elementary method to generate the singularities f_{Δ} systematically, and basically produces a unique list,
- (2) to work over an arbitrary field, and to design the construction as to be independent on any extra assumptions on this (characteristic, algebraically closedness),
- (3) to recover from f_{Δ} or the associated graded ring $k[x, y, z]/(f_{\Delta})$, the Dynkin diagram Δ .

In these notes we are giving a more direct link to finite dimensional representation theory via one associated abelian hereditary category and three related triangulated categories. The link is then established by means of appropriate tilting objects and their endomorphism rings.

2. From Dynkin diagrams to simple singularities

2.1. **Dynkin diagrams.** Assume we are given a triple (p_1, p_2, p_3) of integers $p_i \geq 0$. By the symbol $[p_1, p_2, p_3]$ we denote the star-shaped

graph



with base point, where the number p_i indicates the length (number of vertices) of the *i*th branch (which for $p_i = 1$ degenerates to the base point). Here the length of the *i*th branch counts the number of vertices in the branch including the fat base point. In this notation, a **Dynkin diagram** Δ is just a star $[p_1, p_2, p_3]$ satisfying the inequality

$$(2.1) 1/p_1 + 1/p_2 + 1/p_3 > 1.$$

We thus have $\mathbb{D}_n = [2, 2, n-2]$ with $n \geq 4$, $\mathbb{E}_6 = [2, 3, 3]$, $\mathbb{E}_7 = [2, 3, 4]$ and $\mathbb{E}_8 = [2, 3, 5]$. For \mathbb{A}_n , there is some ambiguity, since any triple (p, q, 1), with p + q - 1 = n, produces the Dynkin diagram \mathbb{A}_n . Taking the base point into account, what we are going to do consistently, the ambiguity obviously disappears. Any triple (p_1, p_2, p_3) satisfying the inequality (2.1) is called a **Dynkin triple** or, following Klein [13], a Platonic triple.

Triangle singularities We work over an arbitrary field k and fix a triple (p_1, p_2, p_3) of integers ≥ 1 , called **weight triple**. Let $\mathbb{L} = \mathbb{L}(p_1, p_2, p_3)$ be the abelian group given by generators $\vec{x}_1, \vec{x}_2, \vec{x}_3$ and the defining relations $p_1\vec{x}_1 = p_2\vec{x}_2 = p_3\vec{x}_3 =: \vec{c}$. The element \vec{c} is called the *canonical element* of \mathbb{L} . As is easily seen, the group \mathbb{L} has rank one, and thus has shape $\mathbb{L} \cong \mathbb{Z} \oplus F$, where F is a finite (abelian) group. As a group, \mathbb{L} is not particularly interesting. We are therefore putting an additional structure on \mathbb{L} .

First of all, \mathbb{L} is an ordered group with the members from $\mathbb{N}\vec{x}_1 + \mathbb{N}\vec{x}_2 + \mathbb{N}\vec{x}_3$ forming its positive cone. Thus, $\vec{x} \leq \vec{y}$ if and only if $\vec{y} - \vec{x}$ is a positive integral linear combination of the generators \vec{x}_1, \vec{x}_2 and \vec{x}_3 . Putting $\bar{p} = \text{lcm}(p_1, p_2, p_3)$, there is a uniquely defined homomorphism of groups, actually a homomorphism of ordered groups $\delta : \mathbb{L} \longrightarrow \mathbb{Z}$ sending each generator \vec{x}_i to \bar{p}/p_i . We further note that $\delta : \mathbb{L} \to \mathbb{Z}$ is surjective and its kernel is the (finite) torsion subgroup of \mathbb{L} . In order to deal with elements of \mathbb{L} explicitly, it is useful to have the following

property: each element \vec{x} of \mathbb{L} can be uniquely written in **normal form**

(2.2)
$$\vec{x} = \sum_{i=1}^{3} \ell_i \vec{x}_i + \ell \vec{c}$$
 with $0 \le \ell_i < p_i$ and $\ell \in \mathbb{Z}$.

Moreover, if an element \vec{x} is in normal form as above, then $\vec{x} \geq 0$ if and only if $\ell \geq 0$.

There is a further element of \mathbb{L} which is important for reasons becoming clear later. This is the **dualizing element** $\vec{\omega} = \vec{c} - (\vec{x}_1 + \vec{x}_2 + \vec{x}_3)$. For the moment we remark that the dualizing element is useful to determine how far the order \leq on \mathbb{L} is away from a total order. Indeed, it is easily established that an element \vec{x} of \mathbb{L} either satisfies $\vec{x} \geq 0$ or $\vec{x} \leq \vec{c} + \vec{\omega}$. We are now in a position to introduce the **triangle singularity**¹

(2.3)
$$h_{(p_1,p_2,p_3)} = x_1^{p_1} + x_2^{p_2} + x_3^{p_3}$$

over k and the associated algebra $S = k[x_1, x_2, x_3]/(x_1^{p_1} + x_2^{p_2} + x_3^{p_3})$. By forming the k-linear span of all monomials $x_1^{\ell_1} x_2^{\ell_2} x_3^{\ell_3}$ having the same degree $\vec{x} = \ell_1 \vec{x}_1 + \ell_2 \vec{x}_2 + \ell_3 \vec{x}_3$, we obtain a finite dimensional k-space $S_{\vec{x}}$ such that $S = \bigoplus_{\vec{x} \in \mathbb{L}} S_{\vec{x}}$.

Proposition 2.1. Assume (p_1, p_2, p_3) is a weight triple. Then, the following properties hold.

(a) The k-algebra S is positively \mathbb{L} -graded by attaching degree $\vec{x_i}$ to each generator x_i . That is,

$$S = \bigoplus_{\vec{x} \ge 0} S_{\vec{x}}, \quad S_0 = k, \quad S_{\vec{x}} \cdot S_{\vec{y}} \subseteq S_{\vec{x} + \vec{y}} \quad \text{ for all } \vec{x}, \vec{y} \in \mathbb{L}.$$

Moreover, the homogeneous components $S_{\vec{x}}$ of S are finite dimensional over k.

(b) Restricting the grading of S to the subgroup $\mathbb{Z}\vec{c}$, we obtain the **heart**

$$H = S_{|\mathbb{Z}\vec{c}} = \bigoplus_{n \geq 0} S_{n\vec{c}} = k[x_1^{p_1}, x_2^{p_2}]$$

of S which 'is' the polynomial algebra in the 'variables' $x_1^{p_1}$ and $x_2^{p_2}$, which are viewed to be homogeneous of degree one. Accordingly, $H = \bigoplus_{n>0} H_n$, with H_n of k-dimension n+1 for $n \geq 0$.

¹Properly speaking, this requires that all $p_i \geq 2$. By abuse of language, we extend the terminology to the present slightly more general setting.

(c) If $\vec{x} = \ell_1 \vec{x}_1 + \ell_2 \vec{x}_2 + \ell_3 \vec{x}_3 + \ell \vec{c}$ has normal form with $\ell \ge 0$, then

$$(2.4) S_{\vec{x}} = x_1^{\ell_1} x_2^{\ell_2} x_3^{\ell_3} H_{\ell}$$

$$\dim_k S_{\vec{x}} = \ell + 1.$$

Proof. Assertion (c) follows by collecting monomials having the same degree and using the relation $x_3^{p_3} = -(x_1^{p_1} + x_2^{p_2})$. Property (c) then implies assertions (a) and (b).

The next consequence explains the role of the order on \mathbb{L} .

Corollary 2.2. For $\vec{x} \in \mathbb{L}$, we have $\vec{x} \geq 0$ if and only if $S_{\vec{x}} \neq 0$.

Corollary 2.3. As an \mathbb{L} -graded algebra, S satisfies the following properties.

- (a) S is graded-integral, that is, if x and y are non-zero homogeneous elements of S, then also xy is non-zero.
- (b) The k-algebra S is **graded-factorial**, that is, each non-zero homogeneous element is a product of homogeneous prime elements. (Here, a homogeneous element p of S is called **prime** if S/(p) is graded-integral.)
- (c) The non-zero homogeneous prime elements of S naturally form a $\mathbf{P}^1(k)$ -family.

Proof. Concerning (a) it follows from formula (2.4) that each homogeneous element of S has the form $x_1^{l_1}x_2^{l_2}x_3^{l_3}h_l$, where h_l is a homogeneous element of the heart H of S which is an integral domain. Claim (b) follows in a similar way, observing that H is clearly graded-factorial. Using the known structure of homogeneous prime polynomials in H, claim (c) follows along the same lines. (If k is not algebraically closed, we have to interpret the projective line as a scheme, not as a variety.)

Comments 2.4. (1) One should not mix the concepts "graded-integral" and "graded plus integral". For instance, we have in characteristic two that the square of $x = x_1 + x_2 + x_3$ is zero in the $\mathbb{L}(2,2,2)$ -graded algebra $S = k[x_1, x_2, x_3]/(x_1^2 + x_2^2 + x_3^2)$ which, as we have pointed out, is graded-integral. Note, in this context, that x is not a homogeneous element with regard to the \mathbb{L} -grading.

(2) A similar remark replies to all other "graded concepts". So, a k-algebra R is a "graded field" (to be thought of as one word!) if each non-zero homogeneous element has a homogeneous inverse with respect to multiplication. In the graded sense therefore, the algebra of Laurent

polynomials $K = k[X, X^{-1}]$ (considered to be \mathbb{Z} -graded by attaching degree 1 to X) is a graded field. Similar care has to be taken, when dealing with graded modules. For instance, the concepts "graded-injective" module and "graded plus injective" module will usually be different. For instance, dealing with the \mathbb{Z} -graded polynomial algebra k[X], where we give X degree one, the graded module K of Laurent polynomials, graded as above, is an injective object in the category of graded modules, but K is far from being injective in the category of all k[X]-modules.

- (3) If k is an algebraically closed field, then the algebras $S(p_1, p_2, p_3)$ exhaust the graded-factorial affine k-algebras of Krull dimension two which have three generators and are graded by a rank-one abelian group. This follows from a result of Kussin [14]. In that paper, the more general situation of an arbitrary number of weights is treated, yielding the corresponding result. For simplicity, we restrict to three weights which is simplifying notation and nevertheless allows to cover the most interesting singularities.
- **Lemma 2.5.** The degree $\delta(\vec{\omega})$ is negative if and only if the triple (p_1, p_2, p_3) is, up to reordering, one of (1,1,1), (1,1,p), (1,p,q), (2,2,n), (2,3,3), (2,3,4) and (2,3,5). Moreover, we have $\delta(\vec{\omega}) = 0$ if and only if the triple is one of the triples (3,3,3), (2,4,4) and (2,3,6), called tubular. For all the remaining triples, called wild, we have $\delta(\vec{\omega}) > 0$.
- 2.2. The simple singularity attached to a Dynkin diagram. We are now going to show how to associate to each Dynkin diagram, equivalently to each triple (p_1, p_2, p_3) of negative degree, a simple singularity. This is simply done by restricting the \mathbb{L} -graded algebra to the (infinite cyclic) subgroup $\mathbb{Z}\vec{\omega}$. This restriction is defined to be the $\mathbb{Z}\vec{\omega}$ -graded algebra $R = \bigoplus_{n \in \mathbb{Z}} S_{n\vec{\omega}}$. By our assumption on the degree of $\vec{\omega}$, we can have $n\vec{\omega} \geq 0$ only if $n \leq 0$. By means of the bijection $\mathbb{Z} \to \mathbb{Z}\vec{\omega}$, $n \mapsto -n\vec{\omega}$, and we may thus view R as the positively \mathbb{Z} -graded algebra with homogeneous components $R_n = S_{-n\vec{\omega}}$.

For the base field of complex numbers, an alternative interpretation of this passage from S to R is important. Due to Klein [13], each Dynkin diagram $\Delta = [p_1, p_2, p_3]$ yields an action of the corresponding binary polyhedral group G_{Δ} on the polynomial algebra $\mathbb{C}[X, Y]$. Klein's central result states that the subalgebra of relative invariants is isomorphic to S and the subalgebra of absolute invariants is isomorphic to R; see [7, Subsection 5.4.1].

Theorem 2.6. For any weight triple (p_1, p_2, p_3) with $\delta(\vec{\omega}) < 0$, let $\Delta = [p_1, p_2, p_3]$ denote the attached Dynkin diagram. Then, the restriction of the grading of $S = S(p_1, p_2, p_3)$ to $\mathbb{Z}\vec{\omega}$ yields a \mathbb{Z} -graded algebra having a (minimal) system of three homogeneous generators x, y, z, all being monomials in x_1, x_2, x_3 . With this choice of generators, we have

$$R := S_{|\mathbb{Z}\vec{\omega}} = k[x, y, z]/(f_{\Delta}),$$

where f_{Δ} is the simple graded singularity from the table below. Moreover, with the above assumptions, the singularity f_{Δ} can be chosen as a sum of monomials in x, y, z and thus is unique:

Dynkin diagram Δ	generators(x, y, z)	$\deg(x,y,z)$	relation f_{Δ}	$deg(f_{\Delta})$			
$\mathbb{A}_{p+q} = [p, q]$	$(x_1 x_2, x_2^{p+q}, x_1^{p+q})$	(1, p, q)	$x^{p+q} - yz$	p+q			
$\mathbb{D}_{2l+2} = [2, 2, 2l]$	$(x_3^2, x_1^2, x_1 x_2 x_3)$	(2, 2l, 2l + 1)	$z^2 + x(y^2 + yx^l)$	4l + 2			
$\mathbb{D}_{2l+3} = [2, 2, 2l+1]$	$(x_3^2, x_1 x_2, x_1^2 x_3)$	(2, 2l+1, 2l+2)	$z^2 + x(y^2 + z x^l)$	4l + 4			
$\mathbb{E}_6 = [2, 3, 3]$	$(x_1, x_2 x_3, x_2^3)$	(3, 4, 6)	$z^2 + y^3 + x^2 z$	12			
$\mathbb{E}_7 = [2, 3, 4]$	$(x_2, x_3^2, x_1 x_3)$	(4, 6, 9)	$z^2 + y^3 + x^3 y$	18			
$\mathbb{E}_8 = [2, 3, 5]$	(x_3, x_2, x_1)	(6, 10, 15)	$z^2 + y^3 + x^5$	30			
The simple graded surface singularities (arbitrary base field)							

We postpone a discussion of the proof for a moment in order to point out an interesting consequence.

Corollary 2.7. Each algebra $R_{\Delta} = k[x,y,z]/(f_{\Delta})$, with f_{Δ} from the above list, where k is an arbitrary field, is an integral domain (both in the graded and ungraded sense). Accordingly, the polynomial f_{Δ} is a prime element in k[x,y,z] (both in the graded and ungraded sense).

(Just judging from the form of the relations f_{Δ} , this is not obvious at all; compare, for example Comments 2.4 (2). With Theorem 2.6 at hand, this proof becomes very easy.)

Proof. Since S is graded integral, the same holds true for its restriction $R = S_{|\mathbb{Z}\vec{\omega}}$. Now, any positively \mathbb{Z} -graded algebra R which is graded integral is also integral. For this, write two non-zero elements x and y as a sum of homogeneous elements $x = x_0 + x_1 + \cdots + x_m$ and $y = y_0 + y_1 + \cdots + y_n$, with leading terms x_m and y_n different from zero. Then, the product of x and y has the non-zero leading term $x_m y_n$ and hence is non-zero.

We now sketch the **proof of Theorem 2.6**. The proof relies on two useful lemmas. Recall for this that the **Poincaré series** or the Hilbert-Poincaré series of a positively \mathbb{Z} -graded algebra $R = \bigoplus_{n\geq 0} R_n$, with finite dimensional components R_n , is the formal power series in x given as $P_R = \sum_{n\geq 0} \dim_k R_n x^n$.

Lemma 2.8. Let (p_1, p_2, p_3) be a weight triple, where each $p_i \geq 2$.

(a) If $\delta(\vec{\omega}) < 0$, then the Poincaré series of $R = S_{|\mathbb{Z}\vec{\omega}}$ with $R_n = S_{-n\vec{\omega}}$ is

(2.6)
$$P_R = -\frac{1}{1-x} - \frac{1}{(1-x)^2} + \frac{1}{1-x} \sum_{i=1}^3 \frac{1}{1-x^{p_i}}.$$

(b) If $\delta(\vec{\omega}) > 0$, then the Poincaré series of $R = S_{|\mathbb{Z}\vec{\omega}}$ with $R_n = S_{n\vec{\omega}}$ is

(2.7)
$$P_R = x + \frac{1}{1-x} + \frac{x}{(1-x)^2} - \frac{x}{1-x} \sum_{i=1}^{3} \frac{1}{1-x^{p_i}}.$$

Proof. (a): We sketch the argument assuming $\delta(\vec{\omega}) < 0$ and all $p_i \geq 2$. In this case, the element $-\vec{\omega} = x_1 + x_2 + x_3 - \vec{c}$ is already in normal form. For $n \geq 0$, the element $n\vec{x}_i = \text{has normal form } (n-p_i[\frac{n}{p_i}])\vec{x}_i + [\frac{n}{p_i}]\vec{c}$, where the bracket notation [q] denotes the integral part of a rational number q. For $n \geq 0$, it follows that the normal form of $-n\vec{\omega}$ is given by

$$-n\vec{\omega} = \sum_{i=1}^{3} \left(n - p_i \left[\frac{n}{p_i} \right] \right) \vec{x}_i + \left(-n + \sum_{i=1}^{3} \left[\frac{n}{p_i} \right] \right) \vec{c}.$$

One then uses Proposition 2.1 and takes care of what happens for small values of n. The claim follows.

(b): The proof is similar, calculating this time the normal form of $n\vec{\omega}$ and using that the normal form of $\vec{\omega}$ is $(\sum_{i=1}^{3} (p_i - 1) \vec{x}_i) - 2\vec{c}$.

Lemma 2.9. Assume the algebra $A = k[u_1, u_2, u_3]/(f)$ is positively \mathbb{Z} -graded such that the generators u_i and the relation f are homogeneous of degree $c_i \geq 1$ and d, respectively. Then, the Hilbert series P_A of A is given by the rational function

(2.8)
$$P_A = \frac{1 - x^d}{(1 - x^{c_1})(1 - x^{c_2})(1 - x^{c_3})}.$$

Proof. The polynomial ring $A_i = k[u_i]$ with $\deg(u_i) = c_i$ has Hilbert series

 $1/(1-x^{c_i})$. As the tensor product of the A_i , the polynomial algebra $B=k[u_1,u_2,u_3]$ thus gets the Hilbert series $\sum_{n\geq 0}b_nx^n=\prod_{i=1}^3\,1/(1-x^{c_i})$. Finally, since f has degree d, we get exact sequences

$$0 \to B_{n-d} \xrightarrow{f} B_n \longrightarrow A_n \to 0$$
, yielding $\dim_k A_n = b_n - b_{n-d}$ and then $P_A = (1 - x^d)P_B$. This proves the claim.

Proof of Theorem 2.6. To prove the theorem, each row of the table is separately dealt with. For two cases, we show the arguments involved; the remaining cases can be dealt with in a similar fashion. First, we deal with the case $\mathbb{E}_8 = [2,3,5]$. Here, practically nothing is to show, since in this case, $\mathbb{Z}\vec{\omega} = \mathbb{L}$, that is, up to renaming the grading group, the algebra R coincides with S. In more detail, we have $-6\vec{\omega} = \vec{x}_3$, $-10\vec{\omega} = \vec{x}_2$ and $14\vec{\omega} = \vec{x}_1$. The corresponding components are $R_6 = kx_3$, $R_{10} = kx_2$ and $R_{15} = kx_1$. Thus, $x = x_3$, $y = x_2$ and $z = x_1$ are homogeneous generators for R satisfying the relation $f_{\Delta} = z^2 + y^3 + x^5$. Thus, $R = k[x, y, z]/(f_{\Delta})$, as claimed. Of course, in this case we do not need the two lemmas stated above.

Next, we deal with the case $\mathbb{E}_6 = [2, 3, 3]$. In calculating the normal form of $-n\vec{\omega}$, $n = 0, 1, 2, \cdots$, we first determine for small values of n those multiples $-n\vec{\omega}$ which are ≥ 0 and then, by means of Proposition 2.1 (c), we determine the members of R_n . Here, we get

$$\begin{array}{lll} -3\vec{\omega} & = & \vec{x}_1 & \text{hence } R_3 = kx_1 \\ -4\vec{\omega} & = & \vec{x}_2 + \vec{x}_3 & \text{hence } R_4 = kx_2x_3 \\ -6\vec{\omega} & = & \vec{c} & \text{hence } R_6 = kx_2^3 + kx_3^3. \end{array}$$

Restricting to monomials, we have no choice in the first two cases, obtaining $x = x_1$ of degree 3 and $y = x_2x_3$ of degree 4. Concerning the third case, we have three monomials in x_1, x_2, x_3 lying in R_6 , namely x_1^2 , x_2^3 and x_3^3 . Since x_1^2 equals x^2 , only the choices $z=x_2^3$, and respectively $z=x_3^3$, make sense. Because of weight type (2,3,3), these two choices are equivalent, and so let $z=x_2^3$. As is easily checked, the elements x, y, z are indeed generators for R (use some almost-periodicity of the expression $-n\vec{\omega}$ with "period" 6 = lcm(2,3,3), resulting in some almostperiodic building law for R_n of the same "period"). The canonical homomorphism ϕ from the polynomial algebra $k[u_1, u_2, u_3]$ to R sending u_1, u_2, u_3 to x, y, z is therefore surjective; moreover, ϕ is a homomorphism of graded algebras if we put $deg(u_1, u_2, u_3) = (3, 4, 6)$. Finally, x,y,z satisfy the relation $f_{\Delta}(x,y,z)=0$ since using $x_1^2+x_2^3+x_3^3=0$, we get $z^2=-x_2^3(x_1^2+x_2^3)=-zx^2-y^3$. Hence, we obtain a surjective algebra homomorphism $\psi: k[u_1,u_2,u_3]/(f_{\Delta}(u_1,u_2,u_3)) \to R$, which preserves degrees. Lemmas 2.9 and 2.8 show that both algebras have the same Poincaré series, and it follows that ψ is an isomorphism, as claimed.

2.3. Preliminary conclusions.

Comments 2.10. We comment on various aspects of the theorem.

- (1) In Theorem 2.6, we show uniqueness of the relation f_{Δ} if the minimal generators x, y and z for R_{Δ} are chosen from the set of monomials in x_1, x_2, x_3 . However, any choice of a minimal triple of homogeneous generators x, y, z yields a valid relation g_{Δ} . For the types $\mathbb{E}_7 = [2, 3, 4]$ and $\mathbb{E}_8 = [2, 3, 5]$, each such system x, y, z is formed by monomials in x_1, x_2, x_3 , up to multiplication with non-zero scalars. For the remaining cases, we have a choice. This explains what in the literature one often finds relations which are different from those derived here.
- (2) For instance, for type $\mathbb{E}_6 = [2,3,3]$, the usual form of the relation is given as $g = z^2 + y^4 + x^3$ and not as $f = z^2 + y^3 + x^2z$. We show how by a simple base change, equation f transforms into equation g provided the base field k is algebraically closed of characteristic $\neq 2$. First, we note that for an arbitrary $\lambda \in k$, the elements x, y and $\overline{z} = z + \lambda x^2$ are again a minimal set of generators for R having degrees 3, 4 and 6, respectively. Substitution into f yields the new relation $(\lambda^2 \lambda)x^4 + (1 2\lambda)x^2 + y^3 + \overline{z}^2$. We now put $\lambda = 1/2$ such that the quadratic term in x disappears and introduce the new variable $\overline{x} = \mu x$, where μ is a 4th root of -1/4. This yields, as claimed, for the new generators $\overline{x}, y, \overline{z}$, the relation $\overline{z}^2 + y^3 + \overline{x}^4$. With similar arguments, the relations f_{Δ} for \mathbb{D}_n can be transformed into those from the previous list. Again, this works for fields of characteristic $\neq 2$, which are algebraically closed.
- (3) The degrees of the generators x, y, z and the degree of f_{Δ} are important numerical invariants of the graded singularity f_{Δ} . For instance, the degree of f_{Δ} equals the Coxeter number h_{Δ} of the Dynkin diagram Δ , which equals the period of the Coxeter transformation of the Dynkin diagram Δ , and thus reflects important homological information of the Auslander-Reiten translation for mod- $k\Delta$, where $k\Delta$ is the path algebra. If S_1, \ldots, S_n denote the simple $k\Delta$ -modules (up to isomorphism) and P_1, \ldots, P_n (respectively I_1, \ldots, I_n) are their projective covers (respectively injective hulls), then the Coxeter transformation Φ of Δ is the automorphism of the Grothendieck group $K_0(\text{mod-}k\Delta)$ sending each class $[P_i]$ to the class $-[I_i]$.

Another more conceptual description states that the Auslander-Reiten translation τ is a self-equivalence of the bounded derived category $D^b(\text{mod-}k\Delta)$ inducing the transformation Φ on the Grothendieck $K_0(D^b(\text{mod-}k\Delta)) = K_0(\text{mod-}k\Delta)$. It is well-known that Φ is periodic in the Dynkin situation with the numbers n+1, 2(n-1), 12, 18 and 30 being

the periods for \mathbb{A}_n , \mathbb{D}_n , \mathbb{E}_6 , \mathbb{E}_7 and \mathbb{E}_8 , respectively. Here, the equality of the numbers deg f_{Δ} and h_{Δ} occurs, a surprising coincidence. Assuming a more advanced level we will return to this question in Section 5, where the relationship will easily follow on a conceptual level.

(4) The table contains further interesting information. The sum of the degrees of the generators x, y, z always equals $1 + \deg(f_{\Delta})$.

Summary 2.11. We summarize what we have achieved and also address some obvious questions.

- (1) To each Dynkin diagram $\Delta = [p_1, p_2, p_3]$, the restriction of the $\mathbb{L}(p_1, p_2, p_3)$ -graded triangle singularity $S = S(p_1, p_2, p_3)$ to the infinite cyclic group $\mathbb{Z}\vec{\omega}$, identified by \mathbb{Z} via $n\vec{\omega} \leftrightarrow -n$, yields a positively \mathbb{Z} -graded k-algebra $R_{\Delta} = k[x, y, z]/(f_{\Delta})$, where f_{Δ} is a homogeneous prime polynomial, whose degree is the Coxeter number h_{Δ} of Δ . Moreover, R_{Δ} is a \mathbb{Z} -graded integral domain which is noetherian of Krull dimension two. Unlike the \mathbb{L} -graded k-algebra S, the k-algebra R is no longer graded factorial (except for the Dynkin diagram $\mathbb{E}_9 = [2, 3, 5]$).
- (2) Our treatment still leaves important questions open: what is the conceptual role of the grading group \mathbb{L} ? What is the special role of the dualizing element $\vec{\omega}$ in \mathbb{L} that makes the correspondence $\Delta \mapsto R_{\Delta}$ work? So far, our correspondence $\Delta \mapsto R_{\Delta}$ looks a bit ad-hoc. On the other hand, we have seen that the restriction of S to the subgroup $\mathbb{Z}\vec{c}$ of \mathbb{L} , generated by the canonical element, always yields the polynomial algebra $k[x,y], x = x_1^{p_1}, y = x_2^{p_2}$, where x and y both get degree one. So, what are the properties that are making $\vec{\omega}$ so special that for $\vec{\omega}$ the correspondence works?
- (3) Later, we describe a setting, where the above questions get a natural answer. To give a brief indication already, we first comment how to think of the \mathbb{L} -graded algebra S. For many questions, it is natural to replace the \mathbb{L} -graded algebra by its **companion category**, which is equipped with a natural shift-action of \mathbb{L} . This means to consider the k-linear category $[\mathbb{L}; S]$ given by the following data:
 - the objects are just the elements \vec{x} of the grading group \mathbb{L} ,
 - the morphism space $\operatorname{Hom}(\vec{x}, \vec{y})$ equals $S_{\vec{y}-\vec{x}}$,
 - composition of morphisms corresponds to the multiplication of S,
 - an element $\vec{x} \in \mathbb{L}$ sends an object \vec{y} to the object $\vec{y}(\vec{x}) := \vec{x} + \vec{y}$ and yields on morphisms the mapping

$$S_{\vec{z}-\vec{y}} = \operatorname{Hom}(\vec{y}, \vec{z}) \to \operatorname{Hom}(\vec{y}(\vec{x}), \vec{z}(\vec{x})) = S_{\vec{z}-\vec{y}},$$

corresponding to the identity map on $S_{\vec{z}-\vec{y}}$.

In Section 3, we are going to construct the category $\operatorname{coh-X}$ of coherent sheaves on the weighted projective line $\mathbb X$ of weight type (p_1,p_2,p_3) , and we will see there that our companion category $[\mathbb L;S]$ is equivalent to the category of all line bundles on $\mathbb X$. Under this equivalence, moreover, the action of $\vec{\omega}$ on $[\mathbb L;S]$ corresponds to the (restriction of the) Auslander-Reiten translation of $\operatorname{coh-X}$. In that sense, the $\mathbb L$ -graded algebra S embodies the properties of the Auslander-Reiten translation of the category $\operatorname{coh-X}$. It thus comes as a surprise that studying the simple singularity f_{Δ} means to study the Auslander-Reiten translation on the category $\operatorname{coh-X}$ and vice versa.

(4) We mention in this context that weighted projective lines (also those with more than three weights) appear in the following context: assume that the base field k is algebraically closed, and \mathcal{H} is a connected hereditary abelian k-linear category, which is Ext-finite and has a tilting object. Then, \mathcal{H} is derived equivalent to the module category $\operatorname{mod-}k\vec{\Delta}$ over the path algebra of a quiver $\vec{\Delta}$ or to a category $\operatorname{coh-}\mathbb{X}$ of coherent sheaves on a weighted projective line \mathbb{X} ; see [9]. If k is not algebraically closed, then the class of weighted projective lines has to be enlarged to take many further cases into account; compare for this, [15], [19] and [27].

3. From singularities to diagrams


3.1. An analysis of the problem. We have seen in Section 2 how to attach to a Dynkin diagram $\Delta = [p_1, p_2, p_3]$, that is, to a weight triple (p_1, p_2, p_3) whose dualizing element satisfies $\delta(\vec{\omega}) < 0$, a simple \mathbb{Z} -graded surface singularity R by forming the restriction of the \mathbb{L} -graded triangle singularity $S = k[x_1, x_2, x_3]/(x_1^{p_1} + x_2^{p_2} + x_3^{p_3})$ to the subgroup $\mathbb{Z}\vec{\omega}$ of \mathbb{L} . Under the present assumptions, the group $\mathbb{Z}\vec{\omega}$ is infinite cyclic; we identify $\mathbb{Z}\vec{\omega}$ with the integers by means of the correspondence $-n\vec{\omega} \leftrightarrow n$. This way we have obtained a list (f_{Δ}) , Δ Dynkin, of simple graded surface singularities, a list working for any field.

Remark 3.1. Certain aspects of the theory, nevertheless, still need clarification.

(1) We need a conceptual understanding why it is natural to consider the restriction of the \mathbb{L} -grading of S to $\mathbb{Z}\vec{\omega}$ and not to another


- infinite cyclic subgroup of \mathbb{L} . (Usually, there are many such subgroups around.)
- (2) Assume we are presented a (graded) singularity f_{Δ} from the list (but without the list itself and, of course, without the attached Dynkin label). How can we recover the Dynkin diagram, giving rise to it?
- (3) More generally, and this time not restricting to the Dynkin triples, we want to analyze the complexity (or shape) of an isolated graded surface singularity R by attaching suitable canonical invariants, which, in the special case of Dynkin triples, will contain the information on the Dynkin diagram in question.
- 3.2. Dynkin and extended Dynkin diagrams. For the discussion to follow, it is useful to have a clear conception of the natural bijection between Dynkin and extended Dynkin diagrams. (Note that they are not just lists of graphs! More structure is around.) The correspondence is given by looking at subadditive (respectively additive) functions.
- (a) Recall that a positive integral function λ on a graph Δ is additive in a vertex v provided that $2\lambda(v) = \sum_{v-p} \lambda(p)$, where the sum is over all vertices which are incident to v. Subadditivity in v means that we weaken the condition to $2\lambda(v) \geq \sum_{v-p} \lambda(p)$.
- (b) Dynkin diagrams are exactly the connected finite graphs such that there is a unique (normalized) subadditive function which fails to be additive in a single vertex. Given a Dynkin graph Δ , let v be the vertex where a subadditive function fails to be additive. Attaching to v a new edge with a vertex yields an extended Dynkin diagram, denoted by $\tilde{\Delta}$.
- (c) Extended Dynkin diagrams are exactly the finite connected graphs admitting an additive function. These function are all proportional, and it is possible to choose one, called normalized, attaining value 1. Deleting any vertex (and adjacent edges) then yields a Dynkin diagram.
- (d) The two procedures in (b) and (c) are inverse to each other (on the level of isomorphism classes of graphs). We illustrate this by an example.

The Dynkin graph \mathbb{D}_6 admits the subadditive function depicted below:



It is additive, except in the framed vertex. Adding a new edge here yields the extended Dynkin graph $\tilde{\mathbb{D}}_6$.

Conversely, the extended Dynkin graph \mathbb{D}_6 has a unique normalized additive function λ as depicted below:



There are four vertices v with $\lambda(v) = 1$. As we see, it does not matter which one we delete. The four choices give rise to the "same" Dynkin diagram.

3.3. Sheafification by the Serre construction. We now describe how to attach to the \mathbb{L} -graded singularity S a hereditary K-linear category which is Hom-finite. (A similar construction will later be discussed for the \mathbb{Z} -graded algebra R if $\delta(\vec{\omega})$ is non-zero.)

First, we form the abelian k-linear category of finitely generated \mathbb{L} -graded S-modules, which we denote by $\mathrm{mod}^{\mathbb{L}}$ -S. The objects of this category are the finitely generated \mathbb{L} -graded S-modules. We thus have $M = \bigoplus_{\vec{x} \in \mathbb{L}} M_{\vec{x}}$ such that $S_{\vec{x}} M_{\vec{y}} \subseteq M_{\vec{x}+\vec{y}}$, for all \vec{y} and \vec{y} from \mathbb{L} . It follows that all components $M_{\vec{x}}$ are finite-dimensional over k. Equipped with the degree-preserving morphisms (morphism of degree zero), the category $\mathrm{mod}^{\mathbb{L}}$ -S is Hom-finite, that is, it is a k-linear abelian category with finite dimensional Hom-spaces.

An important feature of this category is the **action of the grading group** \mathbb{L} **by shift**: if $M = \bigoplus_{\vec{x} \in \mathbb{L}} M_{\vec{x}}$ is a graded S-module and $\vec{y} \in \mathbb{L}$, then we define $M(\vec{y})$ to be the graded module with $M(\vec{y})_{\vec{x}} = M_{\vec{x}+\vec{y}}$. In particular, each indecomposable projective object in $\text{mod}^{\mathbb{L}}$ -S has the form of the module $S(\vec{x})$ with $\vec{x} \in \mathbb{L}$. Viewed from a graded point of view, the algebra S is **graded-local**², that is, it has a unique maximal graded ideal $\mathbf{m} = (x_1, x_2, x_3)$. Accordingly, $S/\mathbf{m} = k$ is simple in $\text{mod}^{\mathbb{L}}$ -S; moreover, each simple graded S-module has the form $k(\vec{x})$ for a unique \vec{x} in \mathbb{L} . We conclude that a graded S-module has finite length if and only if it is finite dimensional. We denote by $\text{mod}_0^{\mathbb{L}}$ -S the full subcategory of $\text{mod}^{\mathbb{L}}$ -S consisting of all finite length objects. It is a **Serre subcategory**, that is, it is closed under subobjects, factor objects and extensions.

²See the previous remarks for how to interpret concepts in the graded sense!

The setting allows us to deal with an L-grading variant of the socalled **Serre construction** [28], going back to Serre (1955). This is done by forming the **quotient category**

$$\mathcal{H} = \frac{\operatorname{mod}^{\mathbb{L}} - S}{\operatorname{mod}_{0}^{\mathbb{L}} - S},$$

which is again an abelian category defined as follows (for details of the construction we refer to [6]):

- the objects of \mathcal{H} are just the objects of $\operatorname{mod}^{\mathbb{L}}$ -S,
- the morphisms of \mathcal{H} are obtained from the morphisms of $\operatorname{mod}^{\mathbb{L}}$ -S by **formally inverting** all morphisms having a kernel and a cokernel of finite length,
- the composition in \mathcal{H} is induced by the composition in $\operatorname{mod}^{\mathbb{L}}$ -S.

3.4. Coherent sheaves on a weighted projective line. The category \mathcal{H} has an interpretation as the category of coherent sheaves on a weighted projective line \mathbb{X} having three weighted points of weights (p_1, p_2, p_3) . For this reason, we are to some extent using sheaf-theoretic language for concepts related to \mathcal{H} : in the above setting, we have a natural quotient functor $q: \mathrm{mod}^{\mathbb{L}} - S \to \mathcal{H}$, which is exact. Further more, the \mathbb{L} -action on $\mathrm{mod}^{\mathbb{L}} - S$ induces an \mathbb{L} -action on \mathcal{H} , which we also denote in shift notation by $(\vec{x}, X) \mapsto X(\vec{x})$. Moreover, we use the notation $\mathcal{O} = q(S)$ and, for reasons becoming transparent later, call this the structure sheaf. Moreover, we call $q(S(\vec{x}) = \mathcal{O}(\vec{x}))$ to be a twisted structure sheaf. These form a nice set of 'generators' for the category \mathcal{H} . We have the following result, see [7] for further details. For an alternative access to weighted projective lines, we refer to [3, 20].

Theorem 3.2 (Geigle-Lenzing, 1987). The category \mathcal{H} has the following properties:

- (1) \mathcal{H} is a Hom-finite abelian category which is **noetherian**, that is, any ascending chain of subobjects becomes stationary.
- (2) H satisfies the **Serre duality** in the form

$$D \operatorname{Ext}^1(X, Y) = \operatorname{Hom}(Y, X(\vec{\omega})).$$

(This implies that the category \mathcal{H} is hereditary and further that it has almost-split sequence with the Auslander-Reiten translation τ , given by twist with $\vec{\omega}$.)

(3) The indecomposable objects from \mathcal{H} come in two parts:

$$\mathcal{H}_0 = \{X \in \mathcal{H} | \operatorname{length}(X) < \infty\},\$$

 $\mathcal{H}_+ = \{Y \in \mathcal{H} | Y \text{ has no simple subobject}\}.$

Moreover, $\operatorname{Hom}(\mathcal{H}_0, \mathcal{H}_+) = 0$. Members of \mathcal{H}_0 will be called **torsion** (or **finite length**) **sheaves**, and those of \mathcal{H}_+ will be called **vector bundles**.

- (4) We have natural isomorphisms $\operatorname{Hom}(\mathcal{O}(\vec{x}), \mathcal{O}(\vec{y})) = S_{\vec{y}-\vec{x}}$.
- (5) There is a \mathbb{Z} -linear form on $K_0(\mathcal{H})$, called rank, which is 0 exactly on the objects of \mathcal{H}_0 and greater than 0 otherwise.
- (6) For each **line bundle** L, that is, an indecomposable object of rank one, there exists a unique \vec{x} from \mathbb{L} such that L is isomorphic to $\mathcal{O}(\vec{x})$.
- (7) The indecomposables of \mathcal{H}_0 decompose into a $\mathbf{P}^1(k)$ -family of uniserial (standard stable) tubes with three distinguished ones having p_1, p_2, p_3 simple objects, respectively, and the remaining ones containing exactly one simple object.

Proof. We give a few indications concerning the proof.

- Ad (1): Abelianness is a general feature of the quotient category with respect to a Serre subcategory. As is easy to see, noetherianness of $\text{mod}^{\mathbb{L}}$ -S is preserved when passing to the quotient category.
- Ad (2): This is technically the most difficult part. On the other hand, it is a general technique in algebraic geometry. If one deals with a graded complete intersection S having a minimal set of homogeneous generators in degrees a_1, \ldots, a_n and a minimal set of homogeneous relations in degrees b_1, \ldots, b_m , then one gets the Serre duality in the form $D \operatorname{Ext}^n(X,Y) = \operatorname{Ext}^{n-d}(Y,X(\omega))$, where d=n-m-1 and $\omega = \sum_{i=1}^m d_i \sum_{j=1}^n a_i$. The techniques use either a Koszul complex associated with the complete intersection or alternatively a minimal graded resolution of S. For details, we refer to the literature.
- Ad (3): Let X be an object in \mathcal{H} . By noetherianness, X has a largest noetherian subobject X_0 of finite length such that we obtain a short exact sequence η : $0 \to X_0 \to X \to X/X_0 \to 0$ with X_0 from \mathcal{H}_0 and X/X_0 from \mathcal{H}_+ . Invoking the Serre duality, one now shows that η splits, which yields the result.
- Ad (4) and (6): This is another important feature following directly from graded factoriality of S.
- Ad (5): A quick way to define the rank is the following. Let \mathcal{H}_0 denote the full subcategory of \mathcal{H} consisting of all objects of finite length. Then,

the quotient category $\mathcal{H}/\mathcal{H}_0$ is an abelian category, where each object has a finite length. For X in \mathcal{H} , now define the rank of X as the length of X in $\mathcal{H}/\mathcal{H}_0$. It is then easy to verify the claim.

Ad (7): This uses the classification of homogeneous prime elements in the algebra S. If p is a homogeneous prime in S, then $q(S/(p))(\vec{x})$ is a simple object in \mathcal{H} , and each simple object U has this form. Here p is uniquely determined by S, while \vec{x} is not.

Corollary 3.3. We can recover the weight triple (p_1, p_2, p_3) , and hence the \mathbb{L} -graded algebra S from the category \mathcal{H} .

Proof. This follows directly from part (7) in Theorem 3.2.

Corollary 3.4. The companion category $[\mathbb{L}; S]$ of the \mathbb{L} -graded algebra S is equivalent to the full subcategory \mathcal{L} of coh- \mathbb{X} , which is formed by all the line bundles on \mathbb{X} .

Proof. This is an immediate consequence of (4) and (6) in Theorem 3.2.

4. Link to algebras and the Cohen-Macaulay modules

- 4.1. Singularities and finite dimensional algebras. Summarizing the present status, we have applied the Serre construction to the L-graded algebra S and obtained the category $\mathcal{H} = \text{coh-}\mathbb{X}$, which is an abelian, Hom-finite k-linear category which is Krull-Schmidt and which has almost-split sequences. So, \mathcal{H} is already quite close to the features of a category of finite dimensional modules. This relationship is not only a formal one, but it can also be made very explicit, since \mathcal{H} also has a tilting object. (In fact, it does have plenty of them!) Since \mathcal{H} is hereditary, it is convenient to say that an object T of \mathcal{H} is a **tilting object** if the following two conditions are satisfied:
 - (a) T has no self-extensions, that is, $\operatorname{Ext}^1_{\mathcal{H}}(T,T) = 0$;
 - (b) T generates \mathcal{H} homologically, that is, whenever $X \in \mathcal{H}$ satisfies $\operatorname{Hom}_{\mathcal{H}}(T,X) = 0 = \operatorname{Ext}^1_{\mathcal{H}}(T,X)$, then X = 0.

Theorem 4.1. Assume $X = X(p_1, p_2, p_3)$ is the weighted projective line given by the weight triple (p_1, p_2, p_3) . Then, the object

$$T = \bigoplus_{0 \le \vec{x} \le \vec{c}} \mathcal{O}(\vec{x})$$

is a tilting object in coh-X, whose endomorphism ring $\Lambda = \operatorname{End}_{\mathcal{H}}(T)$ is the **canonical algebra** given by the same weight triple; we write $\Lambda = \Lambda(p_1, p_2, p_3)$. That is, Λ is given by the quiver

with three respective arms of lengths p_1, p_2, p_3 and the single relation $x_1^{p_1} + x_2^{p_2} + x_3^{p_3} = 0$.

Proof. That T has no self-extensions, it uses the Serre duality combined with the formula $\operatorname{Hom}(\mathcal{O}(\vec{x}),\mathcal{O}(\vec{y})) = S_{\vec{y}-\vec{x}}$. To show that T generates \mathcal{H} homologically, the key point is to show that for each simple object S in \mathcal{H} at least one of the $\mathcal{O}(\vec{x})$ with \vec{x} in the range $0 \leq \vec{x} \leq \vec{c}$ admits a non-zero homomorphism to S.

Corollary 4.2. The bounded derived categories of $D^b(\text{mod }\Lambda)$ and $D^b(\mathcal{H})$ are triangle-equivalent.

This result allows a number of strong consequences, since the abelian category \mathcal{H} is hereditary, and hence the bounded derived category of \mathcal{H} can be identified with the **repetitive category** of \mathcal{H} . Recall that the repetitive category is the additive closure of the disjoint union

$$\bigvee_{n\in\mathbb{Z}}\mathcal{H}[n],\quad\text{ where each }\mathcal{H}[n]\text{ is a copy of }\mathcal{H},$$

with objects of $\mathcal{H}[n]$ written as X[n], and where morphism are given by

$$\operatorname{Hom}(X[m], Y[n]) = \operatorname{Ext}_{\mathcal{H}}^{n-m}(X, Y).$$

A particular consequence of this setting follows here.

Corollary 4.3. The category mod Λ of modules over the canonical algebra Λ is equivalent to (the additive closure of) the union of

$$\{X \in \mathcal{H} \mid \operatorname{Ext}^1_{\mathcal{H}}(T, X) = 0\} \vee \{Y[1] \mid Y \in \mathcal{H} \text{ with } \operatorname{Hom}_{\mathcal{H}}(T, Y) = 0\}$$


viewed as a full subcategory of $\mathcal{H} \vee \mathcal{H}[1] \subset D^b(\mathcal{H})$.

Remark 4.4. We discuss briefly the relationship between the categories \mathcal{H} and mod Λ as far it is relevant for the matter of singularities.

- As Corollary 4.3 states, the category H contains all the information on the category of Λ-modules (via the repetitive category of H). In particular, the representation type of H determines the representation type of Λ.
- (2) The complexity of the classification problem for coh- $\mathbb{X} = \mathcal{H}$ is completely determined by the numerical invariant $\delta(\vec{\omega})$
 - $\bar{p}(1-(1/p_1+1/p_2+1/p_3))$. Since indecomposables of \mathcal{H}_0 are explicitly classified by means of 1-parameter families, indexed by the projective line, the complexity is determined by the category vect- $\mathbb{X} = \mathcal{H}_+$ of vector bundles on \mathbb{X} .
 - (a) If $\delta(\vec{\omega}) < 0$, then the Auslander-Reiten quiver for the indecomposable vector bundles consists of a single component of shape $\mathbb{Z}\tilde{\Delta}$, where $\tilde{\Delta}$ is the extended Dynkin diagram corresponding to $\Delta = [p_1, p_2, p_3]$.
 - (b) If $\delta(\vec{\omega}) = 0$, then the classification problem for coh- \mathbb{X} is still tame (but complicated). The indecomposable vector bundles decompose into a rational family $(\mathcal{T}_q)_{q \in \mathbb{Q}}$, where, in turn, each \mathcal{T}_q is a $\mathbf{P}^1(k)$ -family of tubes, each one being of tubular type (p_1, p_2, p_3) . We express this by saying that coh- \mathbb{X} , (correspondingly Λ) has tubular type. Note that tubular algebras play an important role in tame representation theory.
 - (c) If $\delta(\vec{\omega}) > 0$, then we deal with a wild situation. Here, all AR-components for vect- \mathbb{X} are of type $\mathbb{Z}\mathbb{A}_{\infty}$. Moreover, there is a natural bijection between the set of all such components to the set of all AR-components of regular modules for the path algebra C of the star $[p_1, p_2, p_3]$, endowed with an arbitrary orientation; see [22]. Note that in this case the algebra C is of wild representation type.
- 4.2. Shape of the category of vector bundles. For the moment, our main interest is in the case $\delta(\vec{\omega}) < 0$. We illustrate the situation by an example.

Example 4.5. For the weight type (2,3,4), the corresponding Dynkin diagram is $\mathbb{E}_7 = [2,3,4]$. Here, the Auslander-Reiten quiver for vect-X

is given by $\mathbb{Z}\tilde{\mathbb{E}}_7$, and therefore looks as follows:



Note that the line bundles form two τ -orbits sitting at the border of the component. We have marked the corresponding vertices by circles \circ while the other vertices are marked by fat dots \bullet . We have 8 τ -orbits whose corresponding orbit graph yields the extended Dynkin diagram $\tilde{\mathbb{E}}_7$. Since the rank, introduced in Theorem 3.2, is constant on τ -orbits, it yields a function on $\tilde{\mathbb{E}}_7$ which turns out to be the unique (normalized) additive function for this extended Dynkin graph. The rank function on vect- \mathbb{X} is thus determined by the following diagram with the attached rank values

$$\begin{smallmatrix} 2 \\ | \\ 1-2-3-4-3-2-1. \end{smallmatrix}$$

We thus rediscover that there are two line bundle components sitting at the border of the AR-quiver.

More is true, also in the other cases of weight triples of Dynkin type $\Delta = [p_1, p_2, p_3]$. Namely, the category vect- \mathbb{X} , not just its Auslander-Reiten quiver is completely determined by the mesh category $\mathbb{Z}\tilde{\Delta}$, since the path category of this mesh category is equivalent to the category of indecomposable vector bundles on \mathbb{X} . This can be derived from the fact that coh- \mathbb{X} has a tilting bundle T whose endomorphism ring is the path algebra $k\tilde{\Delta}$ of the path algebra of an extended Dynkin quiver of type $\tilde{\Delta}$. For the case [2,3,4], we have depicted such a tilting bundle above by framing the vertices of a section in the Auslander-Reiten quiver.

Concerning the position of a suitable tilting object, we can be more specific. We have already defined the rank which is constant on τ -orbits and which is 0 on finite length sheaves and greater than 0 on non-zero vector bundles. There is another \mathbb{Z} -linear form with somewhat complementary properties, the **degree**. The degree is greater than 0

on objects of \mathcal{H}_0 and there it is also constant on τ -orbits. Moreover, it vanishes on \mathcal{O} . Each non-zero vector bundle X then has a well-defined **slope** given by $\mu X = \deg X/\operatorname{rk} F$. The following result is due to Hübner (unpublished), and a proof can be found in [23].

Theorem 4.6. Assume the weight triple (p_1, p_2, p_3) satisfies $\delta(\vec{\omega}) < 0$. Let $\Delta = [p_1, p_2, p_3]$ be the corresponding Dynkin diagram. Then, there is only a finite system T_1, \ldots, T_n of pairwise nonisomorphic indecomposable vector bundles E with slope in the range $0 \le \mu E < -\delta(\vec{\omega})$. Moreover, $T = T_1 \oplus \cdots T_n$ is a tilting object whose endomorphism algebra is isomorphic to the path algebra kQ of a quiver Q whose underlying graph is the extended Dynkin diagram $\tilde{\Delta}$.

If all the three weights are bigger than 2, then (as in the preceding example) Q has bipartite orientation.

Corollary 4.7. The path algebra kQ of a quiver Q without oriented cycles is derived equivalent to a canonical algebra if and only if the graph underlying Q is extended Dynkin.

4.3. From singularities to weights. After this digression on some representation-theoretic link of singularity theory, we come back to our main subject.

Theorem 4.8. We assume a weight triple (p_1, p_2, p_3) with $\delta(\vec{\omega})$ different from zero. Let $R = S_{|\mathbb{Z}\vec{\omega}}$, considered as a \mathbb{Z} -graded k-algebra (with $\vec{\omega} \leftrightarrow -1$ for $\delta(\vec{\omega}) < 0$ and $\vec{\omega} \leftrightarrow 1$ for $\delta(\vec{\omega}) > 0$). Then, the restriction functor res: $\text{mod}^{\mathbb{L}}$ - $S \to \text{mod}^{\mathbb{Z}}$ -R, $M \mapsto M_{|\mathbb{Z}\vec{\omega}}$ induces an equivalence

$$\mathcal{H} = \frac{\operatorname{mod}^{\mathbb{L}} - S}{\operatorname{mod}_{\mathbb{Q}}^{\mathbb{L}} - S} \xrightarrow{\sim} \frac{\operatorname{mod}^{\mathbb{Z}} - R}{\operatorname{mod}_{\mathbb{Q}}^{\mathbb{Z}} - R}.$$

Proof. We first observe that the restriction of \mathbb{L} -graded S-modules to \mathbb{Z} -graded R-modules preserves finite length, and thus induces a restriction functor for the two quotient categories. The main ingredients of the proof then are the following two facts:

- (1) For each simple object E in \mathcal{H} , the image is non-zero (and then also simple).
- (2) Each finitely generated $\mathbb{Z}\vec{\omega}$ -graded R-module M extends to a finitely generated \mathbb{L} -graded S-module \bar{M} (such that the restriction of \bar{M} to $\mathbb{Z}\vec{\omega}$ equals M). This part of the proof uses the left Kan-extension or, in a different terminology, the graded tensor product $S \otimes_R -$.

With Theorem 4.8 at hand, we have solved our problem to discover the Dynkin diagram from the \mathbb{Z} -graded simple surface singularity f_{Δ} .

Corollary 4.9. Let f_{Δ} be a \mathbb{Z} -graded simple surface singularity f_{Δ} and $R = k[x, y, z]/(f_{\Delta})$. Then, the quotient category $\text{mod}^{\mathbb{Z}}$ - $R/\text{mod}_0^{\mathbb{Z}}$ -R is equivalent to the category of coherent sheaves $\text{coh-}\mathbb{X}$ on the weighted projective line of weight type (p_1, p_2, p_3) , where $\Delta = [p_1, p_2, p_3]$.

As we have seen before, the weight type of \mathbb{X} can be recovered as the tubular type of \mathcal{H} , that is, by determining the τ -periods of the tubes in the AR-quiver of \mathcal{H}_0 .

4.4. The link to the graded Cohen-Macaulay modules. We start with a definition of the graded maximal Cohen-Macaulay modules for graded-local algebras of dimension two (like the \mathbb{L} -graded algebra S or the \mathbb{Z} -graded algebra R).

Definition 4.10. A finitely generated \mathbb{L} -graded S-module M is called (maximal) Cohen-Macaulay if

$$\operatorname{Hom}_S(E, M) = 0 = \operatorname{Ext}_S^1(E, M)$$

holds for each simple \mathbb{L} -graded S-module E. (Recall these are of the form $k(\vec{x})$.) By $CM^{\mathbb{L}}$ -S, we denote the category of all \mathbb{L} -graded CM-modules as a full subcategory of $\mathrm{mod}^{\mathbb{L}}$ -S.

A similar definition applies to \mathbb{Z} -graded R-modules. We remark here that for algebras one always has the implications (hypersurface) \Rightarrow (complete intersection) \Rightarrow (Gorenstein) \Rightarrow (Cohen-Macaulay) in the graded and ungraded sense. Hence, the algebra S is always graded Gorenstein. For $\delta(\vec{\omega}) \neq 0$, one can show the same for the \mathbb{Z} -graded algebra R. It is not so obvious for $\delta(\vec{\omega}) > 0$, while for $\delta(\vec{\omega}) < 0$ it follows from the list of simple graded singularities, which all are hypersurfaces.

Theorem 4.11. Let (p_1, p_2, p_3) be a weight triple.

(a) If $q: \operatorname{mod}^{\mathbb{L}} - S \to \operatorname{mod}^{\mathbb{L}} - S/\operatorname{mod}^{\mathbb{Z}} - S = \operatorname{coh-}\mathbb{X}$ denotes the natural quotient functor, then q induces an equivalence $q: \operatorname{CM}^{\mathbb{L}} - S \xrightarrow{\sim} \operatorname{vect-}\mathbb{X}$. This equivalence sends the indecomposable projective $S(\vec{x})$ to $\mathcal{O}(\vec{x})$ and induces an equivalence between the category $\operatorname{proj}^{\mathbb{L}} - S$ of finitely generated \mathbb{L} -graded projective S-modules and the full subcategory \mathcal{L} of $\operatorname{vect-}\mathbb{X}$, consisting of all line bundles.

(b) If we exclude the tubular weights, then the restriction functor from \mathbb{L} -graded S-modules to \mathbb{Z} -graded R-modules induces an equivalence

res :
$$CM^{\mathbb{L}} - S \xrightarrow{\sim} CM^{\mathbb{Z}} - R$$
, $M \mapsto M_{|\mathbb{Z}\vec{\omega}}$.

This equivalence sends the indecomposable projective R(n) to $\mathcal{O}(-n\vec{\omega})$ if $\delta(\vec{\omega}) < 0$, respectively to $\mathcal{O}(n\vec{\omega})$ if $\delta(\vec{\omega}) > 0$, and thus induces an equivalence between $\operatorname{proj}^{\mathbb{Z}}$ -R and the τ -orbit $\tau^{\mathbb{Z}}\mathcal{O}$.

Proof. The first assertion (a) follows from the existence of an inverse $\Gamma: \text{vect-}\mathbb{X} \to \text{CM}^{\mathbb{L}}\text{-}S$ to q, where $\Gamma(E) = \bigoplus_{\vec{x} \in \mathbb{L}} \text{Hom}_{\mathbb{X}}(\mathcal{O}(-\vec{x}), E)$. The point here is to prove that $\Gamma(M)$ is finitely generated over S which uses that E is a vector bundle. The first part of (b) then follows from Theorem 4.8. The remaining assertions in (a) and (b) are obvious. \square

The combination of Theorem 4.11 with Remark 4.4 immediately yields the following theorem.

Theorem 4.12. Assume $[p_1, p_2, p_3]$ is a Dynkin diagram. With the notation introduced previously, the k-linear categories

vect-
$$\mathbb{X}$$
, $\mathrm{CM}^{\mathbb{L}}$ - S and $\mathrm{CM}^{\mathbb{Z}}$ - R


can be naturally identified. Their Auslander-Reiten quiver forms a single component of shape $\mathbb{Z}\tilde{\Delta} = \{(n,v) \mid n \in \mathbb{Z}, v \in \tilde{\Delta}_0\}.$

Moreover, in this component the indecomposable \mathbb{Z} -graded projective R-modules form a single τ -orbit lying at the boundary of the component if all the weights are ≥ 2 .

As an immediate consequence we obtain the information on the shape of the Auslander-Reiten quiver of (maximal) CM-modules over the complete simple surface singularities.

Corollary 4.13. Assume $\Delta = [p_1, p_2, p_3]$ is Dynkin and f_{Δ} is the corresponding singularity. Then, the Auslander-Reiten quiver of $\hat{R} = k[[x, y, z]]/(f_{\Delta})$ is obtained from the extended Dynkin diagram $\tilde{\Delta}$ by replacing each edge $\circ - \circ$ by a 2-cycle $\circ \rightleftharpoons \circ$.

For instance, the Dynkin diagram \mathbb{E}_8 with the corresponding singularity $f_{\Delta} = z^2 + y^3 + x^5$ yields a category $\mathrm{CM}(k[[x,y,z]]/(f_{\Delta}))$ with Auslander-Reiten quiver



We have marked by \circ the indecomposable projective module \hat{R} .

Proof of Corollary 4.13. The proof uses the completion functor $\hat{}$: $CM^{\mathbb{Z}}$ - $R \to CM(\hat{R})$, $M \mapsto \prod_{n \in \mathbb{Z}} M_n$ as studied by Auslander-Reiten in [1]. In the present setting, completion preserves indecomposability and almost-split sequences; moreover, two indecomposable graded modules have the same image if and only if they belong to the same τ -orbit of \mathbb{Z} -graded CM-modules. Hence, the image of the completion functor is a finite component of $CM(\hat{R})$. By a Brauer-Thrall type argument, it then follows that the functor is dense. This proves the claim.

Corollary 4.14. Let k be a field. Then, the k-algebra $\hat{R} = k[[x, y, z]]/(x^2 + y^3 + z^5)$ is a factorial domain, that is, \hat{R} is a domain and each non-zero element is a product of prime elements.

Proof. Since completion preserves the rank, it follows that \hat{R} is the only CM-module over \hat{R} having rank one. It is a well-known fact that, in the present setting, this property implies factoriality of \hat{R} .

Corollary 4.15. Let Q be an extended Dynkin quiver associated with the Dynkin diagram Δ ; we fix a vertex v of Q, where 'the' additive function for the graph underlying Q attains value 1. Denote by P the indecomposable projective kQ-module corresponding to the vertex v. Then, the orbit algebra

$$\mathbb{A}(\tau^-, P) := \bigoplus_{n > 0} \operatorname{Hom}_{kQ}(P, \tau^{-n}P)$$

with multiplication $u_n \cdot v_m = \tau^{-m} u_n \circ v_m$ is a positively \mathbb{Z} -graded algebra which is isomorphic to the graded simple surface singularity $k[x, y, z]/(f_{\Delta})$.

Proof. Using the identifications of the theorem, we may identify $\mathbb{A}(\tau^-, P)$ with the orbit algebra of the \mathbb{Z} -graded R-module R with regard to the grading shift $M \mapsto M(1)$, which obviously brings us back to the \mathbb{Z} -graded algebra R.

Remark 4.16. From a general perspective, the last result is quite remarkable. It tells us that a study of the (graded) simple singularities is equivalent to the analysis of the Auslander-Reiten translation for path algebras of an extended Dynkin quiver, or alternatively for the category of coherent sheaves on a weighted projective line whose weight triple determines the Dynkin diagram. Thus, the simple singularities can be

thought of as mathematical objects capturing the homological information on either category, there given by either Auslander-Reiten or Serre duality.

A K-theoretic shadow of this is contained in the list of simple graded singularities, where we have seen in Comments 2.10 that the degree of the singularity f_{Δ} agrees with the Coxeter number h_{Δ} .

5. Vector bundles as a Frobenius category

Now, we are dealing with a recent joint work with de la Peña [21], and Kussin and Meltzer [17]; see also [16]. In the previous section, we have seen in Theorem 4.8 that for each non-tubular weight triple (p_1, p_2, p_3) we have a commutative diagram

$$\begin{array}{c} \operatorname{CM}^{\mathbb{L}}\text{-}S \xrightarrow{q} \operatorname{vect}\text{-}\mathbb{X} \\ \cong \operatorname{|res} & \| \\ \operatorname{CM}^{\mathbb{Z}}\text{-}R \xrightarrow{q'} \operatorname{vect}\text{-}\mathbb{X} \end{array}$$

where, q and q' are equivalences induced by the natural quotient functors. Since S and R are graded Gorenstein, each of the categories $\mathrm{CM}^{\mathbb{L}}$ -S and $\mathrm{CM}^{\mathbb{Z}}$ -R inherits an exact structure from the ambient abelian categories of finitely generated graded modules $\mathrm{mod}^{\mathbb{L}}$ -S and $\mathrm{mod}^{\mathbb{Z}}$ -R, respectively, which turns the two categories of graded CM-modules into Frobenius categories have the category of indecomposable graded projective modules as their indecomposable projective-injective objects. By transport of structure, we thus obtain on vect- \mathbb{X} two, usually different, structures of Frobenius categories.

In more detail, we arrive at the following setting

- (a) From the \mathbb{L} -graded setting, we obtain that vect- \mathbb{X} is a Frobenius category with the system \mathcal{L} of line bundles being the indecomposable projective-injective objects.
- (b) In the non-tubular case, we also obtain from the \mathbb{Z} -graded setting that vect- \mathbb{X} is a Frobenius category with the τ -orbit $\tau^{\mathbb{Z}}\mathcal{O}$ of the structure sheaf, that is, a single τ -orbit of line bundles, being the indecomposable projective-injective objects.

Explanations 5.1.

(1) A Frobenius category is defined to be an exact category which has sufficiently many (relative) projective and (relative) injective objects and where the projectives coincide with the injectives.

- (2) The term exact category is used here in the sense of Quillen. An exact category C, by definition, admits a full embedding as an extension-closed subcategory into an abelian category A. The exact structure on C then is induced from A consisting of all short exact sequences in A with all their terms in C.
- (3) More concretely, a sequence $\eta: 0 \to A \to B \to C \to 0$ in vect- \mathbb{X} is exact with regard to the exact structure (a) if and only if for each line bundle L the sequence $\operatorname{Hom}(L,\eta)$ is exact. The Serre duality then implies that it is equivalent to request exactness of $\operatorname{Hom}(\eta,L)$ for each line bundle L. By contrast, in case (b) the sequence η is exact if and only if $\operatorname{Hom}(\tau^n\mathcal{O},\eta)$, equivalently $\operatorname{Hom}(\eta,\tau^n\mathcal{O})$, is exact for each integer n.
 - (4) By a result due to Happel [8], the associated stable categories

$$\operatorname{vect-X}/[\mathcal{L}]$$
 and $\operatorname{vect-X}/[\tau^{\mathbb{Z}}\mathcal{O}]$

are triangulated. Here, a notation like vect- $\mathbb{X}/[\mathcal{L}]$ means the factor category of vect- \mathbb{X} modulo the ideal generated by \mathcal{L} . In more detail, this is the category having the same objects as vect- \mathbb{X} with morphisms given by the quotient $\underline{\mathrm{Hom}}(X,Y) = \mathrm{Hom}(X,Y)/\{u: X \to Y \mid u \text{ factors through an object of } \mathrm{add}(\mathcal{L})\}.$

(5) The stable categories (a) vect- $\mathbb{X}/[\mathcal{L}]$ and (b) vect- $\mathbb{X}/[\tau^{\mathbb{Z}}\mathcal{O}]$ are triangulated categories with the **Serre duality** induced from the Serre duality of coh- \mathbb{X} . In particular, the categories (a) and (b) have almost-split triangles and the Auslander-Reiten translation is induced from the Auslander-Reiten translation of coh- \mathbb{X} .

It follows that the Auslander-Reiten quiver for (a) and (b) is obtained from the Auslander-Reiten quiver of vect- \mathbb{X} in case (a) by deleting all orbits of line bundles and in case (b), assuming non-tubular type, by deleting just a single τ -orbit of line bundles.

(6) The stable categories of vector bundles (type (a) or (b)) will always have a tilting object. For $\delta(\vec{\omega}) > 0$, this will be a highly non-trivial matter. We are going to return to this aspect later.

Here, we mainly concentrate on the stable category vect- $\mathbb{X}/[\tau^{\mathbb{Z}}\mathcal{O}]$.

Remark 5.2. We assume a non-tubular weight triple. What is then the role of vect- $\mathbb{X}/[\tau^{\mathbb{Z}}\mathcal{O}]$, equivalently of $\underline{\mathrm{CM}}^{\mathbb{Z}}$ - $R = \mathrm{CM}^{\mathbb{Z}}$ - $R/[\mathrm{proj}^{\mathbb{Z}}-R]$? By old work of Buchweitz (1986), see [2], revived by Orlov in 2005,

see [24], 'this' stable category is a measure for the complexity of the graded singularity R. It is equivalent moreover to the triangulated category of graded singularities of R defined as $D_{sing}^{\mathbb{Z}}(R) = D^b(\text{mod}^{\mathbb{Z}}-R)/D^b(\text{proj}^{\mathbb{Z}}-R)$. For instance, the polynomial algebra R = k[x,y], with x and y homogeneous of positive degree, becomes graded-regular-local, yielding $CM^{\mathbb{Z}}-R=0$.

Because of the canonical equivalences between the categories vect- $\mathbb{X}/\tau^{\mathbb{Z}}\mathcal{O}$], $\underline{\mathrm{CM}}^{\mathbb{Z}}$ -R and $\mathrm{D}_{sing}^{\mathbb{Z}}(R)$ it is advisable to think of all three as being incarnations of a single triangulated category. We will encounter further triangulated categories, which are triangle equivalent to the above, but in a non-canonical way.

5.1. The Case $\delta(\vec{\omega}) < 0$ shape of the stable category. Concerning the existence of a tilting object we start with the case $\delta(\vec{\omega}) < 0$.


Theorem 5.3 (Kajiura-Saito-Takahashi, 2006). Assume a weight triple (p_1, p_2, p_3) such that $\Delta = [p_1, p_2, p_3]$ is a Dynkin diagram. Then, the triangulated category vect- $\mathbb{X}/[\tau^{\mathbb{Z}}\mathcal{O}]$ has a tilting object T such that $\operatorname{End}(T)$ is isomorphic to the path algebra $k\vec{\Delta}$ of a quiver $\vec{\Delta}$ with underlying graph Δ . In particular, we have the equivalence,

$$\operatorname{vect-X}/[\tau^{\mathbb{Z}}\mathcal{O}] \cong \operatorname{D}^b(\operatorname{mod} k\vec{\Delta}).$$

Proof. We will give two proofs which is different from the proof in [11] and actually is much shorter. The first proof is inspired by the proof of Theorem 4.6, where we have shown that the direct sum T of a representative system T_1, \ldots, T_n of pairwise indecomposable vector bundles in the slope range $0 \le \mu(E) \le -\delta(\vec{\omega})$ is tilting in coh- \mathbb{X} . Now, observe that the structure sheaf \mathcal{O} belongs to this system. Let's assume that $\mathcal{O} = T_1$. It is then not difficult to check that $\overline{T} = T_2 \oplus \cdots T_n$ is a tilting object in vect- $\mathbb{X}/[\tau^{\mathbb{Z}}\mathcal{O}]$ with endomorphism ring $k\vec{\Delta}$.

We next present another proof that perhaps is providing more insight in the mechanism. By way of illustration, we restrict to the weight type (2,3,4); the other cases can be dealt with in a similar fashion. The relevant facts, we are going to use, are all present in Example 4.5. We recall that the category of indecomposable vector bundles for this weight

type is given by the mesh category $k\mathbb{Z}\tilde{\mathbb{E}}_7$ depicted below:



We depicted the vertices corresponding to the line bundles from the τ -orbit of \mathcal{O} by small circles \circ and all arrows starting or ending in such a vertex are marked by dotted arrows. Passing to the stable category vect- $\mathbb{X}/[\tau^{\mathbb{Z}}\mathcal{O}]$ just kills the orbit $\tau^{\mathbb{Z}}\mathcal{O}$ and the morphisms factor through a finite direct sum of those. On the level of mesh-categories this means to kill the marked vertices and the adjacent arrows, yielding the mesh category of $k\Delta$ of the Dynkin diagram Δ . It is a fundamental result by Happel [8] that the mesh category of $k\Delta$ is equivalent to the bounded derived category $D^b(\text{mod }k\vec{\Delta})$ for any orientation $\vec{\Delta}$ of Δ . It is further well-known that each slice in the AR-quiver $k\mathbb{Z}\Delta$ yields a tilting object in the triangulated category $D^b(\text{mod }k\vec{\Delta})$. In the above picture, we have marked one such tilting object.

5.2. The case $\delta(\vec{\omega}) = 0$.

Comment 5.4. What is going to happen for the tubular weight triples (3,3,3), (2,4,4) and (2,3,6)? In this case, $\mathbb{Z}\vec{\omega}$ is a finite cyclic group and so restriction of S to $\mathbb{Z}\vec{\omega}$ does not yield a \mathbb{Z} -graded algebra. Moreover, it can be shown that the restriction $R = S_{|U}$ of S to any infinite cyclic subgroup U of \mathbb{L} is never \mathbb{Z} -graded Gorenstein; in particular, it will never be generated by three homogeneous elements.

The conclusion from this is that for tubular weight triples it only makes sense to study alternatively the stable category of vector bundles $\mathcal{T} = \text{vect-}\mathbb{X}/[\mathcal{L}]$, where one factors out all line bundles. Here, Ueda [30] shows that \mathcal{T} is triangle-equivalent to the category $D^b(\text{coh-}\mathbb{X})$. Ueda's proof uses an \mathbb{L} -graded version of a recent theorem of Orlov [24]. It is also possible to directly construct a tilting object in \mathcal{T} whose endomorphism algebra is the canonical algebra of the corresponding weight type; see [17].

Ueda's result (actually Orlov's result underlying Ueda's proof) looks paradoxical. Namely, we start with the category coh- \mathbb{X} of coherent sheaves, and then pass to the subcategory vect- \mathbb{X} of vector bundles and in the next step make the category additionally smaller when passing to the stable category $\mathcal{T} = \text{vect-}\mathbb{X}/[\mathcal{L}]$. This, by Ueda's result, is triangle-equivalent to the bounded derived category $D^b(\text{coh-}\mathbb{X})$ which, being equivalent to the repetitive category $\bigvee_{n\in\mathbb{Z}} \text{coh-}\mathbb{X}[n]$, looks much bigger than the category we started with. Note, in this context, that for a tubular weight type the category coh- \mathbb{X} has tame representation type.

5.3. The case $\delta(\vec{\omega}) > 0$, the Arnold's strange duality list. We are now going to discuss what happens with the restriction procedure if we apply it to the weight triples with $\delta(\vec{\omega}) > 0$. The following result is taken from [18] and [21], where additional information is available.

Proposition 5.5. Let k be a field and assume (p_1, p_2, p_3) is a weight triple with $\delta(\vec{\omega}) > 0$. Let $R = S_{|\mathbb{Z}\vec{\omega}}$ be the \mathbb{Z} -graded restriction of the \mathbb{L} -graded triangle singularity S to the subgroup $\mathbb{Z}\vec{\omega}$ which we identify with \mathbb{Z} by the correspondence $\vec{\omega} \leftrightarrow 1$. Then, the followings holds.

- (a) The algebra R is always graded Gorenstein.
- (b) Exactly for the weights triples of Arnold's strange duality list, the algebra R is generated by three homogeneous elements x, y, z and then has the form

$$R = k[x, y, z]/(f),$$

where the generators x, y, z, the relation f and their degrees are given by the list below.

(p_1,p_2,p_3)	generators (x, y, z)	$\deg(x,y,z)$	relation f	$\deg f$	N		
(2, 3, 7)	(x_3, x_2, x_1)	(6, 14, 21)	$z^2 + y^3 + x^7$	42	12		
(2, 3, 8)	(x_3^2, x_2, x_1x_3)	(6, 8, 15)	$z^2 + x^5 + xy^3$	30	13		
(2, 3, 9)	(x_3^3, x_2x_3, x_1)	(6, 8, 9)	$y^3 + xz^2 + x^4$	24	14		
(2,4,5)	$(x_3, x_2^2, x_1 x_2)$	(4, 10, 15)	$z^2 + y^3 + x^5y$	30	11		
(2, 4, 6)	$(x_3^2, x_2^2, x_1 x_2 x_3)$	(4, 6, 11)	$z^2 + x^4y + xy^3$	22	12		
(2, 4, 7)	$(x_3^3, x_2^2x_3, x_1x_2)$	(4, 6, 7)	$y^3 + x^3y + xz^2$	18	13		
(2, 5, 5)	(x_2x_3, x_1, x_2^5)	(4, 5, 10)	$z^2 + y^2z + x^5$	20	12	•	
(2, 5, 6)	$(x_2x_3^2, x_1x_3, x_2^4)$	(4,5,6)	$xz^2 + y^2z + x^4$	16	13		
(3, 3, 4)	$(x_3, x_1 x_2, x_1^3)$	(3, 8, 12)	$z^2 + y^3 + x^4 z$	24	10	•	
(3, 3, 5)	$(x_3^2, x_1x_2, x_3x_1^2)$	(3, 5, 9)	$z^2 + xy^3 + x^3z$	18	11	•	
(3, 3, 6)	$(x_2^3, x_1x_2x_3, x_2^3)$	(3, 5, 6)	$y^3 + x^3z + xz^2$	15	12	•	
(3, 4, 4)	$(x_2x_3, x_1^2, x_1x_2^4)$	(3,4,8)	$z^2 - y^2z + x^4y$	16	11	•	
(3, 4, 5)	$(x_2x_3^3, x_1^2x_3, x_1x_2^3)$	(3,4,5)	$x^3y + xz^2 + y^2z$	13	12		
(4, 4, 4)	$(x_1x_2x_3, x_1^4, x_2^4)$	(3, 4, 4)	$x^4 - yz^2 + y^2z$	12	12	•	
Arnold's strange duality list							

Here, the bullet marks the cases where one has a choice for the monomial generators. Further more, N denotes the sum of the three weights, whose mathematical significance will be revealed later.


Remark 5.6.

(1) For each graded singularity f from Arnold's list, the original weight type can be recovered by the procedure discussed already for the weight triples of the Dynkin type: the Serre construction, when applied to the \mathbb{Z} -graded algebra R = k[x, y, z]/(f), yields back the category coh- \mathbb{X} on the weighted projective line $\mathbb{X}(p_1, p_2, p_3)$, and the tubular type of coh- \mathbb{X} just coincides with (p_1, p_2, p_3) . In the classical context, where $k = \mathbb{C}$, this triple runs under the name of the Dolgachev numbers of f.

(2) For the base field $k = \mathbb{C}$, this list is (equivalent to) Arnold's list of the 14 exceptional unimodular singularities.

This list, slightly extended by the so-called **Gabrielov numbers**, gives rise to what is called **Arnold's strange duality**, which is also related to mirror symmetry. As pointed out before, the weight triples, we are using, will in this context be called the **Dolgachev numbers**. On an ad-hoc basis, we point out that the above list is equipped with an involution, keeping all the weight triples (p_1, p_2, p_2) with $\sum_{i=1}^{3} p_i = 12$ fixed and otherwise sends a weight triple (p_1, p_2, p_3) (the Dolgachev numbers) to the conjugate triple (p'_1, p'_2, p'_3) (the Gabrielov numbers) such that $\sum_{i=1}^{3} p_i + \sum_{i=1}^{3} p'_i = 24$, and moreover the degrees of the relations attached to the two weight triples are identical. We refer to the introductory account of Ebeling [4] for the definition and properties of the Gabrielov numbers.

Our next theorem is taken from a joint work with de la Peña [21]. We point out that the work of Kajiura-Saito-Takahashi [12] is related by subject and results; language and setting are however different. Before stating the result, we need to introduce the concept of an **extended canonical algebra**. By definition, an extended canonical algebra $\bar{\Lambda} = \bar{\Lambda}(p_1, p_2, p_3)$ arises from a canonical algebra $\Lambda = \Lambda(p_1, p_2, p_3)$ by attaching one new arrow (with a new vertex) to a vertex of the quiver of Λ , keeping the relation for Λ , and not introducing any new relation. The algebra given by the quiver below,



having three arms of lengths p_1, p_2, p_3 , respectively, and with the single relation $x_1^{p_1} + x_2^{p_2} + x_3^{p_3} = 0$ is thus an extended canonical algebra $\bar{\Lambda}(p_1, p_2, p_3)$. (We have marked the extension vertex by \star .) Any other attachment of a new arrow and a new vertex would have led to a derived-equivalent algebra, which indicates already some important feature of the extended canonical algebras.

Theorem 5.7. For any weight triple (p_1, p_2, p_3) with $\delta(\vec{\omega}) > 0$ there exists a tilting object T in the stable category vect- $\mathbb{X}/[\tau^{\mathbb{Z}}\mathcal{O}]$ whose endomorphism ring is the extended canonical algebra $\bar{\Lambda}(p_1, p_2, p_3)$. Accordingly, we have equivalences of triangulated categories

$$\underline{\mathrm{CM}}^{\mathbb{Z}} - R = \underline{\mathrm{vect}} - \mathbb{X} \cong \mathrm{D}^b(\mathrm{mod}\,\bar{\Lambda}),$$

and the Grothendieck group $K_0(\underline{\text{vect}}\text{-}\mathbb{X})$ is finitely generated free of rank $\sum_{i=1}^3 p_i$.

Proof. We fix a weight triple (p_1, p_2, p_3) with $\delta(\vec{\omega}) > 0$. Let us say first that we do not know any 'concrete' vector bundle T in vect-X producing a tilting object in the stable category vect-X. Our construction of such a tilting object T is thus done by a theoretical argument using a recent theorem of Orlov: actually we need to apply the proof of Orlov's theorem to the present situation. Since the details are quite technical, we only describe the basic idea of the proof. It follows from our assumption on the weight type and from the construction of R that R is always graded Gorenstein and moreover, the so-called Gorenstein parameter equals -1. Orlov's theorem then states that there is an exceptional object E in vect- \mathbb{X} such that its right perpendicular category, that is, the triangulated subcategory consisting of all objects X such that $\operatorname{Hom}(E[n],X)=0$ for each integer n, is equivalent to $D^b(\text{coh-}\mathbb{X})$. Choosing in coh- \mathbb{X} a tilting object T with endomorphism algebra $\Lambda = \Lambda(p_1, p_2, p_3)$ it then can be shown that the direct sum $\bar{T} = T \oplus E$ is a tilting object in <u>vect-X</u>.

It follows that the Grothendieck group of the triangulated category vect- \mathbb{X} is finitely generated free. Since, moreover, the quiver of the extended canonical algebra has $N = \sum_{i=1}^{3} p_i$ vertices, it follows that the rank of $K_0(\text{vect-}\mathbb{X})$ equals the sum N of the weights.

The theorem has interesting applications; and up to now no other method is known to derive these assertions. We assume throughout that $\delta(\vec{\omega}) > 0$.

Corollary 5.8. Each Auslander-Reiten component in vect- $\mathbb{X} \cong D^b \pmod{\bar{\Lambda}}$ has shape $\mathbb{Z}\mathbb{A}_{\infty}$.

Proof. The corresponding statement is known for the category vect- \mathbb{X} ; see [22]. By a stability argument, all line bundles form AR-orbits belonging to the boundary of $\mathbb{Z}\mathbb{A}_{\infty}$ -components. It follows that after stabilization the components still have shape $\mathbb{Z}\mathbb{A}_{\infty}$.

Corollary 5.9. The set of Auslander-Reiten components of vect- $\mathbb{X} \cong D^b(\text{mod }\overline{\Lambda})$ is in a natural bijection with the set of regular Auslander-Reiten components for any path algebra kQ of a quiver Q with underlying graph $[p_1, p_2, p_3]$.

Proof. For the set of AR-components for vect- \mathbb{X} this is shown in [22]. By the previous argument, stabilization does not change the set of AR-components.

Remark 5.10.

- (1) As for the simple graded singularities, the degrees of the relations f from Arnold's list have an interpretation as the period of the Coxeter transformation for $D^b(\text{mod }\bar{\Lambda})$, and equivalently as the period of the Coxeter transformation for the triangulated category vect- \mathbb{X} . In fact, the two triangulated categories (which are equivalent) are fractional Calabi-Yau, yielding a conceptual reason for the observed periodicity.
- (2) However, the fractional Calabi-Yau property for vect- $\mathbb{X}/[\tau^{\mathbb{Z}}\mathcal{O}]$ is not true for arbitrary weight triples. The weight type (2,3,11) already yields an example.
- (2) Finally, we remark that the stable categories of vector bundles of shape vect- $\mathbb{X}/[\mathcal{L}]$ are in a certain sense much better behaved than those discussed here. The main reason is that they have more symmetry since the Picard group \mathbb{L} acts on them. For instance, one then has tilting

objects for each weight triple, given by an explicit construction. For details in this direction we refer to [17] and [16].

Acknowledgments

The author takes the opportunity to thank the referee for a thorough and critical study of the paper and for numerous helpful comments.

References

- M. Auslander and I. Reiten, Graded modules and their completions, Topics in algebra, Part 1 (Warsaw, 1988), Banach Center Publ. 26 (1990) 181-192.
- [2] R.-O. Buchweitz, Maximal Cohen-Macaulay Modules and Tate-Cohomology over Gorenstein Rings, Preprint, 1986.
- [3] X.-W. Chen and H. Krause, Introduction to Coherent Sheaves on Weighted Projective Lines, Preprint arXiv:0911.4473.
- [4] W. Ebeling, Functions of Several Complex Variables and Their Singularities, Translated from the 2001 German original by Philip G. Spain. Graduate Studies in Mathematics, 83, American Mathematical Society, Providence, RI, 2007.
- [5] W. Ebeling and D. Ploog, McKay correspondence for the Poincaré series of Kleinian and Fuchsian singularities, Math. Ann. 347 (2010) 689-702.
- [6] P. Gabriel, Des catégories abéliennes, (French) Bull. Soc. Math. France 90 (1962) 323-448.
- [7] W. Geigle and H. Lenzing, A class of weighted projective curves arising in representation theory of finite-dimensional algebras, Singularities, representation of algebras, and vector bundles (Lambrecht, 1985), Lecture Notes in Math. 1273, Springer, Berlin, 1987, pp. 265-297.
- [8] D. Happel, Triangulated Categories in the Representation Theory of Finite-Dimensional Algebras, London Mathematical Society Lecture Note Series 119, Cambridge University Press, Cambridge, 1988.
- [9] D. Happel, A characterization of hereditary categories with tilting object, *Invent. Math.* 144 (2001) 381-398.
- [10] M. Hazewinkel, W. Hesselink, D. Siersma and F. D. Veldkamp, The ubiquity of Coxeter-Dynkin diagrams (an introduction to the A-D-E problem), Nieuw Arch. Wisk. (3) **25** (1977) 257-307.
- [11] H. Kajiura, K. Saito and A. Takahashi, Matrix factorization and representations of quivers. II. Type ADE case, Adv. Math. 211 (2007) 327-362.
- [12] H. Kajiura, K. Saito and A. Takahashi, Triangulated categories of matrix factorizations for regular systems of weights with $\epsilon = -1$, Adv. Math. **220** (2009) 1602-1654.
- [13] F. Klein, Vorlesungen Über das Ikosaeder und die Auflösung der Gleichungen vom Fünften Grade, (German) [Lectures on the icosahedron and the solution

- of equations of fifth degree] Reprint of the 1884 original. Edited, with an introduction and commentary by Peter Slodowy. Birkhäuser Verlag, Basel; B. G. Teubner, Stuttgart, 1993.
- [14] D. Kussin, Graded factorial algebras of dimension two, Bull. London Math. Soc. 30 (1998) 123-128.
- [15] D. Kussin, Noncommutative curves of genus zero: related to finite dimensional algebras, *Mem. Amer. Math. Soc.* **201** (2009) x+128 pp.
- [16] D. Kussin, H. Lenzing and H. Meltzer, Nilpotent Operators and Weighted Projective Lines, Preprint arXiv:1002.3797, 2010.
- [17] D. Kussin, H. Lenzing and H. Meltzer, Triangle Singularities, ADE-chains and weighted projective lines, in preparation.
- [18] H. Lenzing, Wild canonical algebras and rings of automorphic forms, Finitedimensional algebras and related topics (Ottawa, ON, 1992), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci. 424 (1994) 191-212.
- [19] H. Lenzing, Representations of finite-dimensional algebras and singularity theory, *Trends in ring theory* (Miskolc, 1996), CMS Conf. Proc. **22**, *Amer. Math. Soc.*, *Providence*, *RI*, 1998, pp. 71-97.
- [20] H. Lenzing, Hereditary Categories, Handbook of Tilting Theory, London Math. Soc. Lecture Note Ser. 332, Cambridge Univ. Press, Cambridge, 2007, pp. 105-146
- [21] H. Lenzing and J. A. de la Peña, Extended Canonical Algebras and Fuchsian Singularities, arXiv:math.RT/0611532Math. Z., to appear.
- [22] H. Lenzing and J. A. de la Peña, Wild canonical algebras, Math. Z. 224 (1997) 403-425.
- [23] H. Lenzing and I. Reiten, Hereditary Noetherian categories of positive Euler characteristic, Math. Z. 254 (2006) 133-171.
- [24] D. Orlov, Derived Categories of Coherent Sheaves and Triangulated Categories of Singularities, In Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, volume 270 of Progr. Math., pages 503-531. Birkhäuser Boston Inc., Boston, MA, 2009.
- [25] H. Poincare, Mémoire sur les fonctions fuchsiennes. (French) $Acta\ Math.\ {\bf 1}\ (1882)$ 193-294.
- [26] C. M. Ringel. Tame Algebras and Integral Quadratic Forms, Lecture Notes in Mathematics. 1099. Springer-Verlag, Berlin, 1984.
- [27] C. M. Ringel, The Canonical Algebras, Topics in algebra, Part 1, Banach Center Publ. 26, PWN, Warsaw, 1990, With an appendix by William Crawley-Boevey, pp. 407-432.
- [28] J.-P. Serre, Faisceaux algébriques cohérents, (French) Ann. of Math. (2) **61** (1955) 197-278.
- [29] P. Slodowy, Platonic solids, Kleinian singularities, and Lie groups, Algebraic geometry (Ann Arbor, Mich., 1981), Lecture Notes in Math. 1008, Springer, Berlin, 1983, pp. 102-138.
- [30] K. Ueda, A remark on a Theorem of math.ag/0511155, arXiv:math/0508258v3 [math.AG], 2006, Preprint.

Helmut Lenzing

Institut für Mathematik, Universität Paderborn, 33095, Paderborn, Germany Email: helmut@math.uni-paderborn.de