

ON CO-NOETHERIAN DIMENSION OF RINGS

A. HAGHANY AND M. R. VEDADI*

Communicated by Siamak Yassemi

ABSTRACT. We define and study co-Noetherian dimension of rings for which the injective envelope of simple modules have finite Krull-dimension. This is a Morita invariant dimension that measures how far the ring is from being co-Noetherian. The co-Noetherian dimension of certain rings, including commutative rings, are determined. It is shown that the class \mathcal{W}_n of rings with co-Noetherian dimension $\leq n$ is closed under homomorphic images and finite normalizing extensions, and that for each n there exist rings with co-Noetherian dimension n . The possible relations between Krull and co-Noetherian dimensions are investigated, and examples are provided to show that these dimensions are independent of each other.

1. Introduction and Preliminaries

Throughout rings have nonzero identity elements and modules are unitary. Unless otherwise stated, we shall consider right hand properties, e.g., right modules, right Noetherian, etc. The category of all right R -modules is denoted by $\text{Mod-}R$. Dualizing the notion of finitely generated (f.g.), Vamos [12] defined *finitely embedded (f.e.)* module M_R by the condition $E(M) = \bigoplus_{i=1}^n E(S_i)$ where each S_i is simple. In [8], Jans called a ring R *co-Noetherian* if factors of f.e. modules are f.e., and proved that R is co-Noetherian if and only if all f.e. modules are Artinian. More

MSC(2010): Primary: 16D10; Secondary: 16D90, 16P50

Keywords: Co-Noetherian, finitely cogenerated, Krull dimension, normalizing extension.

Received: 5 March 2010, Accepted: 7 September 2010

*Corresponding author

© 2012 Iranian Mathematical Society.

recently the term finitely embedded is replaced by *finitely cogenerated*, and the books [2, 13] contain detailed accounts of this notion. We shall define in §2, the co-Noetherian dimension of a ring R in terms of the Krull dimension of finitely cogenerated R -modules. A useful characterization of co-Noetherian rings states that R is co-Noetherian if and only if for every M_R , the intersection of submodules N with M/N Artinian, is zero; see [8] and [7]. Our first result, Theorem 2.1 (stated below) is a generalization of this fact. Hirano in [7, Theorem 2.2], proved that if R is co-Noetherian and S is a finite normalizing extension of R , then S is also co-Noetherian. This result is generalized in Theorem 2.2 for rings of higher co-Noetherian dimension. We use formal triangular matrix rings to prove in Theorem 2.4 that for a given positive integer n there exists a ring R with co-Noetherian dimension n . Co-Noetherian dimension of certain rings is investigated in Proposition 2.6 and, for a commutative ring R , it is shown in Theorem 2.5 that the co-Noetherian dimension (when it exists) is equal to $\text{Sup}_{P \in X} \{ \text{K.dim}(R/P) \}$ where X is a suitable subset of $\text{Spec}(R)$. In the final part of the paper the existence of rings with infinite co-Noetherian dimension is shown and, by way of examples, some possible relations which may exist between co-Noetherian dimension and Krull dimension of rings are illustrated. We now state some terminology that will be used in the paper, and the reader is referred to [10] and [11] which deal in depth with Krull dimension. However, note the following:

Standing hypothesis. Throughout the paper we shall only deal with modules that have finite Krull dimension, and the symbol “ K.dim ” will stand for Krull dimension. When we write $\text{K.dim}(M)$, it is tacitly assumed that M does have Krull dimension which is n for some non-negative integer n .

For each non-negative integer n , and $M \in \text{Mod-}R$, let $\tau_n(M_R) = \cap N$, where the intersection runs through the set of R -submodules N of M with $\text{K.dim}(M/N) \leq n$. Let \mathcal{W}_n denote the class of rings R with the property that *all finitely cogenerated R -modules have Krull dimension at most n* . Thus \mathcal{W}_0 is the class of co-Noetherian rings.

For M_R let $\tau(M_R) = \cap N$, where the intersection runs through the set of R -submodules N such that M/N has Krull dimension. Moreover, \mathcal{W} will denote the class of rings R such that $E(S_R)$ has Krull dimension for all simple modules $S \in \text{Mod-}R$. If there is no confusion, we simply write

$\tau_n(M)$ (respectively $\tau(M)$) instead of $\tau_n(M_R)$ (respectively $\tau(M_R)$).

Next, we present some facts on formal triangular matrix rings for later use.

Let A and B be rings, ${}_B M_A$ a bimodule and $R = \begin{bmatrix} A & 0 \\ M & B \end{bmatrix}$ the formal triangular matrix ring. We apply the equivalence of category $\text{Mod-}R$ with the category of triples $(X, Y)_f$, and use methods of [4]. Thus if V is a right R -module it is often identified with the triple $(Ve_1, Ve_2)_f$ where $(Ve_1)_A, (Ve_2)_B$ are obtained from the idempotents $e_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$; $e_2 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$. Here, f is an A -linear map $:Ve_2 \otimes_B M \rightarrow Ve_1$, given by the rule $f(ve_2 \otimes m) = ve_2 \begin{bmatrix} 0 & 0 \\ m & 0 \end{bmatrix} = ve_2 \begin{bmatrix} 0 & 0 \\ m & 0 \end{bmatrix} e_1$. Hence, it is easily verified (or use the description of maximal right ideals of R in [6]) that if V_R is simple then either $V = (X, 0)_0$ or $V = (0, Y)_0$ where X_A and Y_B are simple. We are interested in triples that can represent an injective envelope of the simple module V_R . In fact, if $V = (0, Y)_0$ then $E(V) = (0, E(Y))_0$, and if $V = (X, 0)_0$, then $E(V) = [E(X), \text{Hom}_A(M, E(X))]_\delta$ where $\delta : \text{Hom}_A(M, E(X)) \otimes_B M \rightarrow E(X)$ with $\delta(\eta \otimes m) = \eta(m)$ for all $\eta \in \text{Hom}_A(M, E(X)), m \in M$; see [4] or [6] for details.

Given an arbitrary R -module $L = (C, D)_g$, according to [5], $\text{K.dim}(L_R) = \text{Max} \{ \text{K.dim}(C_A), \text{K.dim}(D_B) \}$. In particular, we have $\text{K.dim}(R_R) = \text{Max} \{ \text{K.dim}(A \oplus M)_A, \text{K.dim}(B_B) \}$, and for the simple R -modules V of the form $(0, Y)_0$ we have $\text{K.dim}(E(V))_R = \text{K.dim}(E(Y))_B$ while for the simple R -modules V of the form $(X, 0)_0$, $\text{K.dim}(E(V))_R = \text{Max} \{ \text{K.dim}(E(X))_A, \text{K.dim}[\text{Hom}_A(M, E(X))]_B \}$.

2. Co-Noetherian dimension

In this section we use the notation introduced in §1. Let $R \in \mathcal{W}$. If there exists an integer $n \geq 1$ such that $R \in \mathcal{W}_n \setminus \mathcal{W}_{n-1}$, we say that R has *co-Noetherian dimension* n , and write $\text{co-N.dim}(R) = n$. If on the other hand, $R \notin \mathcal{W}_m$ for all m , we say that R has infinite co-Noetherian dimension, and write $\text{co-N.dim}(R) = \infty$. Finally, we set $\text{co-N.dim}(R) = 0$

whenever R is co-Noetherian. We begin with the following result that characterizes when a ring R belongs to \mathcal{W}_n or to \mathcal{W} .

Theorem 2.1. *If $n \geq 0$ and R is a ring, then $R \in \mathcal{W}_n$ (respectively $R \in \mathcal{W}$) if and only if for all $M \in \text{Mod-}R$, $\tau_n(M) = 0$ (respectively $\tau(M) = 0$).*

Proof. First suppose that $\tau_n(M) = 0$ for all $M \in \text{Mod-}R$. Let M be an f.e. R -module. Because $\tau_n(M) = 0$ and M is f.e., there exist submodules $N_i (i = 1, \dots, t)$ such that $\bigcap_{i=1}^t N_i = 0$ and the Krull dimension of each M/N_i is at most n . It follows that M embedding in $\bigoplus_{i=1}^t (M/N_i)$ has Krull dimension at most n , as desired. For the converse, let $\{S_i \mid i \in I\}$ be a complete set of non-isomorphic simple R -modules. The module $C = \bigoplus_{i \in I} E(S_i)$ is a co-generator for $\text{Mod-}R$, hence if $0 \neq M_R$ is given, there are a set Λ and an embedding $f : M \rightarrow \prod_{\alpha \in \Lambda} C_\alpha$ where $C_\alpha = C$ for all $\alpha \in \Lambda$. For $i \in I$, let $\pi_{(\alpha, i)}$ be the projection of $\prod_{\alpha \in \Lambda} C_\alpha$ into the summand $E(S_i)$ of C_α . By assumption that the Krull dimension of each $E(S_i)$ is at most n , we see that $\text{K.dim}(M/\ker(\pi_{(\alpha, i)}f)) \leq n$. Clearly, $\tau_n(M) \subseteq \bigcap_{\alpha, i} \ker(\pi_{(\alpha, i)}f) = 0$. A slight modification of the above argument will prove the other statement. \square

Theorem 2.2. *All classes \mathcal{W}_n , as well as \mathcal{W} , are closed under Morita equivalence, homomorphic images, and finite normalizing extensions.*

Proof. That \mathcal{W} and each \mathcal{W}_n is closed under Morita equivalence follows from the standard techniques of Morita theory since “simple”, “essential monomorphism”, “injective” and “Krull dimension” are all Morita invariant properties. Next we prove that \mathcal{W}_n ($n \geq 1$) is closed under homomorphic images. Suppose that $R \in \mathcal{W}_n$, I is a proper ideal of R , S is a simple R/I -module and $E := E(S_R)$. It is well known that the set $E' := \{e \in E \mid eI = 0\}$ is the injective envelope of $S_{R/I}$. Hence $\text{K.dim}(E'_{R/I}) = \text{K.dim}(E'_R) \leq \text{K.dim}(E_R) \leq n$. This shows that $R/I \in \mathcal{W}_n$. We now prove that if $R \in \mathcal{W}_n$ and T is a finite normalizing extension of R , then $T \in \mathcal{W}_n$. In view of Theorem 2.1, we need to show that $\tau_n(M_T) = 0$ whenever M is a nonzero T -module. By assumption $T = \sum_{i=1}^t a_i R$ with a_i normalizing R . If N is an R -submodule of M , then according to [10]; p345 $Na_i^{-1} = \{m \in M \mid ma_i \in N\}$ is an R -submodule of M and the group monomorphism $M/Na_i^{-1} \rightarrow M/N$ given by $m + Na_i^{-1} \mapsto ma_i + N$ induces a lattice embedding of the R -submodules of M/Na_i^{-1} into the R -submodules of M/N . If

$b(N) = \cap_{i=1}^t Na_i^{-1}$, then $b(N)$ is the largest T -submodule of M contained in N and there is an R -monomorphism $M/b(N) \hookrightarrow \oplus_i (M/Na_i^{-1})$. Consequently, if $\text{K.dim}(M/N)_R \leq n$ we deduce that $\text{K.dim}[M/b(N)]_R \leq n$, hence $\text{K.dim}[M/b(N)]_T \leq n$. Now by hypothesis and Theorem 1, $\tau_n(M_R) = 0$, thus it follows that $\tau_n(M_T) = 0$. The proof of the implication “ $R \in \mathcal{W} \Rightarrow T \in \mathcal{W}$ ” is now evident. \square

Immediate consequences are given in the following.

Corollary 2.3. (i) *Let A and B be Morita equivalent rings under the category equivalence $\alpha : \text{Mod-}A \rightarrow \text{Mod-}B$. If $M \in \text{Mod-}A$ and $n \geq 1$, then $\alpha(\tau_n(M)) \simeq \tau_n(\alpha(M))$ and $\alpha(\tau(M)) \simeq \tau(\alpha(M))$. Furthermore $\text{co-N.dim}(A) = \text{co-N.dim}(B)$.*

(ii) *If $R \in \mathcal{W}$, I is a proper ideal of R and T is a finite normalizing extension of R , then $\text{co-N.dim}(R) \geq \text{Max} \{ \text{co-N.dim}(R/I), \text{co-N.dim}(T) \}$.*

Theorem 2.4. *For each integer $n \geq 1$, there exists a ring with co-Noetherian dimension n*

Proof. Given a positive integer n , let A be any commutative Noetherian local ring with $\text{K.dim}(A) = n$. If P is the unique maximal ideal of A , let $M = E(A/P)$. It is well-known that M_A is Artinian. Put $B = \text{End}(M_A)$, which by a classical result of Matlis [9] is isomorphic to \hat{A} , the P -adic completion of A , and consequently by [3, Proposition 10.16], B is also a local ring. Now consider the formal triangular matrix ring $R = \begin{bmatrix} A & 0 \\ M & B \end{bmatrix}$. Up to isomorphisms, the only simple modules over R are $V_1 = (A/P, 0)_0$ and $V_2 = (0, B/J(B))_0$. Thus we have the following descriptions: $E(V_1) = (E(A/P), \text{Hom}_A(M, E(A/P)))_\delta$, and $E(V_2) = (0, E(B/J(B)))_0$. It follows that $\text{K.dim}(R_R) = n = \text{Sup} \{ \text{K.dim}(E(V_R)) \mid V_R \text{ is simple} \}$. Thus we deduce that $\text{co-N.dim}(R) = n$. \square

Our next result determines the co-Noetherian dimension of commutative rings when they exist. For any ring R , let $\text{Ass}_E(R) = \{ P \mid P \in \text{Ass}(N) \text{ where } N \text{ is a sub-factor of } E(S) \text{ for some simple } S_R \}$.

Theorem 2.5. *$\text{Co-N.dim}(R) = \text{Sup} \{ \text{K.dim}(R/P) \text{ where } P \in \text{Ass}_E(R) \}$ for every commutative ring R in \mathcal{W} .*

Proof. Let $\text{co-N.dim}(R) = n$. Note that if $P \in \text{Ass}(N)$ for some N_R , then there exists a cyclic submodule C of N such that $P = \text{ann}_R(C)$. It follows that R/P embeds in N . Thus $\text{K.dim}(N) \geq \text{K.dim}(R/P)$ (if they exist). Now since $R \in \mathcal{W}$, $\text{Sup}\{\text{K.dim}(E(S)) \mid S_R \text{ is simple}\} = n$. Hence $n \geq \text{Sup}\{\text{K.dim}(R/P) \mid P \in \text{Ass}_E(R)\}$. On the other hand, if S is a simple R -module, then by a known result, $\text{K.dim}(E(S)) = \text{Sup}\{\text{K.dim}(C) \mid C \text{ is a critical } R\text{-module contained in a sub-factor of } E(S)\}$; see [11, 4.19] or Corollary 3.2.12. Since for every critical R -module C , $\text{K.dim}(xR) = \text{K.dim}(C)$ for all $0 \neq x \in C$, we can conclude that $\text{K.dim}(E(S)) = \text{Sup}\{\text{K.dim}(C) \mid C \text{ is a critical cyclic } R\text{-module contained in a sub-factor of } E(S)\}$. The result is now clear by the well known fact that critical cyclic R -modules have the form R/P for some $P \in \text{Spec}(R)$; see for example [1, Lemma 3.5]. \square

We already know that any commutative Noetherian ring is necessarily co-Noetherian. Thus there are plenty of commutative rings R with $\text{co-N.dim}(R) < \text{K.dim}(R)$. Our next result provides a class of rings that are not necessarily commutative, yet for them the strict inequality holds.

Proposition 2.6. (i) *Let R be a semiprime right Goldie ring. If R has non-zero Krull dimension, then either $R \notin \mathcal{W}$ or $\text{co-N.dim}(R) < \text{K.dim}(R)$.*

(ii) *Let R be a right semi-Artinian ring. Then either $R \notin \mathcal{W}$ or $\text{co-N.dim}(R) = 0$.*

Proof. Let $R \in \mathcal{W}$ and S be a simple R -module.

(i) Let $\text{K.dim}(R) = n \geq 1$. By hypothesis, $\text{K.dim}(E(S))$ and, hence $\text{K.dim}(E(S)/S)$ exist. Let $x \in E(S)/S$. Thus $xR \simeq R/A$ for some essential right ideal A of R . Because R is semiprime right Goldie, A contains a regular element c of R . Thus we have $\text{K.dim}(R/A) \leq \text{K.dim}(R/cR) < n$ by [10, Lemma 6.3.9]. It follows that $\text{K.dim}(E(S)/S) \leq n - 1$ by [10, Lemma 6.2.17]. Consequently, one obtains

$$\text{co-N.dim}(R) = \text{Sup}\{\text{K.dim}(E(S_R)) \mid S_R \text{ is simple}\} \leq n - 1 < n.$$

(ii) Let $B =$ the sum of all Artinian submodules of $E(S)$. Then since $E(S)$ has Krull dimension so does B . It follows that B is an Artinian R -module. Now if $E(S)/B$ is non-zero, it contains a simple submodule L/B by our assumption on R . But then L is an Artinian submodule of $E(S)$ and so $L \subseteq B$, a contradiction. Therefore, $E(S) = B$ is an Artinian R -module, proving that $\text{co-N.dim}(R) = 0$. \square

Corollary 2.7. *Let R be a ring with Krull-dimension and P be a prime ideal of R . If $\text{co-N.dim}(R/P) = \text{K.dim}(R/P)$, then P is a maximal ideal of R .*

Proof. By [10, Proposition 6.3.5], R/P is a prime right Goldie ring. Hence, $\text{K.dim}(R/P)$ must be zero by Proposition 2.6(i). It follows that R/P is a right Artinian ring and hence P is a maximal ideal of R . \square

We note that the converse to the above Corollary is not true in general. For instance, if $R = K[x, D]$ is the Cozzens's example where K is a universal field with derivation D , then it is well known that R is a simple principal right (and left) ideal domain with a unique (up to isomorphism) simple injective R -module and $\text{K.dim}(R) = 1 \neq \text{co-N.dim}(R)$.

3. Examples and concluding remarks

This section is devoted to explore possible relations between Krull dimension and co-Noetherian dimension of rings. If R is a ring, provided that for R both Krull dimension and co-Noetherian dimension exist, we have $\text{co-N.dim}(R) \leq \text{K.dim}(R)$. The proof of Theorem 2.4 shows that equality can occur.

Example 3.1. If p is a fixed prime number, the ring $\begin{bmatrix} \mathbb{Z} & 0 \\ \mathbb{Z}_{p^\infty} & \mathbb{Z} \end{bmatrix}$ is not of the type used in the proof of Theorem 2.4, yet for it the Krull dimension coincides with the co-Noetherian dimension and this common value is 1. This example also shows that the semiprime condition in Proposition 2.6(i) cannot be deleted.

The following examples show that in general co-Noetherian dimension and Krull dimension of a ring are independent of each other.

Example 3.2. There is an Artinian ring with a simple module whose injective envelope has no Krull dimension: Let $B \subseteq A$ denote Cohn's division rings such that A_B has finite dimension while the left B -module A has infinite dimension. Then the ring $R = \begin{bmatrix} A & 0 \\ A & B \end{bmatrix}$ is right Artinian

and the injective envelope of the simple R -module $(A, 0)_0$ has no Krull dimension. Thus $R \notin W$, so $\text{co-N.dim}(R)$ is not defined.

Example 3.3. It is well-known that the first Weyl algebra $A = A_1(k)$ over a field of characteristic zero is a simple Noetherian domain of Krull dimension 1, and that it is not co-Noetherian, as shown in [7, Example 2.1]. Thus by Proposition 2.6(i), for A even the co-Noetherian dimension cannot exist. This fact in particular shows that in general, co-Noetherian dimension is not well behaved under Ore ring extension. Now set $R = \begin{bmatrix} A & 0 \\ M & k \end{bmatrix}$ where M is an infinite direct sum of any non-zero A -module. It is clear that R does not have Krull-dimension. Moreover, as A is a homomorphic image of R , by Corollary 2.3, we deduce that R does not have co-Noetherian dimension.

Example 3.4. We construct a co-Noetherian ring that has no Krull dimension. Let $A = \mathbb{C}[x]$, and consider an infinite sequence of pairwise non-isomorphic simple A -modules $\{S_i\}_{i \geq 1}$. Put $M = \bigoplus_i S_i$ and consider the ring $R = \begin{bmatrix} A & 0 \\ M & \mathbb{C} \end{bmatrix}$. Suppose $V = (S', 0)_0$ is a simple R -module. Then $E(V) = (E(S'), \text{Hom}_A(M, E(S')))_{\delta}$. We have $T := \text{Hom}_A(M, E(S')) \simeq \prod_{i \geq 1} \text{Hom}_A(S_i, E(S'))$. If $S_i \not\simeq S'$ for all i , then $T = 0$. Suppose $S_j \simeq S'$ for some j . In this case, we have:

$$T \simeq \text{Hom}_A(S_j, E(S_j)) \hookrightarrow \text{Hom}_A(S_j, S_j) = \mathbb{C}$$

Thus in any case T is a finite dimensional \mathbb{C} -space. Since $E(S')$ is an Artinian A -module, we deduce that $E(V)$ is an Artinian R -module. All other simple R -modules are of the form $(0, \mathbb{C})_0$ which is Artinian and injective. Therefore R is co-Noetherian.

Remark 3.5. (i) It is possible to modify the definition of co-Noetherian dimension by restricting to those simple modules whose injective envelopes do have Krull-dimensions (not necessarily finite). Then for any ring this new dimension exists, it is less than or equal to the global Krull-dimension of the ring in the sense of Albu and Smith [11], and results similar to Theorems 2.1 and 2.2 hold. But when this new dimension is zero (as it is the case for the first Weyl algebra over a field with zero characteristic) the ring is not necessarily co-Noetherian. Hence it cannot measure distance from being co-Noetherian. On the other hand, our

construction of rings with co-Noetherian dimension n cannot be carried over to the case of infinite ordinal numbers. These considerations indicate the reason for our Standing Hypothesis. But allowing modules with arbitrary Krull dimensions the following problem is naturally imposed:

Open Problem. Given an arbitrary ordinal number γ , does there exist a ring R with a simple module S such that $E(S)$ has Krull-dimension γ ?

(ii) By using “ Noetherian dimension ”, it is possible to define “ co-Artinian dimension ” of a ring and prove results analogous to Theorems 2.1 and 2.2 However, one must first prove the existence of rings with arbitrary co-Artinian dimensions.

Acknowledgments

The authors thank the referee for suggestions towards improvement of the presentation. The first named author also wishes to express his gratitude to Professors K. R. Goodearl, J. T. Stafford and O. A. S. Karamzadeh for email communications regarding the subject matter of this paper.

REFERENCES

- [1] T. Albu and P. F. Smith, Global Krull Dimension, *Interactions Between Ring Theory and Representations of Algebras (Murcia)*, Edited by F. van Oystaeyen and M. Saorin, Dekker, New York, *Lecture Notes in Pure and Appl. Math.* **210** (2000) 1-21.
- [2] F.W. Anderson and K.R. Fuller, Rings and Categories of Modules, Springer-Verlag, New York-Heidelberg, 1974.
- [3] M. F. Atiyah and I. G. MacDonald, Introduction to Commutative Algebra, Addison-Wesley Publication Co., London, 1969.
- [4] A. Haghany, Injectivity conditions over a formal triangular matrix ring, *Arch. Math. (Basel)* **78** (2002), no. 4, 268–274.
- [5] A. Haghany, Krull and Noetherian dimensions for modules over formal triangular matrix rings, *Iran. J. Sci. Technol. Trans. A Sci.* **27** (2003), no. 2, 403–406.
- [6] A. Haghany and K. Varadarajan, Study of formal triangular matrix rings, *Comm. Algebra* **27** (1999), no. 11, 5507–5525.
- [7] Y. Hirano, On injective hulls of simple modules, *J. Algebra* **225** (2000), no. 1, 299–308.
- [8] J. P. Jans, On co-Noetherian rings, *J. London Math. Soc.* **1** (1969) 588–590.

- [9] E. Matlis, Injective Modules over Noetherian rings, *Pacific J. Math* **8** (1958) 511–528.
- [10] J. C. McConnell and J. C. Robson, Noncommutative Noetherian rings, Wiley Interscience Publication, New York, 1987.
- [11] C. Năstăsescu and F. van Oystaeyen, Dimensions of Ring Theory, D. Reidel Publishing Co., Dordrecht, 1987.
- [12] P. Vámos, The dual of the notion of “finitely generated”, *J. London Math. Soc.* **43** (1968) 643–646.
- [13] R. Wisbauer, Foundations of Module and Ring Theory, Gordon and Breach Science Publishers, Philadelphia, 1991.

Ahmad Haghany

Department of Mathematical Sciences, Isfahan University of Technology, 84156-83111
Isfahan, Iran

Email: aghagh@cc.iut.ac.ir

Mohammad Reza Vedadi

Department of Mathematical Sciences, Isfahan University of Technology, 84156-83111
Isfahan, Iran

Email: mrvedadi@cc.iut.ac.ir