

ISSN: 1017-060X (Print)



ISSN: 1735-8515 (Online)

**Bulletin of the**  
**Iranian Mathematical Society**

Vol. 40 (2014), No. 1, pp. 1–7

**Title:**

**A lower estimate of harmonic functions**

**Author(s):**

**G. Pan, L. Qiao and G. Deng**

Published by Iranian Mathematical Society  
<http://bims.ims.ir>

## A LOWER ESTIMATE OF HARMONIC FUNCTIONS

G. PAN\*, L. QIAO AND G. DENG

(Communicated by Javad Mashreghi)

**ABSTRACT.** We shall give a lower estimate of harmonic functions of order greater than one in a half space, which generalize the result obtained by B. Ya. Levin in a half plane.

**Keywords:** Lower estimate, Harmonic function, Half space.

**MSC(2010):** Primary: 31B05; Secondary: 31B10.

### 1. Introduction

Let  $\mathbf{R}$  and  $\mathbf{R}_+$  be the sets of all real numbers and of all positive real numbers, respectively. Let  $\mathbf{R}^n$  ( $n \geq 2$ ) denote the  $n$ -dimensional Euclidean space with points  $x = (x', x_n)$ , where  $x' = (x_1, x_2, \dots, x_{n-1}) \in \mathbf{R}^{n-1}$  and  $x_n \in \mathbf{R}$ . The boundary and closure of an open set  $D$  of  $\mathbf{R}^n$  are denoted by  $\partial D$  and  $\bar{D}$ , respectively. The upper half space is the set  $H = \{(x', x_n) \in \mathbf{R}^n : x_n > 0\}$ , whose boundary is  $\partial H$ .

For a set  $E$ ,  $E \subset \mathbf{R}_+ \cup \{0\}$ , we denote  $\{x \in H : |x| \in E\}$  and  $\{x \in \partial H : |x| \in E\}$  by  $HE$  and  $\partial HE$ , respectively. We identify  $\mathbf{R}^n$  with  $\mathbf{R}^{n-1} \times \mathbf{R}$  and  $\mathbf{R}^{n-1}$  with  $\mathbf{R}^{n-1} \times \{0\}$ , writing typical points  $x, y \in \mathbf{R}^n$  as  $x = (x', x_n)$ ,  $y = (y', y_n)$ , where  $y' = (y_1, y_2, \dots, y_{n-1}) \in \mathbf{R}^{n-1}$  and putting

$$x \cdot y = \sum_{j=1}^n x_j y_j = x' \cdot y' + x_n y_n, \quad |x| = \sqrt{x \cdot x}, \quad |x'| = \sqrt{x' \cdot x'},$$

$$|x'| = |x| \cos \theta \text{ and } x_n = |x| \sin \theta \quad (0 < \theta \leq \pi/2).$$

---

Article electronically published on February 25, 2014.

Received: 10 September 2012, Accepted: 2 November 2012.

\*Corresponding author.

Let  $B_r$  denote the open ball with center at the origin and radius  $r$  ( $> 0$ ) in  $\mathbf{R}^n$ . We use the standard notations  $u^+ = \max\{u, 0\}$  and  $u^- = -\min\{u, 0\}$ . In the sense of Lebesgue measure  $dy' = dy_1 \cdots dy_{n-1}$  and  $dy = dy' dy_n$ . Let  $\sigma$  denote  $(n-1)$ -dimensional surface area measure and  $\partial/\partial n$  denote differentiation along the inward normal into  $H$ .

The estimate we deal with has a long history which can be traced back to Levin's estimate of harmonic functions from below (see, for example, Levin [6, p. 209]).

**Theorem 1.1.** *Let  $A_1$  be a constant,  $u(z)$  harmonic in the upper half space  $\mathbf{C}_+$  and continuous on  $\partial\mathbf{C}_+$ . Suppose that*

$$u(z) \leq A_1 R^\rho, \quad z \in \mathbf{C}_+, \quad R = |z| > 1, \quad \rho > 1$$

and

$$|u(z)| \leq A_1, \quad |z| \leq 1, \quad \text{Im}z \geq 0.$$

Then

$$u(Re^{i\varphi}) \geq -A_2 A_1 (1 + R^\rho) \sin^{-1} \varphi, \quad Re^{i\varphi} \in \mathbf{C}_+,$$

where  $A_2$  is a constant independent of  $A_1$ ,  $R$ ,  $\varphi$  and the function  $u(z)$ .

Further versions and refinements of Theorem 1.1 may be found in the monograph Nikol'skiĭ [7, Ch. 1] and in the paper Krasichkov-Ternovskii [3].

In this article, we will consider functions  $u(x)$  harmonic in  $H$  and continuous on  $\overline{H}$ . In what follows we shall denote by  $M$  various values which does not depend on  $K$ ,  $R$  ( $= |x|$ ),  $\theta$  and the function  $u(x)$ .

In this note we prove analogous estimates for  $u(x)$  in  $H$ .

**Theorem 1.2.** *Suppose that*

$$(1.1) \quad u(x) \leq KR^{\rho(R)}, \quad x \in H, \quad R = |x| > 1, \quad \rho(R) > 1$$

and

$$(1.2) \quad u(x) \geq -K, \quad |x| \leq 1, \quad x_n \geq 0.$$

Then

$$u(x) \geq -MK \left(1 + (2R)^{\rho(2R)}\right) \sin^{1-n} \theta,$$

where  $x \in H$  and  $\rho(R)$  is nondecreasing on  $[1, +\infty)$ .

**Remark 1.3.** *If  $n = 2$  and  $\rho(R) \equiv \rho$ , Theorem 1.2 is just a consequence of Theorem 1.1.*

**Theorem 1.4.** *If (1.1) and (1.2) hold, then*

$$u(x) \geq -MK \left( 1 + \left( \frac{N+1}{N} R \right)^{\rho \left( \frac{N+1}{N} R \right)} \right) \sin^{1-n} \theta,$$

where  $x \in H$ ,  $N(\geq 1)$  is a sufficiently large number and  $\rho(R)$  is as defined in Theorem 1.2.

## 2. Lemmas

Carleman's formula [2] connects the modulus and the zeros of a function analytic in  $\mathbf{C}_+$  (see, for example, [5, p. 224]). Nevanlinna's formula (see [6, p. 193]) refers to a harmonic function in a half disk. Armitage and Kuran obtained a generalized Nevanlinna-type formula in a half space and Poisson integral formula for half balls respectively, which play important roles in our discussions.

**Lemma 2.1.** ([1]). *If  $R > 1$ , then we have*

$$\int_{\{x \in H: |x|=R\}} u(x) \frac{nx_n}{R^{n+1}} d\sigma(x) + \int_{\partial H(1,R)} u(x') \left( \frac{1}{|x'|^n} - \frac{1}{R^n} \right) dx' = c_1 + \frac{c_2}{R^n},$$

where

$$c_1 = \int_{\{x \in H: |x|=1\}} \left( (n-1)x_n u(x) + x_n \frac{\partial u(x)}{\partial n} \right) d\sigma(x)$$

and

$$c_2 = \int_{\{x \in H: |x|=1\}} \left( x_n u(x) - x_n \frac{\partial u(x)}{\partial n} \right) d\sigma(x).$$

**Lemma 2.2.** ([4]). *Let  $R > 1$ ,  $u(x)$  be a function in  $B_R^+ = B_R \cap H$  and continuous in  $\overline{B}_R^+$ . Then*

$$\begin{aligned} u(x) &= \int_{\{y \in H: |y|=R\}} \frac{R^2 - |x|^2}{\omega_n R} \left( \frac{1}{|y-x|^n} - \frac{1}{|y-x^*|^n} \right) u(y) d\sigma(y) \\ &+ \frac{2x_n}{\omega_n} \int_{\partial H[0,R)} \left( \frac{1}{|y'-x|^n} - \frac{R^n}{|x|^n} \frac{1}{|y'-\tilde{x}|^n} \right) u(y') dy', \end{aligned}$$

where  $x \in B_R^+$ ,  $\tilde{x} = R^2 x / |x|^2$ ,  $x^* = (x', -x_n)$  and  $\omega_n = \pi^{\frac{n}{2}} / \Gamma(1 + \frac{n}{2})$  is the volume of the unit  $n$ -ball in  $\mathbf{R}^n$ .

### 3. Proof of Theorem 1

By applying Lemma 2.1 to  $u(x)$ , we have

$$\begin{aligned}
 (3.1) \quad & \int_{\{x \in H: |x|=R\}} u^+(x) \frac{nx_n}{R^{n+1}} d\sigma(x) + \int_{\partial H(1,R)} u^+(x') \left( \frac{1}{|x'|^n} - \frac{1}{R^n} \right) dx' \\
 &= \int_{\{x \in H: |x|=R\}} u^-(x) \frac{nx_n}{R^{n+1}} d\sigma(x) \\
 &+ \int_{\partial H(1,R)} u^-(x') \left( \frac{1}{|x'|^n} - \frac{1}{R^n} \right) dx' + c_1 + \frac{c_2}{R^n}.
 \end{aligned}$$

It immediately follows from (1.1) that

$$(3.2) \quad \int_{\{x \in H: |x|=R\}} u^+(x) \frac{nx_n}{R^{n+1}} d\sigma(x) \leq MKR^{\rho(R)-1}$$

and

$$(3.3) \quad \int_{\partial H(1,R)} u^+(x') \left( \frac{1}{|x'|^n} - \frac{1}{R^n} \right) dx' \leq MKR^{\rho(R)-1}.$$

Hence from (3.1), (3.2) and (3.3) we have

$$(3.4) \quad \int_{\{x \in H: |x|=R\}} u^-(x) \frac{nx_n}{R^{n+1}} d\sigma(x) \leq MKR^{\rho(R)-1}$$

and

$$(3.5) \quad \int_{\partial H(1,R)} u^-(x') \left( \frac{1}{|x'|^n} - \frac{1}{R^n} \right) dx' \leq MKR^{\rho(R)-1}.$$

And (3.5) gives

$$\begin{aligned}
 & \int_{\partial H(1,R)} \frac{u^-(x')}{|x'|^n} dx' \\
 & \leq \frac{2^n}{2^n - 1} \int_{\partial H(1,R)} u^-(x') \left( \frac{1}{|x'|^n} - \frac{1}{(2R)^n} \right) dx' \\
 (3.6) \quad & \leq MK(2R)^{\rho(2R)-1}.
 \end{aligned}$$

Since  $-u(x) \leq u^-(x)$ , by applying Lemma 2.2 to  $-u(x)$  we have

$$(3.7) \quad -u(x) \leq I_1(x) + I_2(x),$$

where

$$I_1(x) = \int_{\{y \in H: |y|=R\}} \frac{R^2 - |x|^2}{\omega_n R} \left( \frac{1}{|y-x|^n} - \frac{1}{|y-x^*|^n} \right) u^-(y) d\sigma(y)$$

and

$$I_2(x) = \frac{2x_n}{\omega_n} \int_{\partial H[0,R]} \left( \frac{1}{|y' - x|^n} - \frac{R^n}{|x|^n} \frac{1}{|y' - \tilde{x}|^n} \right) u^-(y') dy'.$$

We remark that

$$(3.8) \quad \frac{1}{|y - x|^n} - \frac{1}{|y - x^*|^n} \leq \frac{2nx_n y_n}{|y - x|^{n+2}}$$

and

$$(3.9) \quad |y - x|^n \geq x_n^n = |x|^n \sin^n \theta, \quad x \in H, \quad y_n = 0.$$

If we put  $|x| = r > 1/2$  and  $R = 2r$  in (3.7), then we finally have from (3.4), (3.8) and (3.9)

$$(3.10) \quad \begin{aligned} I_1(x) &\leq \int_{\{y \in H: |y|=R\}} \frac{R^2 - r^2}{\omega_n R} \frac{2nx_n y_n}{\omega_n |y - x|^{n+2}} u^-(y) d\sigma(y) \\ &\leq MKR^{\rho(R)} \end{aligned}$$

and

$$(3.11) \quad I_2(x) \leq I_{21}(x) + I_{22}(x),$$

where

$$I_{21}(x) = \frac{2}{\omega_n x_n^{n-1}} \int_{\partial H(1,R)} u^-(y') dy'$$

and

$$I_{22}(x) = \frac{2}{\omega_n x_n^{n-1}} \int_{\partial H[0,1]} u^-(y') dy'.$$

We obtain that

$$(3.12) \quad \begin{aligned} I_{21}(x) &\leq \frac{2R^n}{\omega_n x_n^{n-1}} \int_{\partial H(1,R)} \frac{u^-(y')}{|y'|^n} dy' \\ &\leq MK(2R)^{\rho(2R)} \sin^{1-n} \theta \end{aligned}$$

and

$$(3.13) \quad \begin{aligned} I_{22}(x) &\leq \frac{2K}{\omega_n x_n^{n-1}} \int_{\partial H[0,1]} dy' \\ &\leq MK \sin^{1-n} \theta \end{aligned}$$

from (3.6) and (1.2), respectively.

From (3.7), (3.10), (3.11), (3.12) and (3.13), we have for  $|x| > 1/2$

$$(3.14) \quad -u(x) \leq MK \left( 1 + (2R)^{\rho(2R)} \right) \sin^{1-n} \theta.$$

For  $|x| \leq 1/2$ , we have from (1.2)

$$(3.15) \quad -u(x) \leq K \leq K \left(1 + (2R)^{\rho(2R)}\right) \sin^{1-n} \theta.$$

Thus the conclusion immediately follows from (3.14) and (3.15).

#### 4. Proof of Theorem 2

By modifying (3.6), we have

$$\begin{aligned} & \int_{\partial H(1,R)} \frac{u^-(x')}{|x'|^n} dx' \\ & \leq \frac{(N+1)^n}{(N+1)^n - N^n} \int_{\partial H(1,R)} u^-(x') \left( \frac{1}{|x'|^n} - \frac{1}{\left(\frac{N+1}{N}R\right)^n} \right) dx' \\ & \leq MK \left(\frac{N+1}{N}R\right)^{\rho\left(\frac{N+1}{N}R\right)-1}. \end{aligned}$$

Then (3.12), (3.14) and (3.15) are replaced by the following estimates

$$(4.1) \quad I_{21}(x) \leq MK \left(\frac{N+1}{N}R\right)^{\rho\left(\frac{N+1}{N}R\right)-1} \sin^{1-n} \theta.$$

$$(4.2) \quad -u(x) \leq MK \left(1 + \left(\frac{N+1}{N}R\right)^{\rho\left(\frac{N+1}{N}R\right)}\right) \sin^{1-n} \theta.$$

$$(4.3) \quad -u(x) \leq K \leq MK \left(1 + \left(\frac{N+1}{N}R\right)^{\rho\left(\frac{N+1}{N}R\right)}\right) \sin^{1-n} \theta.$$

All (3.7), (3.10), (3.11), (4.1), (3.12), (4.2) and (4.3) give

$$u(x) \geq -MK \left(1 + \left(\frac{N+1}{N}R\right)^{\rho\left(\frac{N+1}{N}R\right)}\right) \sin^{1-n} \theta$$

from which the conclusion immediately follows.

#### Acknowledgments

The authors are grateful to the referee for her or his careful reading and helpful suggestions which led to an improvement of their original manuscript. This work is supported by the National Natural Science Foundation of China under Grant Nos. 11271045, 11301140 and U1304102 and Specialized Research Fund for the Doctoral Program of Higher Education under Grant No. 20100003110004.

## REFERENCES

- [1] D. H. Armitage, A Nevanlinna theorem for superharmonic functions in half-spaces, with applications, *J. London Math. Soc.(2)* **23** (1981), no. 1, 137–157.
- [2] T. Carleman, Über die approximation analytischer Funktionen durch lineare aggregate von vorgegebenen potenzen, *Ark. Mat., Astr. Och Fysik.* **17** (1923) 1–30.
- [3] I. F. Krasichkov-Ternovskii, Estimates for the subharmonic difference of subharmonic functions. II, *Mat. Sb.* **32** (1977) 32–59.
- [4] Ü. Kuran, Harmonic majorizations in half-balls and half-spaces, *Proc. London Math. Soc. (3)* **21** (1970) 614–636.
- [5] B. Ya. Levin, Distribution of Zeros of Entire Functions, Revised ed., Translations of Mathematical Monographs, 5, Amer. Math. Soc., Providence, 1980.
- [6] B. Ya. Levin, Lectures on Entire Functions, Translations of Mathematical Monographs, 150, Amer. Math. Soc., Providence, 1980.
- [7] N. K. Nikol'skii, Selected problems of the weighted approximation and of spectral analysis, *Trudy Mat. Inst. Steklov. Inst. Steklov.* **21** (1974); *English transl. in Proc. Steklov Inst. Math.* **120** (1976).

(Guoshuang Pan) BEIJING NATIONAL DAY SCHOOL, BEIJING, PEOPLE'S REPUBLIC OF CHINA

*E-mail address:* `gsp1979@163.com`

(Lei Qiao) COLLEGE OF MATHEMATICS AND INFORMATION SCIENCE, HENAN UNIVERSITY OF ECONOMICS AND LAW, ZHENGZHOU, PEOPLE'S REPUBLIC OF CHINA

*E-mail address:* `qiaocqu@163.com`

(Guantie Deng) SCHOOL OF MATHEMATICAL SCIENCE, BEIJING NORMAL UNIVERSITY, LABORATORY OF MATHEMATICS AND COMPLEX SYSTEMS, MOE, BEIJING, PEOPLE'S REPUBLIC OF CHINA

*E-mail address:* `denggt@bnu.edu.cn`