

ISSN: 1017-060X (Print)



ISSN: 1735-8515 (Online)

Bulletin of the
Iranian Mathematical Society

Vol. 41 (2015), No. 5, pp. 1183–1193

Title:

On Sandwich theorems for certain classes of analytic functions

Author(s):

R. Aghalary, A. Ebadian and M. Mafakheri

Published by Iranian Mathematical Society
<http://bims.ims.ir>

ON SANDWICH THEOREMS FOR CERTAIN CLASSES OF ANALYTIC FUNCTIONS

R. AGHALARY*, A. EBADIAN AND M. MAFAKHERI

(Communicated by Ali Abkar)

ABSTRACT. The purpose of this paper is to derive some subordination and superordination results for certain analytic functions in the open unit disk.

Keywords: Subordination, superordination, integral operators, Hadamard product.

MSC(2010): Primary 30C45; Secondary 30C80.

1. Introduction and preliminaries

Let H be the class of all functions analytic in the unit disc $U = \{z \in \mathbb{C} : |z| < 1\}$ and for $a \in \mathbb{C}$ and $n \in \mathbb{N}$, let $H[a, n] = \{f \in H : f(z) = a + a_n z^n + \dots\}$. Let

$$\mathcal{A}_n = \{f \in H : f(z) = z + a_{n+1} z^{n+1} + \dots\},$$

and set $\mathcal{A} := \mathcal{A}_1$. We say that the function $f \in H$ is subordinate to the function $g \in H$ in the unit disc U (written $f \prec g$) if there exists an analytic function w in U with $w(0) = 0$ and $|w(z)| < 1$ for all $z \in U$ such that $f(z) = g(w(z))$ in U . If g is univalent in U , then the following equivalence relationship holds true:

$$f(z) \prec g(z) \Leftrightarrow f(0) = g(0) \text{ and } f(U) \subset g(U).$$

For $f(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=0}^{\infty} b_n z^n$, the Hadamard product (or convolution) of f and g is defined by

$$(f * g)(z) = \sum_{n=0}^{\infty} a_n b_n z^n.$$

Recently, Komatu [5] introduced a certain integral operator I_a^λ defined by

Article electronically published on October 15, 2015.

Received: 1 May 2014, Accepted: 18 July 2014.

*Corresponding author.

$$(1.1) \quad I_a^\lambda f(z) = \frac{a^\lambda}{\Gamma(\lambda)} \int_0^1 t^{a-2} (\log \frac{1}{t})^{\lambda-1} f(tz) dt \quad (z \in U, a > 0, \lambda \geq 0, f \in \mathcal{A}).$$

If $f \in \mathcal{A}_n$ is of the form $f(z) = z + \sum_{k=n+1}^\infty a_k z^k$, then it is easy to see from (1.1) that

$$I_a^\lambda f(z) = z + \sum_{k=n+1}^\infty \left(\frac{a}{a+k-1}\right)^\lambda a_k z^k.$$

We now introduce the operator $L_a^\lambda(b, c, z) : \mathcal{A}_n \rightarrow \mathcal{A}_n$ by

$$(1.2) \quad L_a^\lambda(b, c, z)f(z) = \left[z + \sum_{k=n+1}^\infty \frac{(b)_{k-1}}{(c)_{k-1}} \left(\frac{a+1}{a+k}\right)^\lambda z^k \right] * f(z)$$

where $\lambda \geq 0, b, a \in \mathbb{C}, \operatorname{Re}(a) > -1, c \in \mathbb{C} - \{0, -1, -2, \dots\}$. In view of (1.2), for $f \in \mathcal{A}_n$ and $z \in U$ it follows that

$$(1.3) \quad z(L_a^{\lambda+1}(b, c, z)f)'(z) = (1+a)L_a^\lambda(b, c, z)f(z) - aL_a^{\lambda+1}(b, c, z)f(z),$$

and

$$(1.4) \quad z(L_a^\lambda(b, c, z)f)'(z) = bL_a^\lambda(b+1, c, z)f(z) - (b-1)L_a^\lambda(b, c, z)f(z).$$

Furthermore let Q be the set of all analytic and univalent functions on the set $\bar{U}/E(f)$, such that $f'(\zeta) \neq 0$ for all $\zeta \in \partial U/E(f)$. where

$$E(f) := \{\zeta \in \partial U : \lim_{z \rightarrow \zeta} f(z) = \infty\},$$

The subclass of Q for which $f(0) = a$ is denoted by $Q(a)$. We shall need the following definition and lemmas to prove our results.

Definition 1.1. A function $p(z, t) : U \times [0, \infty) \rightarrow \mathbb{C}$ is called a subordination (or Loewner) chain if $p(\cdot, t)$ is analytic and univalent in U , for all $t \geq 0$ and $p(\cdot, t_1) \prec p(\cdot, t_2)$, for all $0 \leq t_1 \leq t_2$.

The next lemma gives a sufficient condition so that the function $p(\cdot, t)$ to be a subordination chain.

Lemma 1.2. (see [9, p.159]) Let

$$p(z, t) = a_1(t)z + a_2(t)z^2 + \dots \quad (a_1(t) \neq 0, t \geq 0),$$

with $a_1(t) \neq 0$ for all $t \geq 0$ and $\lim_{t \rightarrow \infty} |a_1(t)| = \infty$.

Suppose that $p(\cdot, t)$ is analytic in U for all $t \geq 0$, and $p(z, \cdot)$ is continuously differentiable on $[0, \infty)$ for any $z \in U$. If $p(z, t)$ satisfies

$$\operatorname{Re} \left(z \frac{\partial p(z, t)/\partial z}{\partial p(z, t)/\partial t} \right) > 0 \quad (z \in U, t \geq 0),$$

and

$$|p(z, t)| \leq k_0 |a_1(t)|, \quad |z| < r_0 < 1, t \geq 0$$

for some positive constants k_0 and r_0 , then, $p(z, t)$ is a subordination chain.

Lemma 1.3. (see [7]) Suppose that the function $\psi : \mathbb{C}^2 \times U \rightarrow \mathbb{C}$ satisfies the condition

$$\psi(ix, y; z) \leq \delta,$$

for real x , $y \leq -n(\frac{1+x^2}{2})$ and for all $z \in U$. If the function $\varphi(z) = 1 + a_n z^n + \dots$ is analytic in U and

$$\operatorname{Re}\{\psi(\varphi(z), z\varphi'(z); z)\} > \delta,$$

then $\operatorname{Re}\{\varphi(z)\} > 0$ in U .

Lemma 1.4. (see [8]) Let $q \in H[a, 1]$ and $\phi : \mathbb{C}^2 \rightarrow \mathbb{C}$. Also set

$$\phi(q(z), zq'(z)) \equiv h(z) \quad (z \in U).$$

If $P(z, t) := \phi(q(z), tzq'(z))$ is subordination chain and $p \in H[a, 1] \cap Q(a)$, then

$$h(z) \prec \phi(p(z), zp'(z))$$

implies that

$$q(z) \prec p(z).$$

Furthermore, if $\phi(q(z), zq'(z)) = h(z)$ has a univalent solution $q \in Q(a)$, then q is the best subordinate.

Making use of the principle of subordination between analytic functions, many authors (see for example [4, 6–8]) obtained some interesting subordination-preserving properties for certain subclasses of the class \mathcal{A}_1 . In the present paper we aim to investigate the subordination and superordination-preserving properties of the integral operator $L_a^\lambda(b, c, z)f$ in the class \mathcal{A}_n . Our results extends many results obtained by other authors (see for example [1–3]). Throughout the following section we assume that $b \in \mathbb{C}$, $\lambda \geq 0$ and $c \in \mathbb{C} - \{0, -1, -2, -3, \dots\}$.

2. Main results

Our first result involve the integral operator $L_a^\lambda(b, c, z)$.

Lemma 2.1. Let f be in the class \mathcal{A}_n , $(L_a^{\lambda+1}(b, c, z)f)'(z) \neq 0$ for all $z \in U$ and $a \in \mathbb{C}$ with $\operatorname{Re}(a) > 0$. If

$$(2.1) \quad \operatorname{Re} \left\{ 1 + \frac{z(L_a^\lambda(b, c, z)f)''(z)}{(L_a^\lambda(b, c, z)f)'(z)} \right\} > -\delta$$

then

$$(2.2) \quad \operatorname{Re} \left\{ 1 + \frac{z(L_a^{\lambda+1}(b, c, z)f)''(z)}{(L_a^{\lambda+1}(b, c, z)f)'(z)} \right\} > 0$$

where

$$(2.3) \quad \delta = n \frac{1 + |a|^2 - |1 - a^2|}{4\operatorname{Re}(a)}.$$

Proof. Let $f \in \mathcal{A}_n$ and

$$(2.4) \quad p(z) = 1 + \frac{z(L_a^{\lambda+1}(b, c, z)f)''(z)}{(L_a^{\lambda+1}(b, c, z)f)'(z)}.$$

It is easy to see that p is analytic in U and $p(z) = 1 + b_n z^n + b_{n+1} z^{n+1} + \dots$. Taking the logarithmic differentiation in (2.4) and using identity (1.3) in the resulting equation, we get that

$$(2.5) \quad p(z) + \frac{zp'(z)}{a+p(z)} = 1 + \frac{z(L_a^\lambda(b, c, z)f)''(z)}{(L_a^\lambda(b, c, z)f)'(z)}.$$

To make use of Lemma 1.3 we consider the function

$$\psi(r, s) = r + \frac{s}{r+a} + \delta.$$

In view of (2.1) and (2.5) we have $Re\{\psi(p(z), zp'(z))\} > 0$. Furthermore for $x \in \mathbb{R}$ and $y < \frac{-n}{2}(1+x^2)$ we have

$$Re\{\psi(ix, y)\} = Re\left\{ix + \frac{y}{a+ix} + \delta\right\} = \frac{yRe(a)}{|a+ix|^2} + \delta \leq -\frac{k}{2|a+ix|^2},$$

where

$$k = (nRe(a) - 2\delta)x^2 - 4\delta Im(a)x - 2\delta|a|^2 + nRe(a).$$

But in view of the value of δ given by (2.1), we know that k is a perfect square, which implies that

$$Re\{\psi(ix, y)\} \leq 0 \quad (x \in \mathbb{R}, y \leq -n\frac{1+x^2}{2}).$$

Now by Lemma 1.3, we deduce that

$$Re(p(z)) > 0 \quad (z \in U),$$

this completes the proof. □

Lemma 2.2. *Let f be in the class \mathcal{A}_n , $(L_a^{\lambda+1}(b, c, z)f)'(z) \neq 0$ for all $z \in U$ and $a > -1$. If*

$$(2.6) \quad Re\left\{1 + \frac{z(L_a^\lambda(b, c, z)f)''(z)}{(L_a^\lambda(b, c, z)f)'(z)}\right\} > -(n+a) + \frac{(1+a)}{4}$$

then

$$(2.7) \quad Re\left\{\sqrt{\frac{z(L_a^{\lambda+1}(b, c, z)f)''(z)}{(L_a^{\lambda+1}(b, c, z)f)'(z)} + (1+a)}\right\} > \frac{\sqrt{1+a}}{2}.$$

Proof. Let $f \in \mathcal{A}_n$ and set

$$(2.8) \quad p(z) = \frac{2}{\sqrt{1+a}} \sqrt{1+a + \frac{z(L_a^{\lambda+1}(b, c, z)f)''(z)}{(L_a^{\lambda+1}(b, c, z)f)'(z)}} - 1.$$

Then it is clear that p is analytic in U and $p(z) = 1 + b_n z^n + b_{n+1} z^{n+1} + \dots$. Taking the logarithmic differentiation in (2.8) and using identity (1.3) in the resulting equation, we obtain

$$(2.9) \quad \frac{(1+a)}{4}(1+p(z))^2 + 2\frac{zp'(z)}{1+p(z)} - a = 1 + \frac{z(L_a^\lambda(b, c, z)f)''(z)}{(L_a^\lambda(b, c, z)f)'(z)}.$$

To make use of Lemma 1.3 we consider the function

$$\psi(r, s) = \frac{(1+a)}{4}(1+r)^2 + \frac{2s}{1+r} + n - \frac{(1+a)}{4},$$

then by (2.6), (2.9) we have $Re\{\psi(p(z), zp'(z))\} > 0$. Furthermore for $x \in \mathbb{R}$ and $y < \frac{-n}{2}(1+x^2)$ we have

$$\begin{aligned} Re\{\psi(ix, y)\} &= \frac{(1+a)}{4}(1-x^2) + 2Re\frac{y}{1+ix} + n - \frac{(1+a)}{4} \\ &< \frac{(1+a)}{4}(1-x^2) - 2\frac{n(1+x^2)}{2(1+x^2)} + n - \frac{(1+a)}{4} \leq 0 \end{aligned}$$

Now in view of Lemma 1.3, we get that

$$Re(p(z)) > 0 \quad (z \in U),$$

this completes the proof. \square

By putting $\lambda = 0, b = c = 1$ and $a = 0$ in the Lemma 2.2 we obtain the following corollary which is generalization of well known result in the literature.

Corollary 2.3. *Let f be in the class \mathcal{A}_n and $f(z) \neq 0$ for all $z \in U$. If*

$$Re\left\{1 + \frac{zf''(z)}{f'(z)}\right\} > -n + \frac{1}{4},$$

then

$$Re\sqrt{\frac{zf'(z)}{f(z)}} > \frac{1}{2}.$$

Theorem 2.4. *Let $f, g \in \mathcal{A}_n, a \in \mathbb{C}, Re(a) > 0$ and $(L_a^{\lambda+1}(b, c, z)g)'(z) \neq 0$. If*

$$Re\left\{1 + \frac{z(L_a^\lambda(b, c, z)g)''(z)}{(L_a^\lambda(b, c, z)g)'(z)}\right\} > -\delta,$$

where δ is defined in (2.3). Then the relation

$$L_a^\lambda(b, c, z)f \prec L_a^\lambda(b, c, z)g$$

implies

$$L_a^{\lambda+1}(b, c, z)f \prec L_a^{\lambda+1}(b, c, z)g.$$

Proof. Let F, G and p be defined by

$$(2.10) \quad F(z) := L_a^{\lambda+1}(b, c, z)f, \quad G(z) := L_a^{\lambda+1}(b, c, z)g, \quad p(z) := 1 + \frac{G''(z)}{G'(z)}.$$

In view of Lemma 2.1, we have $Rep(z) > 0$. From definition of G and $L_a^\lambda(b, c, z)g$ and taking differentiation of G we conclude

$$(2.11) \quad G(z)\left(z \frac{G'(z)}{G(z)}\right) + a = (1 + a)L_a^\lambda(b, c, z)g(z).$$

But $Rep(z) > 0$ implies that the function G is convex (univalent) in U . We can assume, without loss of generality, that G is univalent on \bar{U} and that $G'(\zeta) \neq 0 (|\zeta| = 1)$. To prove $F \prec G$, we let function $L(z, t)$ be defined by

$$(2.12) \quad L(z, t) = \frac{a}{1+a}G(z) + \frac{1+t}{1+a}zG'(z) = a_1(t)z + \dots$$

It is easy to see that $L(z, t)$ is analytic in $|z| < 1$, for all $t \geq 0$. It is also continuously differentiable on $[0, \infty)$, for each $|z| < 1$. According to (2.11), we have $L(z, 0) = L_a^\lambda(b, c, z)g(z)$. From (2.12) we obtain

$$(2.13) \quad a_1(t) = \frac{\partial L}{\partial z}(0, t) = \left[\frac{a}{a+1} + \frac{1+t}{a+1}\right]G'(0) = 1 + \frac{t}{a+1} \neq 0$$

and $\lim_{t \rightarrow \infty} |a_1(t)| = \infty$. A simple calculation yields

$$Re\left(z \frac{\partial L(z, t)/\partial z}{\partial L(z, t)/\partial t}\right) = Rea + (1+t)Re\left(1 + \frac{G''(z)}{G'(z)}\right) = Rea + (1+t)Rep(z) > 0.$$

Since G is convex and normalized in the unit disk, therefore the following well-known growth and distortion sharp inequalities (see [9, p.45,48]) are true:

$$(2.14) \quad \frac{r}{1+r} \leq |G(z)| \leq \frac{r}{1-r}, \quad \text{if } |z| \leq r$$

and

$$(2.15) \quad \frac{r}{(1+r)^2} \leq |G'(z)| \leq \frac{r}{(1-r)^2}, \quad \text{if } |z| \leq r.$$

Now by making use of (2.14) and (2.15) in the relations (2.12) and (2.13) we obtain

$$\begin{aligned} \frac{|L(z, t)|}{|a_1(t)|} &\leq \frac{|a||G(z)| + (1+t)|zG'(z)|}{|a+1+t|} \\ &\leq \frac{r}{(1-r)^2} \frac{|a|+1+t}{|a+1+t|} \leq \frac{r}{(1-r)^2} \frac{|a|+1+t}{Re(a)+1+t}. \end{aligned}$$

If we define $h(t) = \frac{|a|+1+t}{Re(a)+1+t}$ it is easy to see that h is a decreasing function for $t \geq 0$, and so

$$h(t) \leq \frac{|a|+1}{Re(a)+1}.$$

Therefore from above inequality we conclude that

$$\frac{|L(z, t)|}{|a_1(t)|} \leq \frac{|a| + 1}{\operatorname{Re}(a) + 1} \frac{r}{(1 - r)^2} \quad |z| \leq r, t \geq 0.$$

Hence both assumptions of Lemma 1.2 hold. Now by Lemma 1.2 we obtain that $L(z, t)$ is a subordination chain. From Definition 1.1, it follows that

$$L_a^\lambda(b, c, z)g(z) = L(z, 0) \prec L(z, t) \quad (t \geq 0),$$

which implies

$$(2.16) \quad L(\zeta, t) \notin L(U, 0) = L_a^\lambda(b, c, z)g(U) \quad (\zeta \in \partial U, t \geq 0).$$

Suppose that $F \not\prec G$. Then there exists $z_0 \in U, \zeta_0 \in \partial U$ such that $F(z_0) = G(\zeta_0)$ and $F(|z| < |z_0|) \subset G(U)$. Hence by Jack's Lemma we have $z_0 F'(z_0) = (1 + t)\zeta_0 G'(\zeta_0)$ with $t \geq 0$. From (2.12), we obtain

$$\begin{aligned} L(\zeta_0, t) &= \frac{a}{1+a} G(\zeta_0) + \zeta_0 \frac{1+t}{1+a} G'(\zeta_0) \\ &= \frac{a}{1+a} F(z_0) + z_0 \frac{1}{1+a} F'(z_0) \\ &= L_a^\lambda(b, c, z)f(z_0) \in L_a^\lambda(b, c, z)g(U). \end{aligned}$$

But this contradicts (2.16), and thus we deduce that $F \prec G$. \square

By setting $b = c = 1$ and $\lambda = 0$ in the Theorem 2.4 we obtain.

Corollary 2.5. *Let $a \in \mathbb{C}, \operatorname{Re}(a) > 0$ and $f, g \in \mathcal{A}_n$. If*

$$\operatorname{Re}\left(1 + \frac{g''(z)}{g'(z)}\right) > -n \frac{1 + |a|^2 - |1 - a^2|}{4\operatorname{Re}(a)},$$

then

$$f \prec g \Rightarrow \frac{1+a}{z^a} \int_0^z t^{a-1} f(t) dt \prec \frac{1+a}{z^a} \int_0^z t^{a-1} g(t) dt.$$

Corollary 2.6. *Let $a > 0$ and $f, g \in \mathcal{A}_n$. If*

$$\operatorname{Re}\left(1 + \frac{z(L_a^\lambda(b, c, z)g)''(z)}{(L_a^\lambda(b, c, z)g)'(z)}\right) > -\delta_1,$$

then

$$L_a^\lambda(b, c, z)f \prec L_a^\lambda(b, c, z)g \Rightarrow L_a^{\lambda+1}(b, c, z)f \prec L_a^{\lambda+1}(b, c, z)g,$$

where

$$\delta_1 = \begin{cases} -\frac{an}{2} & 0 < a < 1 \\ -\frac{n}{2a} & a \geq 1. \end{cases}$$

By putting $b = c = 1$ and $\lambda = 0$ in the Corollary 2.6 we have

Corollary 2.7. *Let $a > 0$ and $f, g \in \mathcal{A}_n$. If*

$$\operatorname{Re}\left(1 + \frac{zg''(z)}{g'(z)}\right) > -\delta_1,$$

then

$$f \prec g \Rightarrow \frac{1+a}{z^a} \int_0^z t^{a-1} f(t) dt \prec \frac{1+a}{z^a} \int_0^z t^{a-1} g(t) dt,$$

where

$$\delta_1 = \begin{cases} -\frac{an}{2} & 0 < a < 1 \\ -\frac{n}{2a} & a \geq 1. \end{cases}$$

By using similar arguments as in the proofs of Lemma 2.1 and Theorem 2.4 we obtain the following results and we omit details.

Lemma 2.8. *Let f be in the class \mathcal{A}_n , $(L_a^\lambda(b, c, z)f)'(z) \neq 0$ for all $z \in U$ and $b \in \mathbb{C}$, $\operatorname{Re}(b) > 1$. If*

$$(2.17) \quad \operatorname{Re} \left\{ 1 + \frac{z(L_a^\lambda(b+1, c, z)f)''(z)}{(L_a^\lambda(b+1, c, z)f)'(z)} \right\} > -\delta_2$$

then

$$(2.18) \quad \operatorname{Re} \left\{ 1 + \frac{z(L_a^\lambda(b, c, z)f)''(z)}{(L_a^\lambda(b, c, z)f)'(z)} \right\} > 0$$

where

$$(2.19) \quad \delta_2 = n \frac{1 + |b-1|^2 - |b(2-b)|}{4\operatorname{Re}(b-1)}.$$

Theorem 2.9. *Let $f, g \in \mathcal{A}_n$, $(L_a^\lambda(b, c, z)g)'(z) \neq 0$ and $b \in \mathbb{C}$, $\operatorname{Re}(b) > 1$. If*

$$\operatorname{Re} \left\{ 1 + \frac{z(L_a^\lambda(b+1, c, z)g)''(z)}{(L_a^\lambda(b+1, c, z)g)'(z)} \right\} > -\delta_2,$$

where δ_2 is defined in (2.16). Then

$$L_a^\lambda(b+1, c, z)f \prec L_a^\lambda(b+1, c, z)g \Rightarrow L_a^\lambda(b, c, z)f \prec L_a^\lambda(b, c, z)g.$$

By putting $b = c = 2$ and $\lambda = a = 1$ in the Lemma 2.8 we get the following result.

Corollary 2.10. *Let $f(z) = z + \sum_{k=n+1}^\infty a_k z^k \in \mathcal{A}_n$ satisfies the condition*

$$\operatorname{Re}\left(1 + \frac{zf''(z)}{f'(z)}\right) > \frac{-n}{2},$$

then the function $h(z) = \frac{2}{z} \int_0^z f(t) dt = z + \sum_{k=n+1}^\infty \frac{2}{k+1} a_k z^k$ is convex.

Theorem 2.11. *Let $f, g \in \mathcal{A}_n, (L_a^{\lambda+1}(b, c, z)g)'(z) \neq 0$ and $a \in \mathbb{C}$ with $Re(a) > 0$. If*

$$Re\left\{1 + \frac{z(L_a^\lambda(b, c, z)g)''(z)}{(L_a^\lambda(b, c, z)g)'(z)}\right\} > -\delta$$

where δ is defined in (2.3) and if $L_a^\lambda(b, c, z)f$ is univalent and $L_a^{\lambda+1}(b, c, z)f \in Q(0)$ then

$$L_a^\lambda(b, c, z)g \prec L_a^\lambda(b, c, z)f \Rightarrow L_a^{\lambda+1}(b, c, z)g \prec L_a^{\lambda+1}(b, c, z)f.$$

and $L_a^{\lambda+1}(b, c, z)g$ is the best subordinate.

Proof. Let the functions F, G and p be defined by (2.10). From Lemma 2.1 we get

$$Re(p(z)) > 0 \quad (z \in U).$$

Next, to arrive at our desired result, we show that $G \prec F$. For this, let the functions $L(z, t)$ and $\phi(r, s)$ are defined by

$$(2.20) \quad \begin{aligned} L(z, t) &= \phi(G(z), tzG'(z)) \\ &= \frac{a}{1+a}G(z) + \frac{t}{1+a}zG'(z) = a_1(t)z + \dots \quad (z \in U). \end{aligned}$$

Then it is easy to see that $L(z, t)$ is analytic in $|z| < 1$, for all $t \geq 0$. It is also continuously differentiable on $[0, \infty)$, for each $|z| < 1$. From (2.20) we obtain

$$(2.21) \quad a_1(t) = \frac{\partial L}{\partial z}(0, t) = \left[\frac{a}{a+1} + \frac{t}{a+1}\right]G'(0) = \frac{a+t}{a+1} \neq 0$$

and $\lim_{t \rightarrow \infty} |a_1(t)| = \infty$. A simple calculation yields

$$Re\left(z \frac{\partial L(z, t)/\partial z}{\partial L(z, t)/\partial t}\right) = Rea + tRe\left(1 + \frac{G''(z)}{G'(z)}\right) = Rea + tRep(z) > 0.$$

Since G is convex and normalized in the unit disk, therefore by using similar argument of Theorem 2.4 we obtain

$$\begin{aligned} \frac{|L(z, t)|}{|a_1(t)|} &\leq \frac{|a||G(z)| + t|zG'(z)|}{|a+t|} \\ &\leq \frac{r}{(1-r)^2} \frac{|a|+t}{|a+t|} \leq \frac{r}{(1-r)^2} \frac{|a|+t}{Re(a)+t}. \end{aligned}$$

If we define $k(t) = \frac{|a|+t}{Re(a)+t}$ it is easy to see that k is a decreasing function for $t \geq 0$, and so

$$k(t) \leq \frac{|a|+1}{Re(a)+1}.$$

Therefore from above inequality we conclude that

$$\frac{|L(z, t)|}{|a_1(t)|} \leq \frac{|a|}{Re(a)} \frac{r}{(1-r)^2} \quad |z| \leq r, t \geq 0.$$

Hence both assumptions of Lemma 1.2 hold. Now by Lemma 1.2 we obtain that $L(z, t)$ is a subordination chain. Now following the same as proof of Theorem 2.4 we conclude $G \prec F$ and Moreover, G is best subordinate. \square

By suitably combining Theorems 2.4 and 2.11 we obtain the following Sandwich type theorem.

Corollary 2.12. *Let $f, g_k \in \mathcal{A}_n (k = 1, 2)$ and $a \in \mathbb{C}$ with $Re(a) > 0$. If*

$$Re\left(1 + \frac{z(L_a^\lambda(b, c, z)g_k)''(z)}{(L_a^\lambda(b, c, z)g_k)'(z)}\right) > -\delta$$

where δ is given by (2.3) and if the function $(L_a^\lambda(b, c, z)f)(z)$ is univalent and $(L_a^{\lambda+1}(b, c, z)f)(z) \in Q(0)$, then

$$L_a^\lambda(b, c, z)g_1 \prec L_a^\lambda(b, c, z)f \prec L_a^\lambda(b, c, z)g_2$$

implies

$$L_a^{\lambda+1}(b, c, z)g_1 \prec L_a^{\lambda+1}(b, c, z)f \prec L_a^{\lambda+1}(b, c, z)g_2.$$

By setting $b = c = 1$ and $\lambda = 0$ in the Corollary 2.12 we obtain

Corollary 2.13. *Let $a \in \mathbb{C}, Re(a) > 0$ and $f, g_k \in \mathcal{A}_n (k = 1, 2)$. If f is univalent and $L_a^1(1, 1, z)f \in Q(0)$. Then*

$$Re\left(1 + \frac{zg_k''(z)}{g_k'(z)}\right) > -n \frac{1 + |a|^2 - |1 - a^2|}{4Re(a)}$$

implies

$$g_1 \prec f \prec g_2 \Rightarrow \frac{1+a}{z^a} \int_0^z t^{a-1} g_1(t) dt \prec \frac{1+a}{z^a} \int_0^z t^{a-1} f(t) dt \prec \frac{1+a}{z^a} \int_0^z t^{a-1} g_2(t) dt.$$

Corollary 2.14. *Let $a > 0$ and $f, g_k \in \mathcal{A}_n, k = 1, 2$. If f is univalent and $L_a^1(1, 1, z)f \in Q(0)$. Then*

$$Re\left(1 + \frac{zg_k''(z)}{g_k'(z)}\right) > \begin{cases} -\frac{an}{2} & 0 < a < 1 \\ -\frac{n}{2a} & a \geq 1, \end{cases}$$

implies

$$g_1 \prec f \prec g_2 \Rightarrow \frac{1+a}{z^a} \int_0^z t^{a-1} g_1(t) dt \prec \frac{1+a}{z^a} \int_0^z t^{a-1} f(t) dt \prec \frac{1+a}{z^a} \int_0^z t^{a-1} g_2(t) dt.$$

Acknowledgments

The authors would like to thank the referee for their careful reading and making some valuable comments which have improved the presentation of this paper.

REFERENCES

- [1] R. Aghalary, A. Ebadian and Z- G. Wang, Subordination and superordination results involving certain convolution operators, *Bull. Iranian Math. Soc.* **36** (2010), no. 1, 137–147.
- [2] M. K. Aouf and T. Bulboaca, Subordination and superordination properties of multivalent functions defined by certain integral operator, *J. Franklin Institute* **347** (2010), no. 3, 641–653.
- [3] N. E. Cho, O. S. Kwon, S. Owa and H. M. Srivastava, A class of integral operators preserving subordination and superordination for meromorphic functions, *Appl. Math. comput.* **193** (2007), no. 2, 463–474.
- [4] T. Bulboaca, M. K. Aouf and R. M. Ashwah, Subordination properties of multivalent functions defined by certain integral operator, *Banach J. Math. Anal.* **6** (2012), no. 2, 69–85.
- [5] Y. Komatu, Analytical prolongation of a family of operators, *Math.(Cluj)* **32** (1990), 141–145.
- [6] S. S. Miller and P. T. Mocanu, Differential subordination and univalent functions, *Michigan Math. J.* **28** (1981) 157–171.
- [7] S. S. Miller and P. T. Mocanu, Differential subordinations Monographs and Textbooks in pure and Applied Mathematic, 225, Marcel Dekker, Inc., New York, 2000.
- [8] S. Miller and P. T. Mocanu, Subordinants of differential superordinations, *Complex Var. Theory Appl.* **48** (2003), no. 10, 815–826.
- [9] C. Pommerenke, Univalent functions, Vanderhoeck and Rupercht, Gottingen, 1975.

(Rasoul. Aghalary) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, URMIA UNIVERSITY, URMIA, IRAN

E-mail address: raghalary@yahoo.com and r.aghalary@urmia.ac.ir

(Ali Ebadian) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, URMIA UNIVERSITY, URMIA, IRAN

E-mail address: a.ebadian@urmai.ac.ir

(Mastore Mafakheri) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, URMIA UNIVERSITY, URMIA, IRAN

E-mail address: M.Mafakheri85@yahoo.com