


ISSN: 1017-060X (Print)



ISSN: 1735-8515 (Online)

Special Issue of the
**Bulletin of the
Iranian Mathematical Society**

in Honor of Professor Heydar Radjavi's 80th Birthday

Vol. 41 (2015), No. 7, pp. 107–116

Title:

Additivity of maps preserving Jordan η_* -products on C^* -algebras

Author(s):

A. Taghavi, H. Rohi and V. Darvish

Published by Iranian Mathematical Society
<http://bims.ims.ir>

ADDITIVITY OF MAPS PRESERVING JORDAN η_* -PRODUCTS ON C^* -ALGEBRAS

A. TAGHAVI*, H. ROHI AND V. DARVISH

(Communicated by Peter Rosenthal)

Dedicated to Professor Heydar Radjavi on his 80th birthday

ABSTRACT. Let \mathcal{A} and \mathcal{B} be two C^* -algebras such that \mathcal{B} is prime. In this paper, we investigate the additivity of maps Φ from \mathcal{A} onto \mathcal{B} that are bijective, unital and that satisfy $\Phi(AP + \eta PA^*) = \Phi(A)\Phi(P) + \eta\Phi(P)\Phi(A)^*$, for all $A \in \mathcal{A}$ and $P \in \{P_1, I_{\mathcal{A}} - P_1\}$ where P_1 is a nontrivial projection in \mathcal{A} . If η is a non-zero complex number such that $|\eta| \neq 1$, then Φ is additive. Moreover, if η is rational, then Φ is $*$ -additive.

Keywords: Maps preserving Jordan η_* -product, Additive, Prime C^* -algebras.

MSC(2010): Primary: 47B48; Secondary: 46L10.

1. Introduction

Let \mathcal{R} and \mathcal{R}' be rings. We say that the map $\Phi : \mathcal{R} \rightarrow \mathcal{R}'$ “preserves products” or is “multiplicative” if $\Phi(AB) = \Phi(A)\Phi(B)$ for all $A, B \in \mathcal{R}$. The question of when a multiplicative map is additive was discussed by several authors: see [20] and references therein. Motivated by this, many authors paid more attention to maps on rings (and algebras) that preserve the Lie product $[A, B] = AB - BA$ or the Jordan product $A \circ B = AB + BA$. (See, for example, [1, 2, 6, 10, 13, 16–18, 24, 25].) These results show that, in some sense, the Jordan product or Lie product structure is enough to determine the ring or algebraic structure. Historically, many mathematicians devoted themselves to the study of additive or linear Jordan or Lie product preservers between rings or operator algebras. Such maps are called Jordan homomorphisms or Lie homomorphisms respectively. See, for example, [3–5, 11, 12, 14, 20–22].

Let \mathcal{R} be a $*$ -ring and η be a non-zero complex scalar. For $A, B \in \mathcal{R}$, denote the Jordan η_* -product of A and B by $A \bullet_{\eta} B = AB + \eta BA^*$ and the

Article electronically published on December 31, 2015.

Received: 26 November 2014, Accepted: 18 April 2015.

*Corresponding author.

Lie η_* -product by $[A, B]_*^\eta = AB - \eta BA^*$. In particular, $A \bullet_1 B = AB + BA^*$ and $[A, B]_*^1 = AB - BA^*$ are the Jordan 1_* -product and the Lie 1_* -product, respectively. These products are playing an increasingly important role, and have been studied by many authors (for example, see [7, 23, 26]). A fundamental problem is to determine the conditions under which maps preserving Jordan η_* -products or Lie η_* -products on rings or algebras are isomorphisms. In [8], J. Cui and C. K. Li proved that a bijective map on a factor von Neumann algebra which preserves the Lie 1_* -product ($[A, B]_*^1$) must be a $*$ -isomorphism. Moreover, in [15] C. Li et al, discussed nonlinear bijective maps preserving Jordan 1_* -product ($A \bullet_1 B$). They proved that such a mapping on factor von Neumann algebras is also a $*$ -ring isomorphism. These two articles also discussed new products for arbitrary operators on factor von Neumann algebras. In addition, A. Taghavi et al [27], proved that a bijective unital map (not necessarily linear) on a prime C^* -algebra which preserves both the Lie 1_* -product and the Jordan 1_* -product in the case where one of the operators is a projection must be $*$ -additive (i.e., additive and star-preserving). In a recent paper [9], L. Dai and F. Lu proved that a bijective map on a von Neumann algebra which preserves the Jordan η_* -product is a linear $*$ -isomorphism if η is not real, it is and is a sum of a linear $*$ -isomorphism and a conjugate linear $*$ -isomorphism if η is real.

In this paper, we will consider when a bijective unital map on a prime C^* -algebra which preserves the Jordan η_* -product $\Phi(A \bullet_\eta P) = \Phi(A) \bullet_\eta \Phi(P)$ is additive. We denote the real part of an operator T by $\Re(T)$; i.e., $\Re(T) = \frac{T+T^*}{2}$. A C^* -algebra \mathcal{A} is prime if $AAB = 0$ for $A, B \in \mathcal{A}$ implies that either $A = 0$ or $B = 0$.

2. Main results

We need the following lemmas.

Lemma 2.1. *Let \mathcal{A} and \mathcal{B} be two C^* -algebras and $\Phi : \mathcal{A} \rightarrow \mathcal{B}$ be a map which satisfies $\Phi(A \bullet_\eta P) = \Phi(A) \bullet_\eta \Phi(P)$ for all $A \in \mathcal{A}$ and some $P \in \mathcal{A}$ where η is a non-zero complex number such that $|\eta| \neq 1$. Let A, B and T be in \mathcal{A} such that $\Phi(T) = \Phi(A) + \Phi(B)$. Then we have*

$$(2.1) \quad \Phi(T \bullet_\eta P) = \Phi(A \bullet_\eta P) + \Phi(B \bullet_\eta P).$$

Proof. Multiply the equations $\Phi(T) = \Phi(A) + \Phi(B)$ and $\Phi(T)^* = \Phi(A)^* + \Phi(B)^*$ by $\Phi(P)$ from the right and $\eta\Phi(P)$ from the left. We get

$$\Phi(T)\Phi(P) = \Phi(A)\Phi(P) + \Phi(B)\Phi(P),$$

and

$$\eta\Phi(P)\Phi(T)^* = \eta\Phi(P)\Phi(A)^* + \eta\Phi(P)\Phi(B)^*.$$

By adding the two equations, we get

$$\Phi(T)\Phi(P) + \eta\Phi(P)\Phi(T)^* = \Phi(A)\Phi(P) + \eta\Phi(P)\Phi(A)^* + \Phi(B)\Phi(P) + \eta\Phi(P)\Phi(B)^*.$$

□

Lemma 2.2. *Let \mathcal{A} be a C^* -algebra. Suppose $T \in \mathcal{A}$ and η is a non-zero complex number such that $|\eta| \neq 1$. If $T + \eta T^* = 0$, then $T = 0$.*

Proof. Let $T + \eta T^* = 0$. Then $T^* + \bar{\eta}T = 0$. By an easy computation we have $(1 - \bar{\eta}\eta)T = 0$. Thus $T = 0$. □

Lemma 2.3. *Let \mathcal{A} and \mathcal{B} be two C^* -algebras with identities and $\Phi : \mathcal{A} \rightarrow \mathcal{B}$ be a bijective map which satisfies $\Phi(A \bullet_\eta P) = \Phi(A) \bullet_\eta \Phi(P)$ for all $A \in \mathcal{A}$ and $P \in \{P_1, I - P_1\}$, where P_1 is a nontrivial projection in \mathcal{A} and η is a non-zero complex number such that $|\eta| \neq 1$. Then $\Phi(0) = 0$.*

Proof. Let $\Phi(T) = 0$. We prove that $T = 0$. To show this, apply Lemma 2.1 to $\Phi(T) = 0$ for P_1 and $P_2 = I - P_1$. This gives

$$\Phi(T \bullet_\eta P_1) = 0,$$

and

$$\Phi(T \bullet_\eta P_2) = 0.$$

So, by injectivity of Φ , we obtain $T \bullet_\eta P_1 = T \bullet_\eta P_2$. By the definition of Jordan η_* -product, we have $TP_1 + \eta P_1 T^* = TP_2 + \eta P_2 T^*$; i.e., $(TP_1 - TP_2) + \eta(TP_1 - TP_2)^* = 0$. Now, applying Lemma 2.2, we obtain $TP_1 = TP_2$. We multiply the latter equation by P_2 on the right; it follows that $TP_2 = 0$. Thus $TP_1 = 0$ also. Now we put $I - P_2$ instead of P_1 in $TP_1 = 0$, obtaining $T(I - P_2) = 0$. Therefore $T = 0$ since $TP_2 = 0$. □

Our main theorem is as follows:

Main Theorem. *Let \mathcal{A} and \mathcal{B} be two C^* -algebras such that \mathcal{B} is prime, with $I_{\mathcal{A}}$ and $I_{\mathcal{B}}$ their respective identities. Let $\Phi : \mathcal{A} \rightarrow \mathcal{B}$ be a unital bijective map which satisfies $\Phi(A \bullet_\eta P) = \Phi(A) \bullet_\eta \Phi(P)$ for all $A \in \mathcal{A}$ and $P \in \{P_1, I_{\mathcal{A}} - P_1\}$, where P_1 is a nontrivial projection in \mathcal{A} and η is a non-zero complex number such that $|\eta| \neq 1$. Then Φ is additive. Moreover, if η is rational, then Φ is $*$ -additive.*

Proof of Main Theorem. Let $P_2 = I_{\mathcal{A}} - P_1$ and let $\mathcal{A}_{ij} = P_i \mathcal{A} P_j$, $i, j = 1, 2$. Then $\mathcal{A} = \sum_{i,j=1}^2 \mathcal{A}_{ij}$. For every $A \in \mathcal{A}$ we may write $A = A_{11} + A_{12} + A_{21} + A_{22}$. In all that follows, when we write A_{ij} , it indicates that $A_{ij} \in \mathcal{A}_{ij}$. To show additivity of Φ on \mathcal{A} we will use the above partition of \mathcal{A} and establish that Φ is additive on each \mathcal{A}_{ij} , $i, j = 1, 2$. □

Claim 2.4. *For every $A_{11} \in \mathcal{A}_{11}$ and $D_{22} \in \mathcal{A}_{22}$, we have*

$$\Phi(A_{11} + D_{22}) = \Phi(A_{11}) + \Phi(D_{22}).$$

Since Φ is surjective, we can find an element $T = T_{11} + T_{12} + T_{21} + T_{22} \in \mathcal{A}$ such that

$$(2.2) \quad \Phi(T) = \Phi(A_{11}) + \Phi(D_{22}).$$

We now show $T = A_{11} + D_{22}$. We apply Lemma 2.1 to (2.2) for P_1 . We write

$$\Phi(T \bullet_\eta P_1) = \Phi(A_{11} \bullet_\eta P_1) + \Phi(D_{22} \bullet_\eta P_1),$$

so

$$\Phi(T_{11} + T_{21} + \eta T_{11}^* + \eta T_{21}^*) = \Phi(A_{11} + \eta A_{11}^*).$$

By the injectivity of Φ , we get $T_{11} + T_{21} + \eta T_{11}^* + \eta T_{21}^* = A_{11} + \eta A_{11}^*$. Multiply the latter equation by P_2 on the right and left side, respectively, to obtain $T_{21} = T_{21}^* = 0$. Thus we have $T_{11} + \eta T_{11}^* = A_{11} + \eta A_{11}^*$, or $(T_{11} - A_{11}) + \eta(T_{11} - A_{11})^* = 0$. By Lemma 2.2, we get $T_{11} = A_{11}$. Similarly, applying Lemma 2.1 to (2.2) for P_2 gives

$$\Phi(T \bullet_\eta P_2) = \Phi(A_{11} \bullet_\eta P_2) + \Phi(D_{22} \bullet_\eta P_2),$$

and hence

$$\Phi(T_{12} + T_{22} + \eta T_{12}^* + \eta T_{22}^*) = \Phi(D_{22} + \eta D_{22}^*).$$

By the injectivity of Φ , we get $T_{12} + T_{22} + \eta T_{12}^* + \eta T_{22}^* = D_{22} + \eta D_{22}^*$. Multiply the latter equation by P_1 on the right and left side respectively, to obtain $T_{12} = T_{12}^* = 0$. Thus we have $T_{22} + \eta T_{22}^* = D_{22} + \eta D_{22}^*$, or $(T_{22} - D_{22}) + \eta(T_{22} - D_{22})^* = 0$. By Lemma 2.2, we get $T_{22} = D_{22}$.

Claim 2.5. For every $B_{12} \in \mathcal{A}_{12}$, $C_{21} \in \mathcal{A}_{21}$, we have

$$\Phi(B_{12} + C_{21}) = \Phi(B_{12}) + \Phi(C_{21}).$$

Let $T = T_{11} + T_{12} + T_{21} + T_{22} \in \mathcal{A}$ be such that

$$(2.3) \quad \Phi(T) = \Phi(B_{12}) + \Phi(C_{21}).$$

Applying Lemma 2.1 to (2.3) for P_1 , we have obtain

$$\begin{aligned} \Phi(T \bullet_\eta P_1) &= \Phi(B_{12} \bullet_\eta P_1) + \Phi(C_{21} \bullet_\eta P_1) \\ &= \Phi(C_{21} \bullet_\eta P_1). \end{aligned}$$

Thus, by the injectivity of Φ , we have $T \bullet_\eta P_1 = C_{21} \bullet_\eta P_1$. It follows that

$$T_{11} + T_{21} + \eta T_{11}^* + \eta T_{21}^* = C_{21} + \eta C_{21}^*.$$

Multiplying the above equation by P_2 on the left, gives $T_{21} = C_{21}$ and $T_{11} = 0$. Similarly, we can obtain $T_{12} = B_{12}$ and $T_{22} = 0$ by applying Lemma 2.1 to (2.3) for P_2 .

Claim 2.6. For every $A_{11} \in \mathcal{A}_{11}$ and $B_{12} \in \mathcal{A}_{12}$, we have

$$\Phi(A_{11} + B_{12}) = \Phi(A_{11}) + \Phi(B_{12}).$$

Since Φ is surjective, we can find an element $T = T_{11} + T_{12} + T_{21} + T_{22} \in \mathcal{A}$ such that

$$(2.4) \quad \Phi(T) = \Phi(A_{11}) + \Phi(B_{12}).$$

We show that $T = A_{11} + B_{12}$. We apply Lemma 2.1 to (2.4) for P_1 , we can write $T_{11} = A_{11}$ and $T_{21} = 0$.

Similarly, we apply Lemma 2.1 to (2.4) for P_2 , we will have $T_{12} = B_{12}$ and $T_{22} = 0$. So, $T = A_{11} + B_{12}$.

Note that the equation $\Phi(C_{21} + D_{22}) = \Phi(C_{21}) + \Phi(D_{22})$, where $C_{21} \in \mathcal{A}_{21}$ and $D_{22} \in \mathcal{A}_{22}$, can be obtained as above.

Claim 2.7. *For every $A_{11} \in \mathcal{A}_{11}$ and $C_{21} \in \mathcal{A}_{21}$, we have*

$$\Phi(A_{11} + C_{21}) = \Phi(A_{11}) + \Phi(C_{21}).$$

Since Φ is surjective, we can find an element $T = T_{11} + T_{12} + T_{21} + T_{22} \in \mathcal{A}$ such that

$$(2.5) \quad \Phi(T) = \Phi(A_{11}) + \Phi(C_{21}).$$

We show that $T = A_{11} + C_{12}$. We apply Lemma 2.1 to (2.5) for P_2 ; then we have $\Phi(T \bullet_\eta P_2) = 0$. This means that $\Phi(T_{12} + T_{22} + \eta T_{12}^* + \eta T_{22}^*) = 0$. By Lemma 2.3 and Lemma 2.2, we obtain $T_{12} = T_{22} = 0$. On the other hand, we apply Lemma 2.1 to (2.5) for P_1 . Then, by Claim 2.6, we have

$$\begin{aligned} \Phi(T \bullet_\eta P_1) &= \Phi(A_{11} \bullet_\eta P_1) + \Phi(C_{21} \bullet_\eta P_1) \\ &= \Phi(A_{11} + \eta A_{11}^*) + \Phi(C_{21} + \eta C_{21}^*) \\ &= \Phi(A_{11} + \eta A_{11}^*) + \Phi(2\Re(C_{21}^*) + (\eta - 1)C_{21}^*) \\ &= \Phi(A_{11} + \eta A_{11}^* + 2\Re(C_{21}^*) + (\eta - 1)C_{21}^*) \end{aligned}$$

We get the following:

$$\Phi(T_{11} + T_{21} + \eta T_{11}^* + T_{21}^*) = \Phi(A_{11} + \eta A_{11}^* + C_{21} + \eta C_{21}^*).$$

Now we use Lemma 2.2 to obtain $T_{11} = A_{11}$ and $T_{21} = C_{21}$.

Note that the fact that $\Phi(B_{12} + D_{22}) = \Phi(B_{12}) + \Phi(D_{22})$, where $B_{12} \in \mathcal{A}_{12}$ and $D_{22} \in \mathcal{A}_{22}$, can be obtained as above.

Claim 2.8. *For every $A_{ij}, B_{ij} \in \mathcal{A}_{ij}$ such that $1 \leq i \neq j \leq 2$, we have*

$$\Phi(A_{ij} + B_{ij}) = \Phi(A_{ij}) + \Phi(B_{ij}).$$

Let $T = T_{11} + T_{12} + T_{21} + T_{22} \in \mathcal{A}$ be such that

$$(2.6) \quad \Phi(T) = \Phi(A_{ij}) + \Phi(B_{ij}).$$

Applying Lemma 2.1 to (2.6) for P_i , we get

$$\Phi(TP_i + \eta P_i T^*) = \Phi(A_{ij}P_i + \eta P_i A_{ij}^*) + \Phi(B_{ij}P_i + \eta P_i B_{ij}^*) = \Phi(0) = 0.$$

Therefore, $\Phi(T_{ii} + T_{ji} + \eta T_{ii}^* + \eta T_{ji}^*) = 0$. So, by Lemma 2.2, we have $T_{ii} = T_{ji} = 0$. On the other hand, we apply Lemma 2.1 to (2.6) for P_j again. By

Claim 2.5, it we see that

$$\begin{aligned}
\Phi(TP_j + \eta P_j T^*) &= \Phi(A_{ij}P_j + \eta P_j A_{ij}^*) + \Phi(B_{ij}P_j + \eta P_j B_{ij}^*) \\
&= \Phi(A_{ij} + \eta A_{ij}^*) + \Phi(B_{ij} + \eta B_{ij}^*) \\
&= \Phi(2\Re(A_{ij}^*) + (\eta - 1)A_{ij}^*) + \Phi(2\eta\Re(B_{ij}) + (1 - \eta)B_{ij}) \\
&= \Phi(2\Re(A_{ij}^*) + (\eta - 1)A_{ij}^* + 2\eta\Re(B_{ij}) + (1 - \eta)B_{ij}) \\
&= \Phi(A_{ij} + \eta A_{ij}^* + B_{ij} + \eta B_{ij}^*).
\end{aligned}$$

So,

$$T_{ij} + T_{jj} + \eta T_{ij}^* + \eta T_{jj}^* = A_{ij} + \eta A_{ij}^* + B_{ij} + \eta B_{ij}^*.$$

By Lemma 2.2, we have $T_{jj} = 0$ and $T_{ij} = A_{ij} + B_{ij}$.

Claim 2.9. For every $A_{11} \in \mathcal{A}_{11}$, $B_{12} \in \mathcal{A}_{12}$, $C_{21} \in \mathcal{A}_{21}$, we have

$$\Phi(A_{11} + B_{12} + C_{21}) = \Phi(A_{11}) + \Phi(B_{12}) + \Phi(C_{21}).$$

Let $T = T_{11} + T_{12} + T_{21} + T_{22} \in \mathcal{A}$ be such that

$$(2.7) \quad \Phi(T) = \Phi(A_{11}) + \Phi(B_{12}) + \Phi(C_{21}).$$

Applying Lemma 2.1 to (2.7) for P_2 , we have

$$\begin{aligned}
\Phi(T \bullet_\eta P_2) &= \Phi(A_{11} \bullet_\eta P_2) + \Phi(B_{12} \bullet_\eta P_2) + \Phi(C_{21} \bullet_\eta P_2) \\
&= \Phi(B_{12} \bullet_\eta P_2).
\end{aligned}$$

Thus, by injectivity of Φ , we have $T \bullet_\eta P_2 = B_{12} \bullet_\eta P_2$. It follows that

$$T_{22} + T_{12} + \eta T_{22}^* + \eta T_{12}^* = B_{12} + \eta B_{12}^*.$$

Multiply the above equation by P_1 on the left to get $T_{12} = B_{12}$ and $T_{22} = 0$. Similarly, we apply Lemma 2.1 to (2.7) for P_1 . By Claim 2.6, we have the following

$$\begin{aligned}
\Phi(T \bullet_\eta P_1) &= \Phi(A_{11} \bullet_\eta P_1) + \Phi(B_{12} \bullet_\eta P_1) + \Phi(C_{21} \bullet_\eta P_1) \\
&= \Phi(A_{11} \bullet_\eta P_1) + \Phi(C_{21} \bullet_\eta P_1) \\
&= \Phi(A_{11} + \eta A_{11}^*) + \Phi(C_{21} + \eta C_{21}^*) \\
&= \Phi(A_{11} + \eta A_{11}^*) + \Phi(2\Re(C_{21}^*) + (\eta - 1)C_{21}^*) \\
&= \Phi(A_{11} + \eta A_{11}^* + 2\Re(C_{21}^*) + (\eta - 1)C_{21}^*) \\
&= \Phi(A_{11} + \eta A_{11}^* + C_{21} + \eta C_{21}^*)
\end{aligned}$$

Injectivity of Φ implies $T_{11} + T_{21} + \eta T_{11}^* + \eta T_{21}^* = A_{11} + C_{21} + \eta A_{11}^* + \eta C_{21}^*$. We have $T_{11} = A_{11}$ and $T_{21} = C_{21}$.

Claim 2.10. For every $A_{11} \in \mathcal{A}_{11}$, $B_{12} \in \mathcal{A}_{12}$, $C_{21} \in \mathcal{A}_{21}$ and $D_{22} \in \mathcal{A}_{22}$, we have

$$\Phi(A_{11} + B_{12} + C_{21} + D_{22}) = \Phi(A_{11}) + \Phi(B_{12}) + \Phi(C_{21}) + \Phi(D_{22}).$$

Assume $T = T_{11} + T_{12} + T_{21} + T_{22}$. Then

$$(2.8) \quad \Phi(T) = \Phi(A_{11}) + \Phi(B_{12}) + \Phi(C_{21}) + \Phi(D_{22}).$$

Applying Lemma 2.1 to (2.8) for P_1 , Claim 2.5 and Claim 2.9, we obtain

$$\begin{aligned} \Phi(T \bullet_\eta P_1) &= \Phi(A_{11} \bullet_\eta P_1) + \Phi(B_{12} \bullet_\eta P_1) + \Phi(C_{21} \bullet_\eta P_1) + \Phi(D_{22} \bullet_\eta P_1) \\ &= \Phi(A_{11} \bullet_\eta P_1) + \Phi(C_{21} \bullet_\eta P_1) \\ &= \Phi(A_{11} + \eta A_{11}^*) + \Phi(C_{21} + \eta C_{21}^*) \\ &= \Phi(A_{11} + \eta A_{11}^*) + \Phi(C_{21}) + \Phi(\eta C_{21}^*) \\ &= \Phi(A_{11} + \eta A_{11}^* + C_{21} + \eta C_{21}^*) \end{aligned}$$

Since Φ is injective we have $T_{11} + T_{21} + \eta T_{11}^* + \eta T_{21}^* = A_{11} + C_{21} + \eta A_{11}^* + \eta C_{21}^*$. We obtain $T_{11} = A_{11}$ and $T_{21} = C_{21}$. Similarly, apply Lemma 2.1 to (2.8) for P_2 and the same computation as above to easily obtain $T_{12} = B_{12}$ and $T_{22} = D_{22}$. So, $\Phi(A_{11} + B_{12} + C_{21} + D_{22}) = \Phi(A_{11}) + \Phi(B_{12}) + \Phi(C_{21}) + \Phi(D_{22})$.

Lemma 2.11. *Let Φ satisfy the assumptions of the Main Theorem. Then, for every $A \in \mathcal{A}$ we have the following*

$$\Phi(A \bullet_\eta I) = \Phi(A) \bullet_\eta \Phi(I).$$

Proof. The definition of the Jordan η_* -product implies that

$$(2.9) \quad \Phi(A \bullet_\eta P_1) = \Phi(A) \bullet_\eta \Phi(P_1),$$

and

$$(2.10) \quad \Phi(A \bullet_\eta P_2) = \Phi(A) \bullet_\eta \Phi(P_2).$$

Add Equations (2.9) and (2.10) together, to get

$$\Phi(A \bullet_\eta P_1) + \Phi(A \bullet_\eta P_2) = \Phi(A) \bullet_\eta (\Phi(P_1) + \Phi(P_2)).$$

By Claims 2.4, 2.8 and 2.10, we obtain

$$\Phi(A \bullet_\eta P_1 + A \bullet_\eta P_2) = \Phi(A) \bullet_\eta \Phi(P_1 + P_2),$$

Equivalently,

$$\Phi(A \bullet_\eta (P_1 + P_2)) = \Phi(A) \bullet_\eta \Phi(P_1 + P_2),$$

so, $\Phi(A \bullet_\eta I) = \Phi(A) \bullet_\eta \Phi(I)$. \square

Lemma 2.12. *Let Φ satisfy the assumptions of the Main Theorem. Then $\Phi(P_1)$ and $\Phi(P_2)$ are nontrivial orthogonal projections in \mathcal{B} .*

Proof. Let $P \in \{P_1, P_2\}$, where P_i for $1 \leq i \leq 2$ are nontrivial projections in \mathcal{A} . By Lemma 2.11 and the definition of the Jordan η_* -product we have

$$\begin{aligned} \Phi(P \bullet_\eta I) &= \Phi(P) \bullet_\eta \Phi(I) \\ \Phi(I \bullet_\eta P) &= \Phi(I) \bullet_\eta \Phi(P). \end{aligned}$$

Since Φ is unital, the above equations give us

$$(2.11) \quad \begin{aligned} \Phi(P + \eta P) &= \Phi(P) + \eta\Phi(P)^* \\ \Phi(P + \eta P) &= \Phi(P) + \eta\Phi(P), \end{aligned}$$

so we obtain $\Phi(P) = \Phi(P)^*$. On the other hand, $\Phi(P \bullet_\eta P) = \Phi(P) \bullet_\eta \Phi(P)$. Then $\Phi(P + \eta P) = \Phi(P)^2 + \eta\Phi(P)^2$. It follows from (2.11) that $\Phi(P) + \eta\Phi(P) = \Phi(P)^2 + \eta\Phi(P)^2$. Now, Lemma 2.2 implies $\Phi(P)^2 = \Phi(P)$. We show that $\Phi(P_1)$ and $\Phi(P_2)$ are orthogonal. Let $\Phi(P_1) = Q_1$ and $\Phi(P_2) = Q_2$. Then, by Claim 2.4, $Q_1 + Q_2 = I$. Also, $Q_1 \cdot Q_2 = \Phi(P_1) \cdot \Phi(P_2) = \Phi(P_1) \cdot (\Phi(I) - \Phi(P_1)) = 0$. \square

Lemma 2.13. *Let Φ satisfy the assumptions of the Main Theorem. Then*

$$(2.12) \quad \Phi(AP_i) = \Phi(A)\Phi(P_i),$$

for $1 \leq i \leq 2$.

Proof. It is easy to check that $\Phi(AP_i \bullet_\eta I) = \Phi(A \bullet_\eta P_i)$. The above equation can be written $\Phi(AP_i) + \eta\Phi(AP_i)^* = \Phi(A)\Phi(P_i) + \eta\Phi(P_i)\Phi(A)^*$. Equivalently,

$$(\Phi(AP_i) - \Phi(A)\Phi(P_i)) + \eta(\Phi(AP_i) - \Phi(A)\Phi(P_i))^* = 0,$$

Applying Lemma 2.2, we get $\Phi(AP_i) = \Phi(A)\Phi(P_i)$. \square

Now, lemma 2.12 ensures that there exist nontrivial projections Q_i ($i = 1, 2$) such that $\Phi(P_i) = Q_i$ and $Q_1 + Q_2 = I$. We can write $\mathcal{B} = \sum_{i,j=1}^2 \mathcal{B}_{ij}$ where $\mathcal{B}_{ij} = Q_i \mathcal{B} Q_j$, $i, j = 1, 2$.

The primeness property of \mathcal{B} is only used in the following claim.

Claim 2.14. *For every $A_{ii}, B_{ii} \in \mathcal{A}_{ii}$, $1 \leq i \leq 2$, we have*

$$\Phi(A_{ii} + B_{ii}) = \Phi(A_{ii}) + \Phi(B_{ii}).$$

First, we will prove that $\Phi(P_i A + P_i B) = \Phi(P_i A) + \Phi(P_i B)$ for every $A, B \in \mathcal{A}$.

By Lemma 2.13, Claim 2.8 and for every $T_{ji} \in \mathcal{B}_{ji}$ such that $i \neq j$ we obtain

$$\begin{aligned} (\Phi(P_i A + P_i B) - \Phi(P_i A) - \Phi(P_i B)) T_{ji} &= (\Phi(P_i A + P_i B) - \Phi(P_i A) \\ &\quad - \Phi(P_i B)) Q_j T Q_i \\ &= (\Phi(P_i A + P_i B) Q_j - \Phi(P_i A) Q_j \\ &\quad - \Phi(P_i B) Q_j) T Q_i \\ &= (\Phi(P_i A P_j + P_i B P_j) - \Phi(P_i A P_j) \\ &\quad - \Phi(P_i B P_j)) T Q_i \\ &= (\Phi(A_{ij} + B_{ij}) - \Phi(A_{ij}) \\ &\quad - \Phi(B_{ij})) T Q_i \\ &= (\Phi(A_{ij}) + \Phi(B_{ij}) - \Phi(A_{ij}) \\ &\quad - \Phi(B_{ij})) T Q_i \\ &= 0. \end{aligned}$$

By the primeness of \mathcal{B} , we have

$$(2.13) \quad \Phi(P_i A + P_i B) = \Phi(P_i A) + \Phi(P_i A).$$

Now, multiply the right side of equation (2.13) by $\Phi(P_i) = Q_i$ and use Lemma 2.13 to obtain

$$\Phi(P_i A P_i + P_i B P_i) = \Phi(P_i A P_i) + \Phi(P_i B P_i).$$

So, additivity of Φ follows from Claims 2.8, 2.10 and 2.14. It remains to prove that Φ is $*$ -preserving for non-zero rational numbers η such that $|\eta| \neq 1$. Since Φ is Jordan η_* -product preserving, we have

$$\Phi(A \bullet_\eta I) = \Phi(A) \bullet_\eta \Phi(I).$$

Since Φ is unital,

$$\Phi(A + \eta A^*) = \Phi(A) + \eta \Phi(A)^*.$$

Additivity of Φ and the above equation imply

$$(2.14) \quad \Phi(\eta A^*) = \eta \Phi(A)^*.$$

Let $\eta = \frac{a}{b}$, where a, b are integers. It is easy to see that

$$(2.15) \quad \Phi\left(\frac{1}{b} A^*\right) = \frac{1}{b} \Phi(A^*).$$

Now, by additivity of Φ and (2.15), we have $\Phi\left(\frac{a}{b} A^*\right) = \frac{a}{b} \Phi(A^*)$. It follows that $\Phi(\eta A^*) = \eta \Phi(A^*)$. Hence, by the latter equation and equation (2.14), we get $\Phi(A^*) = \Phi(A)^*$. □

REFERENCES

- [1] Z. F. Bai and S. P. Du, Multiplicative Lie isomorphism between prime rings, *Comm. Algebra* **36** (2008), no. 5, 1626–1633.
- [2] Z. F. Bai and S. P. Du, Multiplicative $*$ -Lie isomorphism between factors, *J. Math. Anal. Appl.* **346** (2008), no. 1, 327–335.
- [3] K. I. Beidar, M. Brešar, M. A. Chebotar and W. S. Martindale, On Herstein's Lie map conjectures, I, *Trans. Amer. Math. Soc.* **353** (2001), no. 10, 4235–4260.
- [4] K. I. Beidar, M. Brešar, M. A. Chebotar and W. S. Martindale, On Herstein's Lie map conjectures, II, *J. Algebra* **238** (2001), no. 1, 239–264.
- [5] K. I. Beidar, M. Brešar, M. A. Chebotar and W. S. Martindale, On Herstein's Lie map conjectures, III, *J. Algebra* **238** (2002), no. 1, 59–94.
- [6] M. Brešar, Jordan mappings of semiprime rings, *J. Algebra* **127** (1989), no. 1, 218–228.
- [7] M. Brešar, A. Fošner, On ring with involution equipped with some new product, *Publ. Math. Debrecen* **57** (2000), no. 1-2, 121–134.
- [8] J. Cui, C. K. Li, Maps preserving product $XY - YX^*$ on factor von Neumann algebras, *Linear Algebra Appl.* **431** (2009), no. 5-7, 833–842.
- [9] L. Dai and F. Lu, Nonlinear maps preserving Jordan $*$ -products, *J. Math. Anal. Appl.* **409** (2014), no. 1, 180–188.
- [10] J. Hakeda, Additivity of Jordan $*$ -maps on AW $*$ -algebras, *Proc. Amer. Math. Soc.* **96** (1986), no. 3, 413–420.
- [11] I. N. Herstein, Jordan homomorphisms, *Trans. Amer. Math. Soc.* **81** (1956) 331–341.

- [12] N. Jacobson and C. E. Rickart, Jordan homomorphism of rings, *Trans. Amer. Math. Soc.* **69** (1950) 479–502.
- [13] P. Ji and Z. Liu, Additivity of Jordan maps on standard Jordan operator algebras, *Linear Algebra Appl.* **430** (2009), no. 1, 335–343.
- [14] R. V. Kadison, Isometries of operator algebras, *Ann. of Math (2)*. **54** (1951), 325–338.
- [15] C. Li, F. Lu and X. Fang, Nonlinear mappings preserving product $XY + YX^*$ on factor von Neumann algebras, *Linear Algebra Appl.* **438** (2013), no. 5, 2339–2345.
- [16] F. Lu, Additivity of Jordan maps on standard operator algebras, *Linear Algebra Appl.* **357** (2002) 123–131.
- [17] F. Lu, Jordan maps on associative algebras, *Comm. Algebra* **31** (2003), no. 5, 2273–2286.
- [18] F. Lu, Jordan triple maps, *Linear Algebra Appl.* **375** (2003) 311–317.
- [19] L. W. Marcoux and Ahmed R. Sourour, Lie isomorphism of nest algebras, *J. Funct. Anal.* **164** (1999), no. 1, 163–180.
- [20] W. S. Martindale III, When are multiplicative mappings additive?, *Proc. Amer. Math. Soc.* **21** (1969) 695–698.
- [21] C. R. Mires, Lie isomorphisms of operator algebras, *Pacific J. Math.* **38** (1971), no. 3, 717–735.
- [22] C. R. Mires, Lie isomorphisms of factors, *Trans. Amer. Math. Soc.* **147** (1970) 55–63.
- [23] L. Molnár, A condition for a subspace of $B(H)$ to be an ideal, *Linear Algebra Appl.* **235** (1996) 229–234.
- [24] L. Molnár, On isomorphisms of standard operator algebras, *Studia Math.* **142** (2000), no. 3, 295–302.
- [25] X. Qi and J. Hou, Additivity of Lie multiplicative maps on triangular algebras, *Linear Multilinear Algebra* **59** (2011), no. 4 391–397.
- [26] P. Šemrl, Quadratic functionals and Jordan $*$ -derivations, *Studia Math.* **97** (1991), no. 3, 157–165.
- [27] A. Taghavi, V. Darvish and H. Rohi, Additivity of maps preserving products $AP \pm PA^*$, *Mathematica Slovaca*, Accepted.

(Ali Taghavi) DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES,
UNIVERSITY OF MAZANDARAN, P.O. BOX 47416-1468, BABOLSAR, IRAN.
E-mail address: taghavi@umz.ac.ir

(Hamid Rohi) DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES,
UNIVERSITY OF MAZANDARAN, P.O. BOX 47416-1468, BABOLSAR, IRAN.
E-mail address: h.rohi@stu.umz.ac.ir

(Vahid Darvish) DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICAL SCIENCES,
UNIVERSITY OF MAZANDARAN, P.O. BOX 47416-1468, BABOLSAR, IRAN.
E-mail address: v.darvish@stu.umz.ac.ir