

APPLICATIONS OF EPI-RETRACTABLE MODULES

B. M. PANDEYA, A. K. CHATURVEDI* AND A. J. GUPTA

Communicated by Siamak Yassemi

ABSTRACT. An R -module M is called *epi-retractable* if every submodule of M_R is a homomorphic image of M . It is shown that if R is a right perfect ring, then every projective slightly compressible module M_R is epi-retractable. If R is a Noetherian ring, then every epi-retractable right R -module has direct sum of uniform submodules. If endomorphism ring of a module M_R is von-Neumann regular, then M is semi-simple if and only if M is epi-retractable. If R is a quasi Frobenius ring, then R is a right hereditary ring if and only if every injective right R -module is semi-simple. A ring R is semi-simple if and only if R is right hereditary and every epi-retractable right R -module is projective. Moreover, a ring R is semi-simple if and only if R is pri and von-Neumann regular.

1. Introduction

All rings are associative with unit elements and all modules are unitary right modules. Let R be a ring. The ring R is said to be a principal right ideal (pri) ring if every right ideal of R is principal. Ghorbani and Vedadi [3] generalized this concept to modules. An R -module M is called epi-retractable if every submodule of M_R is a homomorphic image of M . Therefore, R is a pri ring if and only if R_R is epi-retractable. An

MSC(2010): Primary: 16D10; Secondary: 16D40, 16L30, 16P40, 16E60.

Keywords: Epi-retractable modules, semi-simple rings, perfect rings, hereditary rings, von-Neumann regular rings.

Received: 17 September 2010, Accepted: 19 February 2011.

*Corresponding author

© 2012 Iranian Mathematical Society.

R -module N is called M -cyclic if it is isomorphic to M/L , for some submodule L of M (see [10]). Note that M_R is epi-retractable if and only if every submodule of M is M -cyclic. Here, we shall investigate epi-retractable modules in terms of M -cyclic submodules and also provide those properties of epi-retractable modules which have not been studied earlier.

By [2, 6.9.3], an R -module M is called *compressible* if for every non-zero submodule N of M there exists a monomorphism from M to N . The concept of epi-retractable modules is dual to the concept of compressible modules. There exist some epi-retractable modules which are not compressible. For example, semi-simple modules are epi-retractable but not compressible.

In Section 2, we study two important properties of epi-retractable modules. We observe that every epi-retractable module is a *slightly compressible module* (see [6]), but the converse need not be true. In Theorem 2.2, we provide a sufficient condition for slightly compressible modules to be epi-retractable. We show that if R is a right perfect ring, then every projective slightly compressible module M_R is epi-retractable. This is a well known problem in the theory of rings and modules when a module has direct sum of uniform submodules. In Theorem 2.3, we show that if R is a Noetherian ring, then every epi-retractable right R -module has direct sum of uniform submodules.

In Section 3, we study the semi-simplicity of epi-retractable modules and pri rings. Note that every semi-simple module is epi-retractable, but the converse need not be true. In some results of that section, we provide sufficient conditions for the epi-retractable modules to be semi-simple by injective modules, projective modules, right hereditary rings, von-Neumann regular rings. We show that if endomorphism ring of a module M is von-Neumann regular, then M is semi-simple if and only if M is an epi-retractable module. If R is a quasi Frobenius ring, then R is a right hereditary ring if and only if every injective R -module is semi-simple. We characterize semi-simple rings by epi-retractable modules so that a ring R is semi-simple if and only if R is right hereditary and every epi-retractable R -module is projective. We end up with a result that states: A ring R is semi-simple if and only if R is pri and von-Neumann regular.

We refer to [10] and [1] for all undefined notions used in the text.

2. Epi-retractable modules

Let $R = \begin{pmatrix} F & F \\ 0 & F \end{pmatrix}$, where F is a ring. Then, $M_R = \begin{pmatrix} F & F \\ 0 & 0 \end{pmatrix}$, $N_R = \begin{pmatrix} 0 & 0 \\ 0 & F \end{pmatrix}$ and $P_R = \begin{pmatrix} 0 & F \\ 0 & 0 \end{pmatrix}$ are right R -modules. It is clear that R_R , N_R , P_R and $(M/P)_R$ are epi-retractable R -modules. But, M_R is not an epi-retractable module. Moreover, submodules of an epi-retractable module need not be epi-retractable and also factors of an epi-retractable module need not be epi-retractable. We begin with the observation that the class of epi-retractable modules is closed under direct sums.

Proposition 2.1. *Let $\{M_i\}_{i \in I}$ be a family of epi-retractable modules. Then, $M = \bigoplus_{i \in I} M_i$ is an epi-retractable module.*

Proof. Let K be a submodule of M . Then, $K \cap M_i$ is a submodule of M_i , for each $i \in I$. Since each M_i is an epi-retractable module, there exists an epimorphism $\alpha_i : M_i \rightarrow K \cap M_i$. Define $\alpha = \sum_{i \in I} \alpha_i : M \rightarrow K$. Then, clearly α is a surjective homomorphism. Hence, $\bigoplus_{i \in I} M_i$ is an epi-retractable module. \square

A projective R -module P together with a small epimorphism $\pi : P \rightarrow M$ is called a *projective cover* of M . A ring R is said to be *right perfect* if every R -module has a projective cover. In [6], Smith calls an R -module M *slightly compressible* if, for every non-zero submodule N of M , there exists a non-zero homomorphism from M to N . An R -module M is called to be *self-generator* if, for each submodule N of M , there exists an index set J and an epimorphism $\theta : M^{(J)} \rightarrow N$. It is clear that every epi-retractable module is self-generator. Moreover, every self-generator is slightly compressible. Then, epi-retractable modules are slightly compressible. In general, every slightly compressible module is not a self-generator (see [6, Proposition 3.1]). Therefore, every slightly compressible module need not be epi-retractable.

The following result shows a sufficient condition for slightly compressible modules to be epi-retractable.

Theorem 2.2. *Let R be a right perfect ring. Then, every projective slightly compressible R -module is epi-retractable.*

Proof. Assume that M is a projective and slightly compressible module. Let K be a submodule of M . Since R is right perfect, there is a projective cover P of K with a small $\text{Ker}(\pi)$, where $\pi : P \rightarrow K$ is an epimorphism.

Then, there exists a non-zero homomorphism $f : M \rightarrow K$. Consider the following diagram:

$$\begin{array}{ccc} & M & \\ h \swarrow & & \downarrow f \\ P & \longrightarrow & K \\ & \pi & \end{array}$$

Since M is projective, f can be lifted to a homomorphism h from M to P such that the above diagram is commutative, that is, $f = \pi h$. It follows that $P = \text{Im}(h) + \text{Ker}(\pi)$. Then, $P = \text{Im}(h)$, because $\text{Ker}(\pi)$ is small. This implies that h is surjective. Therefore, f is also surjective, and hence M is an epi-retractable module. \square

A ring R is called a *right V-ring* if every simple R -module is injective. Moreover, if R is a right V -ring, then every projective R -module is slightly compressible (see [6, Theorem 1.5]). Theorem 2.2 has the following consequence.

Corollary 2.3. *Let R be a right perfect and right V -ring. Then, every projective R -module is epi-retractable.*

Proof. This follows from [6, Theorem 1.5] and Theorem 2.2. \square

Following [9], an R -module M is called *quasi-polysimple* if every non-zero submodule of M contains a uniform submodule of M . Note that over a Noetherian ring R , every R -module is quasi-polysimple (see [5, Theorem 2.2]).

We shall now investigate when a epi-retractable module has direct sum of uniform submodules.

Theorem 2.4. *Let R be a Noetherian ring. If M is an epi-retractable R -module, then M has direct sum of uniform submodules of M .*

Proof. It is clear that M is quasi-polysimple. Therefore, M is an essential extension of the direct sum $\bigoplus_{i \in J} K_i$, where each K_i is the uniform submodule of M and J is some index set (see [5, Lemma 2.1]). Since M is epi-retractable, there exists an endomorphism $f \in S$ such that $f(M) = \bigoplus_{i \in J} K_i$. \square

3. Semi-simplicity of epi-retractable modules

A ring R is called *right hereditary* if every right ideal is projective. Moreover, R is right hereditary if and only if every submodule of every

projective R -module is projective and if and only if quotients of injective, R -modules are injective (see [4, Corollary 2.26] and [4, Theorem 3.22]). There are some modules which are injective, but not epi-retractable. For example, the set of rational numbers Q_Z is an injective module, but is not epi-retractable. Note that every semi-simple module is epi-retractable, but in general the converse is not true.

In the following, we investigate when an epi-retractable module is semi-simple.

Proposition 3.1. *Let R be a right hereditary ring. Then, the followings hold:*

- (1) *Every injective epi-retractable R -module is semi-simple.*
- (2) *Every projective epi-retractable R -module is semi-simple.*

Proof. (1). Assume that R is a right hereditary ring and K is submodule of an epi-retractable injective R -module M . Since M is epi-retractable, $K \cong M/L$, for some submodule L of M . It follows that K is injective. Suppose I is the identity map from K to K . Therefore, I can be extended to a homomorphism from M to K . Hence, K is a direct summand of M . This implies that M is semi-simple.

(2). This is clear. □

Recall that a ring R is said to be a quasi Frobenius ring if it is a (left) right self injective Noetherian ring. Note that if R is a ring such that every injective R -module is epi-retractable, then R is a quasi Frobenius ring (see [3, Proposition 3.2]). In the following, we characterize right hereditary rings.

Proposition 3.2. *Let R be a quasi Frobenius ring. Then, R is a right hereditary ring if and only if every injective R -module is semi-simple.*

Proof. Assume R is a right hereditary ring. Let M be an injective R -module. By [3, Proposition 3.2], M is an epi-retractable module. By Proposition 3.1, it is clear that M is a semi-simple module.

Conversely, assume that every injective R -module is semi-simple. Suppose that K is the homomorphic image of an injective R -module M . Then, K is a direct summand of M , because M is semi-simple. Therefore, K is also injective. This implies that quotients of injective R -modules are injective. This proves that R is a right hereditary ring. □

Theorem 3.3. *If the endomorphism ring S of a module M is von-Neumann regular, then M is semi-simple if and only if M is an epi-retractable module.*

Proof. Suppose M is an epi-retractable module and K is a submodule of M . Then, there is an epimorphism f from M to K . Since $S = \text{End}(M)$ is von-Neumann regular, $f(M) = K$ is a direct summand of M . Hence, M is a semi-simple module. The converse is obvious. \square

Let R be a ring and M be an R -module. We denote $r(x) = \{s \in R : xs = 0\}$, for some $x \in M$. Note that $r(x)$ is a right ideal of R and $R/r(x) \cong xR$, for all $x \in M$. In the following, we characterize semi-simple ring.

Theorem 3.4. *A ring R is semi-simple if and only if R is right hereditary and every epi-retractable R -module is projective.*

Proof. Assume that R is a right hereditary ring and every epi-retractable R -module is projective. Let M be a simple R -module. It follows that M is epi-retractable and projective. For any $x \in M$, $xR \cong R/r(x)$. Then, xR (and hence $R/r(x)$) is projective, because R is a right hereditary ring. Therefore, the exact sequence $0 \rightarrow r(x) \rightarrow R \rightarrow R/r(x) \rightarrow 0$ splits. This implies that $r(x)$ is a direct summand of M . Since $r(x)$ is a maximal right ideal, R is a semi-simple ring. The converse is obvious. \square

An R -module M is said to satisfy *(**)-property* if every non-zero endomorphism of M is an epimorphism (see [11]). In general, epi-retractable modules do not satisfy *(**)-property*. For example, Z as Z -module is epi-retractable, but it does not satisfy *(**)-property*. The following result shows that epi-retractable module with *(**)-property* is simple.

Proposition 3.5. *An R -module M is simple if and only if M is epi-retractable with *(**)-property*.*

Proof. Assume that M is epi-retractable with *(**)-property*. Let K be a proper submodule of M . Then, there is an epimorphism $f : M \rightarrow K$. This implies that f is a non-zero endomorphism from M to M . Since M satisfies *(**)-property*, $f(M) = M = K$. Hence, M is simple. The converse is obvious. \square

Corollary 3.6. *If an R -module M is epi-retractable with *(**)-property*, then $\text{End}(M_R)$ is a division ring.*

An R -module M is said to satisfy *(*)-property* if every non-zero endomorphism of M is a monomorphism (see [7]). This is dual to the concept of *(**)-property* defined earlier.

Proposition 3.7. *Every epi-retractable module with *(*)-property* is a co-Hopfian module.*

Proof. Straightforward. \square

Theorem 3.8. *A ring R is semi-simple if and only if R is a pri and von-Neumann regular ring.*

Proof. Assume that R is a pri ring. Then, every right ideal of ring R is a principal right ideal. This implies that every right ideal is a direct summand of R , because R is von-Neumann regular. It follows by [10, 20.7] that R is a semi-simple ring. \square

Proposition 3.9. *Let R be a ring such that every slightly compressible R -module is pseudo-projective. Then, R is a right V -ring if and only if R is a semi-simple ring.*

Proof. Let M be a slightly compressible R -module. Suppose there is a free R -module F with an epimorphism $g : F \rightarrow M$. By [6, Theorem 1.5], F is a slightly compressible module. Then, $F \oplus M$ is a slightly compressible module by [6, Proposition 1.4]. Consider the exact sequence $0 \rightarrow \text{Ker}(g) \xrightarrow{i} F \xrightarrow{g} M \rightarrow 0$. This sequence splits by [8, Lemma 1.3]. Therefore, M is a direct summand of F . Hence, M is projective. In particular, every simple R -module is projective. It follows by [10, 20.7] that R is a semi-simple ring. \square

Corollary 3.10. *Over a right V -ring R , if every slightly compressible R -module is pseudo-projective, then every R -module is epi-retractable.*

A ring R is called *right semi-artinian* if every non-zero R -module has non-zero socle.

Proposition 3.11. *Let R be a right semi-artinian right V -ring. Then, R is semi-simple if and only if every R -module is pseudo-projective.*

Proof. Assume that over a right semi-artinian right V -ring R , every R -module is pseudo-projective. By [6, Proposition 1.18], every right R -module is slightly compressible. It follows by Proposition 3.9 that R is semi-simple. \square

Corollary 3.12. *Over a right semi-artinian right V -ring, every pseudo-projective module is an epi-retractable module.*

Recall that a ring R is *right PP-ring* if every cyclic right ideal of R is projective. A ring R is called a *regular* if for any $a \in R$ there is an element $b \in R$ with $aba = a$. Note that R is regular if and only if every right principal ideal is a direct summand in R (see [10, 3.10]).

Proposition 3.13. *The followings are equivalent for a pri ring R .*

- (1) R is a right PP-ring.
- (2) R is a right hereditary ring.
- (3) R is a von-Neumann regular ring.

Proof. (1) \Rightarrow (2). Straightforward.

(2) \Rightarrow (3). Assume the condition (2). Let L be a principal right ideal of R . Then, L is projective, because R is a right hereditary ring. Suppose $\pi : R \rightarrow L$ is an epimorphism and $I : L \rightarrow L$ is the identity map. This implies that I can be lifted to a homomorphism f from L to R , that is, $I = \pi f$. It follows that L is a direct summand of R . Hence, R is a von-Neumann regular ring.

(3) \Rightarrow (1). Obvious. □

Acknowledgments

The authors thank the referee for his/her careful considerations.

REFERENCES

- [1] F. W. Anderson and K. R. Fuller, Rings and Categories of Modules, Second edition. Graduate Texts in Mathematics, 13. Springer-Verlag, New York, 1992.
- [2] J. C. Mc Connell, J. C. Robson, Non-commutative Noetherian Rings, Wiley-Interscience, New York, 1987.
- [3] A. Ghorbani and M. R. Vedadi, Epi-retractable modules and some applications, *Bull. Iranian Math. Soc.* **35** (2009), no. 1, 155–166.
- [4] T. Y. Lam, Lectures on Modules and Rings, Graduate Texts in Mathematics, 189. Springer-Verlag, New York, 1999.
- [5] B. M. Pandeya and S. A. Paramhans, Extending Modules and Modules over Noetherian Rings, *J. Indian Math. Soc. (N.S.)* **62** (1996), no. 1-4, 121–128.
- [6] P. F. Smith, Modules with many homomorphisms, *J. Pure Appl. Algebra* **197** (2005), no. 1-3, 305–321.
- [7] A. K. Tiwary and B. M. Pandeya, Modules whose non zero endomorphisms are monomorphisms, in: Proc. Int. Symp. Algebra and its Applications, New Delhi, 1981, in: *Lect. Notes Pure Appl. Math.* **91** Marcel Dekker, Inc., New York, 1984.
- [8] A. K. Tiwary and B. M. Pandeya, Pseudo-projective and Pseudo-injective Modules, *Indian J. Pure Appl. Math.* **9** (1978), no. 9, 941–949.
- [9] A. K. Tiwary and B. M. Pandeya, On poly simple modules, *Progr. Math. (Allahabad)* **5** (1971), no. 1-2, 49–61.
- [10] R. Wisbauer, Foundations of Module and Ring Theory, A handbook for study and research. Revised and translated from the 1988 German edition. Algebra, Logic and Applications, 3. Gordon and Breach Science Publishers, Philadelphia, PA, 1991
- [11] W. Xue, On morita duality, *Bull. Austral. Math. Soc.* **49** (1994), no. 1, 35–45.

Bashishth Muni Pandeya

Department of Applied Mathematics, Institute of Technology, Banaras Hindu University, P.O. Box 221005, Varanasi, India

Email: bmpandeya@bhu.ac.in; bmpandeya@gmail.com

Avanish Kumar Chaturvedi

Department of Mathematics, Jaypee Institute of Information Technology, P.O. Box 201307, Noida(UP), India

Email: akc99@rediffmail.com; akchaturvedi.math@gmail.com

Ashok Ji Gupta

Department of Applied Mathematics, Institute of Technology, Banaras Hindu University, P.O. Box 221005, Varanasi, India

Email: ashokg@bhu.ac.in